

Memo 04: In-Run Mass-Conservation Checkpoints for the PLAST_FLOC Split Path

FVCOM-MP Delaware Test Cases

May 4, 2026

Purpose

Memo 03 (v006) established that the ~ 650 kg mass gap between case **a** and cases **b/c** over ten days is driven entirely by the `PLAST_FLOC=T` split path: with `SOURCE_AG_RATE=0` and zero bed mass throughout, the aggregated component remains zero and all water-column plastic is in `conc_d`, yet the domain total oscillates with 28 sign changes after day 3 while the no-floc case **a** grows monotonically.

The offline MATLAB budget confirms that the saved NetCDF output is internally consistent at each output snapshot. Therefore the mass leakage occurs somewhere inside `Advance_Plast` between two consecutive output records, and is not visible in the NetCDF files.

This memo documents the strategy for inserting *in-run mass checkpoints* directly into `mod_plast.F` so that the guilty operator stage can be identified without additional NetCDF output.

Strategy: Dedicated CSV File

Writing diagnostic totals to `write(*,...)` is impractical because the FVCOM screen log intermingles output from every module on every timestep. Instead, a dedicated file is opened once during `Setup_PLAST` and written exclusively by the master process (`msr`). The file is named

`<casename>_mp_mass_check.csv`

and is placed in the run input directory alongside the existing plastic input files. The overhead is gated behind a single namelist key:

`PLAST_MASS_CHECK = T`

When this key is absent or `F`, the new code has zero runtime cost: all checkpoint calls return immediately after the first guard test.

Checkpoint rows are written at the same cadence as the existing `PLAST_Report` screen statistics (every `n_report` internal steps), keeping file sizes manageable while still providing sub-output-interval resolution.

Formula Verification Against MATLAB Offline Budget

The MATLAB script `extract_mp1_mass_budget_a.m` computes:

$$\begin{aligned}
 V_{i,k} &= \text{art1}(i) \max(h_i + \zeta_i, 0) \Delta\sigma_{i,k} \\
 M_{\text{water}} &= \sum_{i,k} \text{mp1}(i,k) \cdot V_{i,k} \quad (\text{wet nodes only}) \\
 M_{\text{bed}} &= \sum_i \text{bot_mass_mp1}(i) \cdot \text{art1}(i) \quad (\text{all nodes}) \\
 M_{\text{total}} &= M_{\text{water}} + M_{\text{bed}}
 \end{aligned}$$

where `art1` is the node-based control-volume area, $\Delta\sigma_{i,k}$ is the sigma-layer thickness fraction (`dzfrac` in MATLAB), and $h_i + \zeta_i$ is the total water depth.

The Fortran in-run equivalent uses identically named arrays:

$$\begin{aligned}
 M_{\text{water}} &= \sum_{i=1}^m \sum_{k=1}^{k_{bm1}} c(i,k) \cdot \text{ART1}(i) \cdot D(i) \cdot \text{DZ}(i,k) \\
 M_{\text{bed}} &= \sum_{i=1}^m \text{mass}(i) \cdot \text{ART1}(i)
 \end{aligned}$$

where $D(i) = H(i) + \text{EL}(i)$ is the FVCOM total depth and $\text{DZ}(i,k)$ is the sigma-layer fraction — exactly matching $\Delta\sigma_{i,k}$ from MATLAB. The bed sum uses *all nodes* to match MATLAB's `total_mass_all_bed_kg` conservation companion.

The formulas are equivalent. The only future guard required is a `#if WET_DRY` wet-node mask for domains with drying cells; the v006 Delaware domain has no dry nodes, so the Fortran loop over `i=1..m` is identical to the MATLAB wet-node filter.

For the split path, the subroutine computes separate sums for `conc_a` and `conc_d` (and `mass_a`, `mass_d`), writing all components plus combined totals in each CSV row so that per-component budgets can be tracked alongside the grand total.

Checkpoint Table

Thirteen checkpoints are inserted into the `floc_intract=T` path of `Advance_Plast`. Checkpoints CP0–CP3 are guarded by `#if defined (SEDIMENT)` and `if(floc_intract)` at the call site; CP4–CP12 are already inside the relevant conditional blocks.

The CSV row format is:

```
iint, T_days, checkpoint, iplast, wc_a_kg, wc_d_kg, wc_tot_kg, bed_a_kg, bed_d_kg,
      bed_tot_kg, grand.total_kg
```

Code Reconstruction in `mod_plast.F`

Four modification phases were implemented in `FVCOM_source_repo_github/mod_plast.F`.

1. **Module-level variables** (near line 490, after the `floc_intract` flag):

```
logical  :: PLAST_mass_check = .false.
integer, parameter :: mass_check_unit = 77
```

CP#	Label	Arrays passed	Expected conservation
0	CP0.start	conc_a,conc_d,mass_a,mass_d	Reference state at timestep entry
1	CP1.after.deposition	same	Water $\downarrow$ + bed $\uparrow$ = const
2	CP2.after.erosion	same	Water $\uparrow$ + bed $\downarrow$ = const
3	CP3.after.update.bottom	same	Bed bookkeeping; grand total = const
4	CP4.after.advection	cnew_a,cnew_d,mass_a,mass_d	Zero water-column change
5	CP5.after.vdif	same	Zero change
6	CP6.after.kinetics	same	cnew_a+cnew_d conserved
7	CP7.after.upper.clamp	same	Loss only if >100 cap fires
8	CP8.after.OBC	same	Change = OBC flux (open domain)
9	CP9.after.PTsource	same	Change = river source
10	CP10.after.neg.clamp	same	Loss only if negatives exist
11	CP11.after.MPI	same	Zero change
12	CP12.end	conc_a,conc_d,mass_a,mass_d	Must equal CP11

2. **Namelist read** (Read_PLAST_Params, immediately after the PLAST_FLOC read):

```
Call Get_Val(PLAST_mass_check, plastfile,
             'PLAST_MASS_CHECK', line=linenum, echo=.false.)
```

3. **CSV file open and header** (Setup_PLAST, after call report_PLAST_setup): Opens unit 77 to <input_dir>/<casename>_mp_mass_check.csv and writes the column header on the master process only.
4. **Subroutine Write_Mass_Checkpoint** (new subroutine added at the end of the module, before End Module Mod_Plast): Accepts one plastic class at a time via arguments (label, iplast_idx, c_a, c_d, m_a, m_d), computes local sums using the $ART1 \cdot D \cdot DZ$ formula, reduces via `MPI_REDUCE(MPI_SUM)`, and writes one CSV row per class if `msr=.true.` The routine returns immediately if `PLAST_mass_check=F` or if the current iteration is not a reporting step.
5. **Thirteen call sites in Advance_Plast**: CP0–CP3 wrapped in `#if defined (SEDIMENT)` / `if(floc_intract)` guards; CP4–CP12 placed inside the existing `floc_intract` or per-stage conditional blocks.

Findings: Root Causes and Fixing Suggestions

The checkpoint infrastructure was compiled and the model was run for ~ 4 days with `PLAST_MASS_CHECK = T` for cases b and c. The analysis script `analyze_mp_mass_checkpoints.py` produced the stage-mass-change heatmap shown in Figure 2, together with the NaN diagnostic tables.

Root Cause 1 — Structural issues in the split advection path

Diagnostic interpretation: the `adv*` heatmap signal is the expected open-boundary flux, not a conservation bug within the advection operator. The non-zero CP3→CP4 signal in the heatmap oscillates at the tidal period and is present in both cases. Reading `Adv_Scal` in `mod_scal.F` confirms the mechanism: for open-boundary nodes (`isonb= 2`), `Adv_Scal` replaces




Figure 1: Stage mass-change heatmap for cases b (left) and c (right). Each row is one operator stage; each column is one reporting timestep. Red = mass gained, blue = mass lost; the colour scale is clipped at $\pm 1\%$ of the domain total. Dashed horizontal lines mark the five stages that must be exactly mass-conservative (adv^* , vdif^* , kin^* , MPI^* , recomb^*). The adv^* row is the only conservative stage showing non-zero values. Subsequent analysis (Section) confirms that this signal is the expected open-boundary advective exchange flux present in all cases, not a split-path conservation bug.

`xflux` with only the vertical flux, while the horizontal advective flux from those nodes is still accumulated in the adjacent interior nodes. The global mass sum therefore changes by the net horizontal advective exchange across all OBC edges — the expected open-boundary flux. This is also present in case **a** and is not a split-path defect.

Why the multi-class loop is mass-conservative but the flocc split is not (for general parameters). The multi-class loop calls `adv_scal(plast(i)%conc, plast(i)%cnew, ..., plast(i)%cflx, ...)` for each class `i` in sequence. Because each class has its own separate `cflx` storage, no mutable state is shared between calls. A common misconception (including an earlier version of this memo) was that the module-level arrays `PFPXB/PFPYB` are shared between calls and could corrupt successive invocations. Inspection of `mod_scal.F` shows that both arrays are only *written* inside `Adv_Scal` (`pfpxb(i) = pfp(x(i)) at k=kbm1` as a side-effect output for the T/S vertical diffusion in `vdif_ts.F`) and are *never read* within `Adv_Scal` itself. There is therefore no gradient-contamination between successive calls.

`cflx` is concentration-nudge-only: the shared-array overwrite has zero effect on the simulation. Tracing the full call chain of `bcond_scal_OBC` in `mod_scal.F` reveals that `cflx` (`= fflux_obc`) has no effect in either branch of the OBC condition:

- **Outflow branch** (`uard_obcn > 0`): the advective flux is used to compute `f2d_obc`, but the line `fn(j,k) = f2d_obc + (fn(j1,k) - f2d_next)` is commented out in the source code. The live assignment is simply `fn(j,k) = fn(j1,k)`, making `cflx` irrelevant.
- **Inflow branch** (`uard_obcn ≤ 0`): pure concentration nudging `fn(j,k) = f(j,k) - alpha_nudge*(f(j,k) - f_obc(i))` — `cflx` is never referenced.

As a result, the fact that both `adv_scal` calls overwrite the same `plast(iplast)%cflx` array has no effect on the computed concentrations in the current code. The OBC treatment is entirely concentration-based. The earlier proposed Fix A (revised local `cflx_a` array) is therefore **withdrawn** — it would compile correctly but change nothing.

Flux-limiter asymmetry — future issue when `conc_a > 0`. `Adv_Scal` calls the non-linear subroutine `limit_hor_flux_Scal`, which caps each outflow edge flux at $f_{\text{source}} \cdot A_{cv} \cdot D/\Delta t$ (the total available tracer mass at the source node). When applied to `conc_a` and `conc_d` independently, the combined flux cap is $(c_a + c_d) \cdot A_{cv} \cdot D/\Delta t$ — the same as for the combined field. However, if the raw flux exceeds the individual cap for one component but not the other, the two separate calls can permit more total outflow than a single combined call. With `conc_a = 0` (test case) the limiter for `conc_a` always evaluates to zero and the `conc_d` call is identical to the single-field case — no asymmetry. For future runs with non-zero aggregated concentration, this asymmetry will introduce a small but nonzero transport non-conservation.

Proposed Fix A (both versions withdrawn). An earlier draft proposed combining `conc_a+conc_d` for a single `adv_scal` call and redistributing by f_a . This is physically wrong: f_a evolves by its own advection–diffusion–reaction history and cannot be used as a post-hoc partition key. A subsequent revision proposed a local `cflx_a` array to preserve the `conc_a` OBC flux independently. This is also ineffective because `cflx` is irrelevant (see above). Both Fix A variants are withdrawn. No change to the advection block is needed for correctness under the current test-case parameters.

Fix D (future) — multi-species flux limiting. For general runs with `conc_a > 0`, the correct approach is to apply `limit_hor_flux_Scal` using the *total* ($c_a + c_d$) as the available mass and then partition the limited flux between the two components proportionally. This is deferred until `SOURCE_AG_RATE > 0` test cases are designed.

Root Cause 2 — NaN propagating from `eflx` into `conc_a/conc_d`

Observed symptom (Phase 2 run). The short diagnostic run (~ 1 day, cases `b` and `c`) shows 75–82% of reporting steps have NaN at CP2. NaN propagates identically through CP3, CP3p5, CP4–CP9, and is silently zeroed at CP10 (neg-clamp), confirming that mass is destroyed silently on every affected step.

Root cause. Inside `Calc_Erosion`, the erosion flux `eflx(i)` is computed via the call chain:

```
Call Calc_critial_shear_WS(...) ! rho_water -> tau_c_p
Call Calc_Erosion_Rate(...)    ! tau_c_p -> Erate
eflx(i) = DTplast * Erate(i)    ! Erate -> eflx
```

At nodes where `rho_water(i,kbm1)` is zero or NaN (e.g. tidal-flat nodes that are formally wet but have negligible water mass), `tau_c_p` and therefore `Erate` become NaN. Once `eflx = NaN`, the subsequent concentration update

```
conc(i,kbm1) = conc(i,kbm1) + eflx(i)/(d(i)*dz(i,kbm1))
```

writes NaN into `conc_a` and `conc_d` regardless of whether `d(i)` is zero or positive. A secondary depth-guard (`d(i) <= 0 cycle`) was added first (Fix B) but had no effect because the NaN originates in `eflx` itself, before any division.

Implemented Fix B+. An `isnan` sentinel is applied immediately after `eflx` and `pflx` are assigned, intercepting NaN from any upstream source before it can enter the concentration arrays:

```
plast(iplast)%eflx(i) = erosion_flux
plast(iplast)%pflx(i) = precip_flux
! Fix B+: guard against NaN from any upstream source
if (isnan(plast(iplast)%eflx(i)) .or. &
    plast(iplast)%eflx(i) < 0.0_sp) plast(iplast)%eflx(i) = 0.0_sp
if (isnan(plast(iplast)%pflx(i)) .or. &
    plast(iplast)%pflx(i) < 0.0_sp) plast(iplast)%pflx(i) = 0.0_sp
```

The depth guard (`d(i) <= 0 cycle`) is retained in the concentration update loop as a secondary defence.

Phase 2 verification (confirmed). After Fix B+ the NaN rate drops to **0/1338 (0.0%)** for case `b` and **0/1286 (0.0%)** for case `c` across all checkpoints CP2–CP12. Additionally, CP3→CP3p5 (`adv_a*delta`) is **exactly zero** for every step in both cases, confirming that the `conc_a=0` advection call contributes no mass change and the split-advection path is structurally correct. The CP3p5→CP4 (`adv_d*`) row shows the expected tidal-period OBC exchange signal (magnitude ~ 1 g peak), identical to the case `a` single-field behaviour.

Further Diagnostic Instrumentation (Phase 2)

Because the first checkpoint run did not definitively separate the `conc_a` and `conc_d` contributions to the advection stage, two additional diagnostics have been added to `mod_plast.F` and the analysis script before the next model run.

CP3p5 — mid-advection split checkpoint

A new checkpoint CP3p5_after_adv_a is inserted *inside* the do iplast loop, between the two adv_scal calls:

```
call adv_scal(conc_a, cnew_a, ...)      ! first call
call Write_Mass_Checkpoint('CP3p5_after_adv_a', iplast,
    cnew_a, conc_d, mass_a, mass_d)    ! conc_d still holds pre-advection value
call adv_scal(conc_d, cnew_d, ...)      ! second call
```

This splits the previous adv* heatmap row into two:

- **CP3→CP3p5 (adv_a*)**: mass change from advecting conc_a alone. With conc_a=0 in the test case, this delta must be exactly zero. Any non-zero value would implicate a side-effect of adv_scal on shared arrays beyond cflx and PFPXB/PFPYB.
- **CP3p5→CP4 (adv_d*)**: mass change from advecting conc_d alone. This should reproduce the same tidal-period OBC signal previously seen in the combined adv* row, confirming that the behaviour is identical to the case a single-field path.

CFLX_TRACE diagnostic prints — confirmed always zero by design

Three trace lines are written to the checkpoint CSV file at each reporting step, recording $\sum |cflx_{iplast}|$ at three moments: before adv_scal(conc_a), after adv_scal(conc_a), and after adv_scal(conc_d). These rows begin with CFLX_TRACE and are filtered out by the Python analysis script during loading.

Phase 2 result: all 5,340 CFLX_TRACE entries in both cases are exactly zero, from timestep 1 onward. This is correct and expected. Inside adv_scal, cflx (= fflux.obc) is assigned as:

```
fflux_obc(i,k) = xflux_adv(i1,k)      ! at OBC ghost-node boundary edge
```

For isonb==2 open-boundary nodes, adv_scal applies a vertical-flux-only treatment: the horizontal flux at the outward boundary edge is set to zero, so xflux_adv at that edge is zero and cflx = 0 unconditionally. The actual OBC mass exchange — the tidal-period signal visible in the adv* heatmap row — enters through the *interior edges* adjacent to OBC nodes. Those edge fluxes are included in the divergence sum of interior cells but are not stored in cflx. The plastic boundary condition (cdis) prescribes concentration directly; no flux is routed through cflx at any time.

This conclusively confirms that the shared-array overwrite of cflx between the two adv_scal calls has zero effect, and both Fix A variants remain withdrawn.

Adjustment Plan and Further Testing

1. Phase 2 diagnostics — complete.

- CFLX_TRACE: all zeros by design (confirmed). ✓
- Fix B (depth guard): compiled and tested — NaN persisted because the source is in eflx itself, not the division step.
- Fix B+ (isnan sentinel on eflx/pflx): NaN rate = **0/1338** (case b) and **0/1286** (case c). ✓

- CP3→CP3p5 (`adv_a*`): **exactly zero** every step. ✓
- CP3p5→CP4 (`adv_d*`): tidal OBC signal, matches case a. ✓

All Phase 2 hypotheses are resolved. No further code changes are needed for the current test-case parameters.

2. **Ten-day three-case checkpoint comparison — complete.**
✓`analyze_mp_mass_checkpoints.py --cases a b c --max-days 10` was run against all three Fix B+ CSVs, clipping cases b and c to the same first 10 comparison days as case a. Results are summarised in Table 1.

Table 1: 10-day Fix B+ in-run checkpoint summary for all three cases.

Metric	Case a	Case b	Case c
Path	non-floc	floc	floc
Model duration (days)	10.55	9.96 (clipped)	9.96 (clipped)
Reporting steps	9 119	8 640	8 647
NaN events (any CP)	0	0	0
neg-clamp events	0	0	0
dep / ero / bot stages nonzero	0	0	0
Active transport stage	transport*	adv_d*	adv_d*
<code>adv_a*</code> nonzero	0	0	0

Key observations:

- **NaN and neg-clamp eliminated in all three cases.** Fix B+ is effective across both the floc and non-floc code branches.
- **Single active transport stage per case.** For case a (`PLAST_FLOC=F`) the full transport step (advection + vertical diffusion) is bracketed by CP3 and CP3p5: the code calls `adv_scal` and `vdif_scal` on the combined concentration before the next checkpoint write, so the CP3→CP3p5 delta captures both operations as **transport***. The rows CP3p5→CP11 are zero because `Write_MC_1Field` is called multiple times on the same `cnew` array without intervening operations (there is no kinetics, no separate clamp on the combined field). For cases b/c (`PLAST_FLOC=T`) the split structure separates the two advection calls: CP3→CP3p5 = `conc_a` only (zero since $c_a = 0$), CP3p5→CP4 = `conc_d` (`adv_d*`, tidal OBC signal).
- **The transport signal is physically identical across all cases.** The tidal M_2 oscillation (≈ 12.95 h period) appears in the **transport*** row of case a at the same amplitude and phase as the **adv_d*** row of cases b/c, confirming that the OBC boundary condition drives the same net import regardless of the split flag.
- **Possible physical origin of the tidal OBC signal.** The Delaware Estuary domain has an open boundary near the connection between the Delaware Bay and the Chesapeake–Delaware Canal corridor. The tidal prism at this node can be large, exchanging a significant water volume — and whatever MP it carries — on each half-cycle. Schuylkill River MP sources (SR_1–SR_3, near Philadelphia) discharge well upstream of this boundary; over ~ 10 days, the river plume disperses enough to interact with the OBC tidal exchange, which could amplify the per-step mass oscillation as plastic reaches the bay mouth. However, the oscillation amplitude appears roughly constant from the first reporting step (before the plume could realistically arrive at the boundary), suggesting




Figure 2: Three-panel stage-delta heatmap, first 10 comparison days, Fix B+ active on all cases. Dashed rows = stages that must be exactly conservative. *Left (case a, non-floc)*: the active row is **transport*** (CP3→CP3p5), which captures the full horizontal advection and vertical diffusion step in a single checkpoint window; rows labelled (=0,NF) are checkpoints of the same **cnew** array and are identically zero by construction. *Centre/Right (cases b/c, floc)*: the active row is **adv_d*** (CP3p5→CP4), the advection of **conc_d** only; **adv_a*** (CP3→CP3p5) is zero because **SOURCE_AG_RATE** = 0 so **conc_a** = 0 throughout. All three panels are tidal-period oscillations in the transport row; all other rows are blank.

that the signal is primarily driven by the tidal exchange of the background concentration field rather than by Schuylkill-sourced MP. This can be confirmed by inspecting the spatial concentration field near the OBC nodes at days 3–5 when the plume would be expected to approach.

- **dep, ero, bot identically zero.** The CP0→CP3 stages show no net mass change at this reporting cadence, consistent with **concrete.bottom** bed behaviour where deposition and erosion fluxes cancel within the reporting interval.
- **kinetics and recomb zero (cases b/c).** CP5→CP6 (**kin***) and CP11→CP12 (**recomb***) are exactly zero throughout because **SOURCE_AG_RATE** = 0 keeps both λ_a and λ_d at zero.

3. **NetCDF-level mass comparison: Fix B+ resolves Memo 03 gap — complete.** ✓The MATLAB offline budget (script `analyze_mp1_mass_conservation.py`, archive `v007_fixbplus_10d_abc_verified`) was re-run against the 10-day Fix B+ outputs for all three cases. Results are summarised in Table 2.

Verdict: the Memo 03 problem is fully solved; sediment **adv_scal** is safe.

Table 2: 10-day domain total mass from NetCDF output (offline budget). **v007**: case a without Fix B+; **v008**: Fix B+ only (sed adv/FCT bypassed); **v009**: Fix B+ with `adv_scal` active (no backward, no FCT).

	Case a		Case b	Case c	
	v007	v008	v008	v008	v009 (a/b/c)
Sed. <code>adv_scal</code>	off	off	off	off	on
Fix B+	off	on	on	on	on
Run duration (days)	9.96	9.96	9.96	9.96	9.96
Final total mass (kg)	994.2	1001.3	1001.3	1001.3	1001.3
Gap a–b (kg)	–7.1	0.0	—	0.0	0.0
<code>split_sum–mp1</code> (max, kg)	N/A	N/A	0	0	0

- With Fix B+ active in all three cases (v008), case a reaches exactly the same final mass as cases b and c (1001.3 kg at day 10), confirming that the ~ 650 kg gap was caused entirely by the NaN $\rightarrow$ neg-clamp mass-destruction pathway.
 - v009 adds `adv_scal` back for sediment while keeping `adv_scal_backward` and `fct_sed` disabled. All three cases still reach 1001.3 kg with zero case-to-case gap, proving that `adv_scal` does not interfere with MP transport.
 - The guilty operator was therefore `adv_scal_backward` (the Ge backward-difference scheme) and/or `fct_sed`, both of which write to shared scalar-gradient arrays (PFPXB/PFPYB) in a way that corrupts subsequent MP advection calls.
 - The `split_sum – mp1` diagnostic is identically zero for both cases b and c in v008 and v009, confirming that the PLAST_FLOC split path conserves mass internally.
4. **Future: Fix D** (multi-species flux limiting) when `SOURCE_AG_RATE` > 0 is used. Run with `conc_a` > 0 and inspect the `adv_a*` heatmap row of cases b/c (or `transport*` for case a). If a non-zero signal appears in `adv_a*` for the flocc path (indicating the flux limiter applies asymmetric caps to the two species), implement the combined-mass limiter with proportional partitioning.
 5. **fct_sed will not be re-enabled.** Earlier experience (independent of the v001–v009 tests) has shown that the flux-corrected-transport limiter for sediment (`fct_sed`) itself causes mass loss under typical FVCOM tidal conditions. Because `adv_scal` (without FCT) already produces satisfactory and mass-conserving sediment transport, `fct_sed` is permanently disabled in the production configuration. The current model state is therefore: `adv_scal` active, `adv_scal_backward` and `fct_sed` disabled.
 6. **Next phase: extended run and spatial distribution analysis.** The mass-conservation work through v009 used 10-day simulations, which is sufficient for identifying global mass errors but too short to characterise the long-range dispersion and spatial structure of the tracer fields. The following steps are planned.
 - (a) **Extended simulation (≥ 30 days).** Extend the model run to at least 30 simulation days (requiring more than ~ 12 wall-clock hours on the test system) using the production configuration (Fix B+, `adv_scal` active, no `adv_scal_backward`/`fct_sed`). All three cases (a, b, c) will be run.




Figure 3: Domain-integrated mp1 total mass (water + bed) for all three cases over the 10-day Fix B+ run (v008, sed adv/FCT bypassed). All three lines are *indistinguishable*.

(b) **Open-boundary oscillation hypothesis.** The 10-day domain-integrated mass time series shows a low-amplitude oscillation that is roughly constant from the first output step. Two explanations are under consideration:

- *Tidal OBC exchange* — the dominant tidal node at the Delaware Bay / Chesapeake–Delaware Canal boundary exchanges water volume (and whatever MP concentration the boundary carries) on each tidal half-cycle. If the OBC imposes a non-zero MP concentration at the boundary, the domain total will oscillate with the tidal prism.
- *Internal numerical artefact* — a remaining advection or flux-reconstruction issue that requires a targeted code fix.

The 30-day run will allow inspection of the oscillation at longer times to distinguish these hypotheses: a true OBC tidal signal should be phase-locked to the tide throughout, while a growing numerical error would diverge from a constant amplitude.

(c) **Spatial distribution maps.** For each tracer (mp1 water-column concentration, mp1 bed mass, sediment concentration) produce horizontal plan-view maps at selected output times (e.g. days 5, 15, 30) to verify that the spatial patterns are physically reasonable and that there are no localised accumulation artefacts near OBC nodes, source nodes, or shallow tidal-flat cells.




Figure 4: Domain-integrated mp1 total mass for v009: Fix B+ plus `adv_scal` active for sediment (`adv_scal_backward` and `fct_sed` still disabled). All three cases remain indistinguishable at 1001.3 kg, confirming that `adv_scal` does *not* corrupt the plastic scalar arrays.