

# Notes: Nevanlinna analytical Continuation Method

Shuang Liang<sup>\*</sup>

*Institute of Physics, Chinese Academy of Sciences*

(Dated: October 28, 2021)

## Abstract

This is the abstract.

## CONTENTS

I. The analytic continuation problem	2
II. How to solve?	2
III. Nevanlinna analytic continuation	2
A. Schur algorithm	2
References	4

---

<sup>\*</sup> [sliang@iphy.ac.cn](mailto:sliang@iphy.ac.cn)

## I. THE ANALYTIC CONTINUATION PROBLEM

The analytic continuation problem seeks to extract real frequency dynamical information from imaginary-time correlation functions  $G(\tau)$  data. Technically, this is a highly nontrivial task[1]. To see this, we use the relation between  $G(\tau)$  and  $A(\omega)$  [1, 2]:

$$G(\tau) = \int_{-\infty}^{\infty} d\omega \frac{e^{-\tau\omega}}{1 - \lambda e^{-\beta\omega}} A(\omega) = \int_{-\infty}^{\infty} d\omega K(\tau, \omega) A(\omega) \quad (1)$$

where  $K(\tau, \omega) = \frac{e^{-\tau\omega}}{1 - \lambda e^{-\beta\omega}}$  is the kernel. One may consider to solve the problem by firstly discretize  $\tau$  and  $\omega$  and get:

$$G(\tau_i) = \sum_{j=1}^{N_\omega} K_{ij} A(\omega_j) \quad (2)$$

Then do SVD decomposition of rectangular matrix  $K$ , write  $K_{ij} = U_{il} \lambda_l V_{lj}$ . Finally the spectral function reads

$$A(\omega_j) = \sum_{l=1}^{N_\tau} \frac{1}{\lambda_l} V_{lj} \sum_{i=1}^{N_\omega} G(\tau_i) U_{il} \quad (3)$$

It seems fine at the first glance. However, if we consider the properties of  $K(\tau, \omega)$ , we would notice that it is highly singular since it is exponentially small for large  $|\omega|$ , so small errors  $G(\tau)$  would be amplified by exponentially small  $\lambda_l$ . This problem is well-known ill-posed[3, 4] and enormous efforts have been made[].

## II. HOW TO SOLVE?

...

Here we introduce the recently developed Nevanlinna analytic continuation method[5].

## III. NEVANLINNA ANALYTIC CONTINUATION

### A. Schur algorithm

The Nevanlinna analytic continuation method[5] is basically an interpolation method. In this method, one should firstly using conformal transforms to map the Matsubara Green's functions  $\mathcal{G}$ , which is analytic in the upper half of the complex plane  $\mathcal{C}^+$  and contains singularities in the lower half plane, to a closed unit disk  $\bar{\mathcal{D}}$  in the complex plane. The

mappings are shown in fig. 1. It becomes a Schur class( $\mathcal{S}$ ) function and would have a continued fraction expansion where the parameters can be recursively defined[6]. Then one can apply the Nevanlinna iterative algorithm to interpolate the Schur functions[7]. Finally, one can do a inverse conformal transform back to  $\mathcal{C}^+$  and obtains  $\mathcal{G}(z)$ , it's then natural to do analytic continuation  $z \rightarrow \omega + i0^+$ . The calculation process is shown in fig. 2.




FIG. 1. Conformal mappings




FIG. 2. Calculation flow chart

For fermionic Green's functions, the mapping from  $\mathcal{C}^+$  to  $\bar{\mathcal{C}}^+$  is simple. Since  $\text{Im}\mathcal{G}(z) \leq 0$  if  $z \in \mathcal{C}^+$ , the mapping is just to take  $\mathcal{G} \rightarrow -\mathcal{G} = \mathcal{NG}$  and  $\mathcal{NG} \subset N$ . While for bosonic Green's functions, this mapping is a little bit complicated and we will discuss in the next section. The data set we have is  $\{i\omega_n, \mathcal{G}(i\omega_n)\}$ , here we denote  $\mathcal{Y}_n = i\omega_n$  and  $\mathcal{C}_n = \mathcal{NG}(i\omega_n) = -\mathcal{G}(i\omega_n)$ .

Then we use the Möbius transform

$$h(z) = \frac{z - i}{z + i} \quad (4)$$

to map  $\mathcal{C}_n \subset N$  to  $\theta(\mathcal{Y}_n) = h(\mathcal{C}_n) \subset \bar{\mathcal{D}}$ . The recursive final  $\theta(z)$  can conveniently be written in a matrix form:

$$\theta(z)[z; \theta_{M+1}(z)] = \frac{a(z)\theta_{M+1}(z) + b(z)}{c(z)\theta_{M+1}(z) + d(z)} \quad (5)$$

where

$$\begin{pmatrix} a(z) & b(z) \\ c(z) & d(z) \end{pmatrix} = \prod_{n=1}^M \begin{pmatrix} h_1(z, \mathcal{Y}_j) & \phi_j \\ \phi_j^* h_1(z, \mathcal{Y}_j) & 1 \end{pmatrix} \quad (6)$$

where  $h_1(z, \mathcal{Y}_n) = \frac{z - \mathcal{Y}_n}{z - \mathcal{Y}_n^*}$  is a conformal map from  $\mathcal{C}^+$  to  $\mathcal{D}$ .  $\theta_j(z)$  is the interpolation function of  $j$ -th step and  $\phi_j = \theta_j(\mathcal{Y}_j)$ . There is a freedom to choose  $\theta_{M+1}(z)$ . One can use this freedom to select the “best” of all consistent spectral functions.

- 
- [1] Mark Jarrell and James E Gubernatis. Bayesian inference and the analytical continuation of imaginary-time quantum monte carlo data. *Physics Reports*, 269(3):133–195, 1996.
  - [2] Ming-Wen Xiao. *Lecture notes: Linear Response Theory*. School of Physics, Nanjing University.
  - [3] Forman S. Acton. *Numerical Methods that Work (Spectrum)*. The Mathematical Association of America, 1997.
  - [4] Ingo Peschel, Xiaoqun Wang, Matthias Kaulke, and Karen Hallberg. *Density-Matrix Renormalization-a New Numerical Method in Physics: Lectures of a Seminar and Workshop Held at the max-planck-institut für physik komplexer systeme, Dresden, Germany, August 24th to September 18th, 1998*. Springer, 1999.
  - [5] Jiani Fei, Chia-Nan Yeh, and Emanuel Gull. Nevanlinna analytical continuation. *Physical Review Letters*, 126(5):056402, 2021.
  - [6] J Schur. Über potenzreihen, die im innern des einheitskreises beschränkt sind. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 1918(148):122–145, 1918.
  - [7] Rolf Nevanlinna. Über beschränkte funktionen, die in gegebenen punkten vorgeschriebene werte annehmen. *Ann. Acad. Sci. Fenn. Ser. A 1 Mat. Dissertationes*, 1919.