

بسم الله الرحمن الرحيم

هذا الملف تجميع لبعض الأسئلة من ود الشمال 4 المكررة و عزمي
والغامدي يوجد راشونال لبعض الأسئلة سأكملها جميعا لاحقا

ويوجد بالأخير تصحيح لبعض مستنداتي السابقة

احب ان اشكر الطلاب الذين اجتهدوا لتصحيح بعض الأسئلة اسأل الله
التوفيق لهم

لا ابيع من يستخدم هذا الملف لغرض شخصي (مادي)

اتمنى من الجميع حذف الملف السابق

Maryam collection part 1

Respiratory system

1. A nurse is caring for a child with a diagnosis of cystic fibrosis and pneumonia. The plan of care includes nebulizer treatment and chest physiotherapy. The nurse should perform chest physiotherapy:

- a. Continuously during the nebulizer treatment
- b. Prior to the nebulizer treatment
- c. After the nebulizer treatment**
- d. Intermittently during the nebulizer treatment

2. A patient with a pulmonary embolus and a nursing diagnosis of impaired gas exchange has an order to obtain arterial blood gases. The FIRST intervention by the nurse is to:

- a. Perform an Allen's test
- b. Explain the procedure
- c. Gather the equipment**
- d. Document the procedure

<http://www.revise4finals.co.uk/medicine/learn/abg.php>

3. While caring for a patient in the post-anesthesia care unit (PACU), a nurse observes the onset of rapid breathing cyanosis, and narrowing blood pressure. The nurse should plan to:

- a. Administer bolus glucose
- b. Suction the airway**
- c. Turn the patient to the right side
- d. Administer intra venous fluids

4. While caring for a patient with in effective airway clearance related to increased mucus production, the nurse should encourage fluids to:

- a. Maintain nutrition
- b. Prevent boredom
- c. Stimulate coughing

d. Thin secretions

5. A patient with advanced lung cancer is exhibiting cyanosis and edema of the head and upper extremities. Which of the following intervention would MOST likely provide an immediate benefit for this patient?

- a. Place in Trendelenburg position
- b. Position on the right side

c. Elevate the head of the bed

- d. Elevate extremities

6. A nurse is administering tuberculin skin tests purified protein derivative (PPD), which of the following time frames should the nurse tell the patient to return to the clinic for the test to be read?

- a. In 12-24 hours
- b. In 24-36 hours
- c. In 36-48 hours

d. In 48-72 hours

7. A patient who is scheduled for a tonsillectomy is in pre-operative unit. The nurse notes an order for pre anesthetic medication to be given “on call to operation room”. The nurse should give this medication:

- a. Immediately upon being notified to prepare the patient for transport**

- b. When the operation room staff arrive to transport the patient
- c. Only if clearly needed after assessment
- d. Upon the patient's arrival in the operation room

8. Which gauge needle should the nurse use for an intradermal injection on the ventral surface of the forearm for patient with TB?

- a. 16 gauge needle
- b. 20 gauge needle
- c. 22 gauge needle
- d. 26 gauge needle**

9. A nurse makes a home visit to a patient recently diagnosed with chronic obstructive pulmonary disease (COPD), which of the following should the nurse teach the patient about managing COPD?

- a. Recognizing signs of impending respiratory infection**
- b. Limiting fluids intake minimize bronchial secretions
- c. Correct technique to auscultate the lung fields
- d. Importance of starting antibiotic therapy

10. A patient with chronic obstructive pulmonary disease (COPD) experiencing frequent dyspnea which of the following exercise would teach the patient how to BETTER control breathing?

- a. Lower side rib
- b. Segmental
- c. Pursed-lip**
- d. Diaphragmatic

11. Patient with asthma is experiencing thick respiratory secretions resulting in increased work of breathing. The best nursing intervention is to:

a. Encourage fluids

b. Eliminate dairy products

c. Decrease relative humidity of the room

d. Have the child lay on the left side.

12. What would be the long-term goal for a patient with asthma?

a. Quickly reverse airflow obstruction

b. Correct hypoxemia

c. Deliver humidified oxygen via nasal cannula

d. Develop a home and school management plan

13. A nurse administers an albuterol nebulizer on a patient with asthma exacerbation. Which of following indicates effectiveness of the treatment?

a. Adventitious breath sound with cough

b. O₂ saturation 94%

c. Nasal flaring

d. Respiration rate 28

14. A patient is diagnosed with asthma exacerbation . Which of the following nursing diagnoses should be the FIRST priority?

a. In effective airway clearance related to broncho spasm and mucosal edema

b. Fatigue related to hypoxia

c. Anxiety related to illness and loss of control

d. Deficient knowledge related to potential side effect of the medication

15. An asthmatic patient presents with wheezing and coughing. Oxygen saturation is 88% on room air. Which of the following nursing diagnosis would take priority?

a. Imbalanced nutrition related to decreased food intake

- b. Activity intolerance related to inefficient breathing
- c. Anxiety-related dyspnea and concern of illness
- d. Ineffective gas exchange related to broncho spasm

16. A patient presents to the emergency department with complaints of head ache, dizziness and confusion. Clinical symptoms include tachypnea and dyspnea with the use of accessory muscles to facilitate breathing. Which of the following orders would the nurse MOST likely implement to reduce the patient's confusion and disorientation?

- a. Oxygen therapy
- b. Chest physical therapy
- c. Bronchodilators
- d. Hydration fluids

17. The nurse is caring for a patient who just had a chest tube inserted due to spontaneous pneumothorax . An appropriate goal is that the patient will:

- a. Be free of pain within 4 hours
- b. Report decreased pain
- c. Rest quietly
- d. Sleep with few movements

18. A home health nurse visits a patient with chronic obstructive pulmonary disease (COPD) using home oxygen at 2 liters per minute. The patient reports periods of shortness of breath and inquires about increasing the oxygen to 4 liters/minute. The nurse explains that increasing the supplemental oxygen will:

- a. Increased activity tolerance
- b. Suppress the hypoxic drive

- c. Alleviate the shortness of breath
- d. Prevent lung infection

19. A patient exhibits clinical manifestation of a pulmonary embolism. Arterial blood gas (ABG) levels and a chest x-ray are ordered. Which of the following test is used to diagnose this condition?

- a. Computer tomography scan(CT scan)
- b. Magnetic resonance imaging (MRI)
- c. Pulmonary angiography
- d. Pulmonary function test

20. A nurse is caring for a child who is post-tonsillectomy and adenoidectomy. The nurse should plan to assess which of the following complications?

- a. Pulmonary hypertension
- b. Hemorrhage
- c. Hearing loss

21. A nurse caring for a patient with acute pulmonary edema observes that the patient's cough produces white, frothy and that the patient is extremely dyspneic. The patient has inspiratory and expiratory wheezing on auscultation of the lungs. The immediate objective of treatment is to

- a. Improve oxygenation
- b. Decrease anxiety
- c. Improve tissue perfusion
- d. Decrease risk for aspiration

22. While performing an assessment on a post-surgical patient 2 days after surgery, the nurse notes shallow and rapid respirations. What should the nurse do NEXT?

- a. Asses the patient from pain
- b. Obtain an order from supplemental oxygen
- c. Elevate the head of the bed
- d. Place a warmed blanket on the patient

23. The nurse is caring for child admitted with viral pneumonia. Which of the following nursing diagnoses should receive PRIORITY?

- a. Nutrition altered: less than body requirements
- b. Ineffective airway clearance
- c. Fluid volume deficit
- d. Risk for injury

24. The nurse receives an order to obtain an arterial blood gas (ABG) specimen on a patient. The nurse will use the radial artery to obtain the specimen . Which of the following will the nurse assess before puncturing the radial artery?

- a. Allen test
- b. Partial pressure of arterial oxygen
- c. Partial carbon dioxide
- d. Prothrombin time

25. A patient with pneumonia has a temperature, 40 C (104 F); heart rate 20;respiratory rate 32 and dyspnea patient has an ineffective airway clearance related to excessive tracheobronchial secretions. Which of the following interventions would the nurse implement to enhance the patient's airway clearance?

- a. Administer oxygen as ordered
- b. Maintain a comfortable position
- c. Increase fluid intake
- d. Administer prescribed analgesic

26. When a child is brought to the emergency department with acute epiglottitis, which of the following nursing diagnoses should receive PRIORITY?

- a. Ineffective airway clearance**
- b. Activity intolerance
- c. Fluid volume deficit
- d. Impaired verbal communication

27. A patient with pneumonia experiences ineffective airway clearance related to the presence of thick secretions secondary to infection. Oxygen saturation is 89% on room air. Which of the following nursing interventions takes priority?

- a. Deliver oxygen with humidity**
- b. Encourage fluid intake
- c. Assist patient into position of comfort
- d. Inspect sputum for odor and color

28. During the immediate postoperative period, a patient reveals an oxygen saturation level of 91%. The nurse should:

- a. Position the patient on the left side
- b. Administer supplemental oxygen**
- c. Continue to provide supportive care
- d. Lower the temperature of the room

29. A patient presents to the emergency department with difficulty breathing. The patient's Parents report that patient has a history of bronchial asthma and has recently had an Upper respiratory tract infection (URI). Upon auscultation, the nurse decreased Breath sounds in the left-lower lung field. The nurse should NEXT assess the child's:

- a. Oral temperature

b. O2 saturation

c. Apical pulse

d. Level of comfort

30. A patient with pulmonary emboli complains of pain, dyspnea , and a fear of dying. Which of the following interventions would MOST likely help to reduce the patient's anxiety level?

a. Administer oxygen as ordered

b. Administer pain medication as ordered

c. Observe closely for signs of pain and discomfort

d. Listen to the patient's concerns

31. The nurse is caring for a client who has had a chest tube inserted and connected to water seal drainage. The nurse determines the drainage system is functioning correctly when which of the following is observed:

a- Continuous bubbling in the water seal chamber

b- Fluctuation in the water seal chamber

c-Suction tubing attached to a wall unit

d-Vesicular breath sounds throughout the lung fields

32. A patient who underwent hand surgery requiring general anesthesia presents to the post anesthesia care unit (PACU) after extubation, The nurse should FIRST assess

a. Circulatory status

b. Wound status

c. Respiratory status

d. Hydration status

33. Prior to administration of an albuterol nebulizer, the nurse should help the patient assume what position?

- a. Sitting and leaning forward
- b. Feet elevated above level of heart

c. High fowler's

d. Standing

34. A 32 year old female comes in for evaluation 14 days after an uncomplicated caesarean section . The patient is very anxious and complaining of sharp stablign pain in her chest .The patient has dyspnea , tachypnea , and hypoxemia .Which of the following postoperative complications is likely?

a. Pulmonary embolism

- b. Atelectasis
- c. Pneumonia
- d. Aspiration

35. A conscious victim of motor vehicle accident arrives at the emergency department. The patient gasping of air , is extremely anxious , and has a deviated trachea . What diagnosis should the nurse anticipate?

a. Pleural effusion

b. Tension pneumothorax

- c. Pneumothorax
- d. Hemothorax

Brunner and Suddarth's Textbook of Med.-Surg.
Nursing 12th ed chapter 23/ page 597

Essentials of Critical Care Nursing: A Holistic
Approach by Patricia Gonce Morton, Dorrie
K. Fontaine chapter 17/ page 243

36. A patient with chronic obstructive pulmonary disease complains of a frequent cough, bilateral wheezing is auscultated in the lung fields. The nurse administers albuterol nebulizer treatment, as

ordered and educates the patient on way to decrease exacerbation. Which of the following actions indicate that the patient understands the instruction?

- a. The patient reduces number of cigarettes smoked per day
- b. The patient requested a pneumococcal vaccination**
- c. The patient increases sodium and potassium intake
- d. The patient exercises whenever experiencing shortness of breath

Brunner and Suddarth's Textbook of Med.-Surg. Nursing 12th ed chapter 24/ page 617

37. A child with asthma has an order for albuterol . Prior to administration of medication the nurse must:

- a. Pre-oxygenate the patient
- b. Assess the patient's heart rate**
- c. Obtain venous access
- d. Feed the patient a snack

38. A nurse is caring for a patient who is 6 hours post left lobectomy. On assessment the nurse observes that the patient has become very restless and the nail beds are blue. the vital signs reveal tachycardia, tachypnea and blood pressure is rising. Which of the following complication is MOST likely?

- a. Pneumonia
- b. Hypoxia**
- c. Postoperative bleeding
- d. Broncho pleural fistula

39. A nurse is caring for a patient who had a pneumonectomy 2 days ago for lung cancer. Which observation would indicate that the patient is progressing towards discharge goal?

- a. Cough productive of serosanguineous fluid

Symptoms of Bronchopleural fistula
common complication of
pneumonectomy

b. 1+ pretibial edema

c. Nap after completing bed bath

d. Frequent premature ventricular contractions (PVC)

40. A patient is scheduled for a pneumonectomy in the morning. Which of the following diagnosis is the MOST likely indication for this type of surgery?

a. Lung carcinoma

b. Pulmonary tuberculosis

c. Benign pulmonary nodule

d. Mediastinal shift

41. Patient came to emergency department with coughing and difficulty in breathing, you suspect patient has asthma attack, the sound you expect to hear is:

a. Wheezing

b. Chronic

c. Crackle

42. Upon assessment of an obese patient's back, the nurse observes a forward curvature of the lumbar spine. These findings are indicative of:

a- Kyphosis

b- List

c- Lordosis

d- Scoliosis

43. A 32 year-old man is brought to the Emergency Department by a friend with whom he had been playing football with earlier that evening. His symptoms came on after twenty minutes of the football match. He appears anxious and restless. Auscultation of the lungs reveals wheezing on exhalation. The patient states that he is unable to get a full breath of air. He had this problem as a child a couple of times after exercising in cold weather.

Blood pressure 126/72 mmHg

Heart rate 90/min

Respiratory rate 28/mm

Temperature 37.0°C

Oxygen Saturation 94 % on room air **Which position is best for this patient?**

- a- Supine
- b- Reverse Trendelenburg
- c- Prone
- d- High fowler's**

44. The nurse is caring for a client who has just had a chest tube attached to a water seal drainage system. To ensure that the system is functioning effectively the nurse should:

- a- Observe for intermittent bubbling in the water seal chamber**
- b- Flush the chest tubes with 30-60 ml of NSS every 4-6 hours
- c- Maintain the client in an extreme lateral position
- d- Strip the chest tubes in the direction of the client

45. A 30 year-old woman has been prescribed albuterol PNR1 and prophylactic inhaled corticosteroids to be taken once per day. She has been taking the prophylactic as prescribed but has needed to use the albuterol more often than usual. She has a chronic cough and often has air hunger.

Which intervention is initially most appropriate?

- a- Refer for a chest X-ray
- b- Administer magnesium sulfate
- c- Assess peak flow measurement
- d- Perform arterial blood gas analysis**

46. A patient on a medical floor is diagnosed with pneumonia. Which of the following interventions is appropriate for a

nursing diagnosis of ineffective airway clearance related to tracheobronchial secretions secondary to infection?

- a- Explain diagnostic procedures and address any of patient's concerns
- b- Assess patient for signs of discomfort and pain
- c- Monitor oxygen therapy and encourage to cough and deep breathe**
- d- Monitor patient for shock

47. A patient with bacterial pneumonia reports increased pain during deep breathing and coughing exercises. To minimize the patient's pain, the nurse should teach the patient to.

- a- Take a cough suppressant prescribed P.R.N at regular time intervals
- b- Turn onto the side before doing the respiratory exercises
- c- Hold a pillow tightly against the chest while coughing**
- d- Drink warm liquids right before taking deep breaths

48. A patient with a history of chronic obstructive pulmonary disease prepares for a colonoscopy with conserve sedation. Due to the patient's history the nurse should very cautious about administering.

- a- Antibiotics
- b- Sedation
- c- Oxygen**
- d- Intravenous fluid

49. A home care patient with chronic obstructive pulmonary disease (COPD) reports an upset stomach. The patient is taking theophylline (Theo-Dur) and triamcinolone acetanide (Azmecort), the nurse should instruct the patient to take

- a- Theo-Dur on an empty stomach
- b- Theo- Dur and Azmecort at the same time
- c- Theo- Dur and Azmecort 12 hours apart
- d- Theo-Dur with milk or crackers**

50. The nurse is assessing a patient who just had surgery under general anesthesia. The patient's respiration rate is 4 per minute and the O₂

saturation on 3L per minute of oxygen via nasal cannula is 84%. The nurse is awaiting the results of an arterial blood gas (ABG) and anticipates that which of the following will be **ELEVATED?**

- a- Arterial oxygen saturation (SaO₂)
- b- Hydrogen ion concentration (pH)
- c- Partial pressure of arterial oxygen (PaO₂)

d- Partial pressure of arterial carbon dioxide (PaCO₂)

52. A patient presents with pneumonia. Which of the following intervention would the nurse implement to improve the patient's gas exchange?

Heart rate 80/min

Respiratory rate 32/min

Temperature 40°C

- a- Encourage the patient to increase fluid intake**
- b- Position the patient in an upright position
- c- Instruct the patient the patient to splint when coughing
- d- Apply heat and or cold to the chest as ordered

53.



A nurse is instructing parents to perform chest Physiotherapy (CPT) on their son with pneumonia the nurse has the parents place their son in a prone position with the body tilted downward before performing CPT.

Based on this information, the nurse is instructing the parents to remove drainage from which of the following lung lobe segment?

- a- Upper posterior
- b- Lower anterior

c- Lower posterior

d- Upper anterior

54. The charge nurse enters the nursing diagnosis "Risk for ineffective airway clearance related to an inability to swallow" on the client's care plan. Which nursing intervention is most appropriate for managing the identified problem?

a- Keeping the client supine

b- Removing all head pillows

c- Performing oral suctioning

d- Providing frequent oral hygiene

55. Which of the following intervention would the nurse implement to enhance the patient's airway clearance ?

Heart rate 80/min

Respiratory rate 32/min

Temperature 40°C

A. Administer oxygen as ordered .

B. Maintain a comfortable position .

C. Increase fluid intake .

D. Administer prescribed analgesics

56. A 7-year-old client is brought to the E.R. He's tachypneic and afebrile and has a respiratory rate of 36 breaths/minute and a nonproductive cough. He recently had a cold. From his history, the client may have which of the following?

a- Acute asthma

b- Bronchial pneumonia

c- Chronic obstructive pulmonary disease (COPD)

e- Emphysem

57. The term "blue bloater" refers to which of the following conditions?

a- Adult respiratory distress syndrome (ARDS)

b- Asthma

c- Chronic obstructive bronchitis

d- Emphysem


58. A nurse is suctioning fluids from a client via a tracheostomy tube. When suctioning, the nurse must limit the suctioning to a maximum of:

- a- 5 seconds
- b- 10 seconds**
- c- 30 seconds
- d- 1 minute

59. Which of the following diets would be most appropriate for a client with COPD?

- a- Low fat, low cholesterol
- b- Bland, soft diet
- c- Low-Sodium diet
- d- High calorie, high-protein diet**

60.



Which of the above location represent the best sites to inspect chest retraction in child with lower respiratory tract disorder?

- a- A&B
- b- B&C
- c- C&D
- d- D&E**

61. Following picture suggest patient has:

- a- Renal disease
- b- Hepatic disease
- c- Pulmonary disease**
- d- Cerebral disease



62. A patient with COPD is admitted to hospital, how can the nurse best position patient to improve gas exchange?

- a- Setting up at bedside on chair
- b- Resting in bed with HOB 45 to 60 degrees
- c- Trendelenburg position
- d- High fowler position with knee flexed

63. proper position for patient during oropharynx suctioning:

- a- Prone
- b- Supine
- c- Semi fowler
- d- Trendelenburg

Cardiac system

1. A patient is diagnosed with pulmonary hypertension. Which of the following nursing diagnoses should be the **PRIORITY**?
 - a. Impaired gas exchanged related to altered blood flow secondary to pulmonary capillary constriction
 - b. Fatigue related to hypoxia
 - c. Anxiety related to illness and loss of control
 - d. Activity intolerance related to imbalance between oxygen supply and demand due to right and left ventricular failure

2. A patient with congestive heart failure and severe peripheral edema has a nursing diagnosis of fluid volume excess. What are the two **MOST** important interventions for the nurse to initiate?
 - a. Diuretic therapy and intake and output
 - b. Nutritional education and low-sodium diet
 - c. Daily weights and intake output
 - d. Low-sodium diet and elevate legs when in bed

3. A patient has exacerbation of congestive heart failure, with one of the nursing diagnosis being excess fluid (Lasix). The nurse closely monitors fluid intake and output and administers furosemide (Lasix). Which of the following indicates the efficacy of the nursing intervention?
 - a. The patient has leg edema
 - b. The patient has shortness of breath
 - c. The patient has decreased in weight
 - d. The patient has jugular vein distention

4. A patient is admitted to the hospital with congenital heart disease. Which of the following nursing diagnoses should receive **PRIORITY**?
 - a. Decreased cardiac output related to decreased myocardial function
 - b. Activity intolerance related to cachexia
 - c. Impaired gas exchanged related to altered pulmonary blood flow
 - d. Imbalanced nutrition: less than body requirements related to excessive energy demands

- 5. Prior to initiating therapy with unfractionated heparin for a patient hospitalized with a deep vein thrombosis, this treatment requires:**
- a. Bed rest**
 - b. Aspirin therapy
 - c. Fluid restrictions
 - d. A high protein diet
- 6. A 59-year-old patient arrives in the emergency department diaphoretic and complains of chest pain and shortness of breath. The patient's sibling states that this has happened before and it is just anxiety. Upon evaluation the physician diagnosis unstable angina and prescribes anti-anginal medications. What is the expected results of this drug therapy**
- a. Balanced between oxygen supply and demand
 - b. Increase in blood flow to the heart
 - c. Reduction in oxygen demand and consumption
 - d. Vessel relaxation**
- 7. A patient with end-stage cardiomyopathy and angina pectoris to the office complaining of frequent chest pain and severe dyspnea. With a nursing diagnosis of alteration in comfort, what is the BEST long term goal for this patients?**
- a. Perform all activities of daily living without complaints of chest pain or shortness of breath**
 - b. Verbalize and employ strategies to decrease pain and increase coronary blood flow
 - c. Take pain medications around the clock and use supplemental oxygen at all times
 - d. Understand the disease process and accept the limitation that it places on his lifestyle
- 8. A patient has an order for a pneumatic compression device. Which of the following is an appropriate goal?**
- a. Reduce the risk deep vein thrombosis**
 - b. Reduce lower extremity edema

- c. Reduce lower extremity pain
 - d. Reduce the risk of phlebitis
- 9. A patient has an elevated prothrombin (PT) time. Which medication should the Nurse consider as a possible cause of the elevated PT Time?**
- a. Rifampin
 - b. Vitamin K
 - c. Birth control pills
 - d. Phenytoin (Dilantin)**
- 10. A healthy patient is in doctor's office for a pre-operative visit before a total replacement. The nurse interviewing the patient charts the following medications: aspirin 81 mg once a day, vitamin E 260 international units once a day, and unknown amount of a herbal supplement once a day, based on the patient's medication list which of the following labs would be important pre operatively?**
- a. Prostate specific antigen(PSA)
 - b. Blood glucose
 - c. Creatine phosphokinases enzymes (CPK enzymes)
 - d. Prothrombin time**
- 11. A patient with the deep vein thrombosis (DVT) is being treated with a low-molecular weight heparin.(LMWH). The patient reports increased pain in the affected extremity. The nurse observe the affected extremity has increased in size by 0.2 cm (0.8 inches) during the past 24 hours. Which of the following actions should the nurse take?**
- a. Administer the next dose of LMWH before the scheduled time.
 - b. Apply dry heat to the site .
 - c. Elevate the extremity**
 - d. Reinforce the importance of ankle circling exercises
- 12. A patient has pacemaker implanted. Which of the following interventions is appropriate for the nursing diagnosis of risk for injury?**

- a. Have patient avoid exposure to magnetic resonance imaging(MRI)
- b. Observe incision site for redness, purulent drainage,
- c. Offer back rubs to promote relaxation
- d. Instruct patient in dorsiflexion exercises of ankles

13.A patient is who is prepared for hip surgery has an order for external pneumatic compression devices. The nurse teaches the patient that pneumatic compression can help prevent:

- a. Upper respiratory infection
- b. Decreased breath sounds
- c. Deep vein thrombosis
- d. Bleeding at the surgical site

14.A patient presents with a productive cough with a moderate amount of white Frothy sputum and dispend. The patient is anxious and the nurse notices on assessment that the patient is using accessory muscle including intercostals spaces to breathe and has jugular vein distention. The patient has a history of hypertension and heart failure. What should the nurse administer FIRST?

- a. Digoxin (lanoxin) to improve the ability of the heart to pump effectively
- b. Oxygen therapy to combat hypoxemia
- c. Furosemide (lasix) to reduce blood volume and pulmonary congestion
- d. Morphine sulfate (Duramorph) to reduce anxiety

15.A nurse is caring for a postoperative patient who is on subcutaneous, low dose heparin. This medication is used to prevent:

- a. Deep vein thrombosis
- b. Congestive heart failure
- c. Paralytic Ileus
- d. Pneumonia

16. A patient required long-term antibiotic has a central line catheter inserted into the right subclavian vein by the physician . Which of the following must be verified prior to the first use of the catheter?

- a. Blood return
- b. X-ray**
- c. Catheter potency
- d. Length of catheter

17. Which of the following tests measures the total quantity of Prothrombin In the blood and monitors the effectiveness of warfarin sodium (Coumadin) therapy and prolonged deficiencies in the extrinsic factor?

- a. Thrombin time (TT)
- b. Prothrombin time (PT)**
- c. Partial Prothrombin time (PTT)
- d. Activated partial thromboplastin time (aPTT)

18. One month after starting new medications for hypertension, a patient returns to the clinic with blood pressure in the range. The patient admits to taking the medications only when “feeling bad” Which of the following actions would the nurse take?

- a. Assess further determine the reason the reason for the patient's Actions
- b. Add a new diagnosis of non-compliance
- c. Re-educate the patient about the importance of following his medication plan**
- d. Reevaluate the need for daily medication since the blood pressure is acceptable

19. When caring for a patient who is receiving anticoagulant medications, the nurse MUST monitor the patient, for signs of:

- a. Skin breakdown
- b. Bleeding**
- c. Pain
- d. Confusion

- 20. A nurse is caring for a post-operative patient who is on subcutaneous, low dose of heparin. When administering injection on the abdomen, the nurse avoids the umbilicus area because of the possibility of :**
- a. Entering a larger body vessel**
 - b. Causing increased pain
 - c. Precipitating hyper ventilation
 - d. umbilical infection
- 21. The nurse is caring for patient with deep vein thrombosis (DVT) . The patient's heparin sodium infusion has been discontinued and the patient is receiving prescribed warfarin sodium (Coumadin). The nurse should advise the patient that which of the following needs to be continued?**
- a. Daily complete blood count (CBC)
 - b. Laboratory tests for partial thromboplastin time (PTT)
 - c. Strict bed rest
 - d. Wearing elasticized support stockings**
- 22. A nurse is discharging a patient after hospitalization due to myocarditis. Which of the following statements should be included in discharge teaching?**
- a. There is usually some residual heart enlargement**
 - b. May resume previous activities as before hospitalization
 - c. Avoid immunizations against infectious disease
 - d. Rapidly beating heart is a common side effect of the illness and is not dangerous
- 23. Which of the following discharge planning instructions takes PRIORITY in patient with congestive heart failure?**
- a. Maintaining a low cholesterol, low sodium and low potassium diet
 - b. Recognizing signs and symptoms that require immediate medical attention**
 - c. The importance of remaining physically active
 - d. The importance of drinking plenty of fluid

24. A patient is to receive heparin sodium, 5,000 U, subcutaneous on call to the operating room . Prior to administering this medication, the nurse should advise patient that this will help to prevent:

- a. Infections
- b. Atelectasis
- c. Thrombosis formation**
- d. Positioning injuries

25. While caring for a patient with potassium deficiency, the nurse should expect that the patient may exhibit:

- a. Dysrhythmias**
- b. Oliguria
- c. Diminished deep-tendon reflexes
- d. Hypertension

26. A Patient has a dissection aortic aneurysm. The patient's surgery would be categorized as:

- a. Elective
- b. Urgent
- c. Emergency**
- d. Diagnostic

27. A patient scheduled for an abdominal aneurysm repair. This is what type of surgical intervention?

- a. Diagnostic
- b. Transplant
- c. Curative**
- d. Palliative

28. When discussing dietary choice with a patient who is receiving heparin therapy, the nurse should state that which of the following foods affect the clotting time?

- a. High protein foods
- b. Soy- based foods

c. Foods high in vitamin K

d. Foods containing goat's milk

29. A patient with heart failure has the following vital signs: blood pressure level, 136/84 mmHg, heart rate 48, temperature 37.1 C (98.8 F); and respiration rate 20 per minute. Which of these vital signs should be reported to the physician prior to administering the next dose of digoxin?

a. Blood pressure

b. Pulse

c. Temperature

d. Respiration rate

30. The nurse is caring for a patient two hours after a pacemaker placement. The patient suddenly starts complaining of chest pain. The nurse observes dyspnea, cyanosis and absent breath sounds on the right side. The nurse should anticipate what complications?

a. Hemothorax

b. Perforation of the heart

c. Pneumothorax

d. Hemorrhage

31. A patient is 24 hours post-operative after having a right total hip arthroplasty, the patient complains of pain in the right calf rated 6 on a scale of 0 no pain to 10 severe pain. The nurse observes that the right calf is warm and tender to touch, while the right foot is pale and cool. There is edema from the toes up the knee. The nurse recognizes that these are the classic signs of:

a. Ineffective tissue perfusion

b. Fluid overload

c. Arterial occlusion

d. Deep vein thrombosis

32. What is the first intervention for a client experiencing MI?

a. Administer morphine

b. Administer oxygen (wasn't from the list)

- c. Administer sublingual nitroglycerin
- d. Obtain an ECG

33. A 50 year-old male presents to the medical office 3 weeks after cardiac surgery with complaints of a feeling of weakness, difficulty breathing, and joint pains. Upon examination the nurse finds a fever and a friction rub on auscultation of the chest. The nurse recognizes that the MOST likely surgical complication is:

- a. Neuropsychological dysfunction
- b. Postpericardiotomy syndrome**
- c. Cardiac tamponade
- d. Phrenic nerve damage

34. Aortic aneurism operation what type of surgical intervention:

- a. Elective
- b. Urgent
- c. Emergency**
- d. Diagnostic

35. A patient is scheduled for a bowel resection. The preoperative plan of care includes putting on anti-embolism stockings prior to the transferring the patient to the operating room. What these stockings will do?

- a. Promote venous return**
- b. Minimize joint stillness
- c. Encourage sustained maximum inspiration
- d. Support intestinal peristalsis

36. On the auscultation, the physician heard the patient's heart with crackle sound and wheezing, where could the problem exist?

- a. left ventricular**
- b. left atrium
- c. right ventricular

37. Patient has swelling on both legs, to assess and diagnosis this, which of the following i should do?

- a. auscultation
- b. palpitation**
- c. inspection

38. A client with angina complains that the angina pain is prolonged and severe and occurs at the same time each day, most often in the morning, On further assessment a nurse notes that the pain occurs in the absence of precipitating factors. This type of anginal pain is best described as:

- a- Stable angina
- b- Unstable angina
- c- Variant angina**
- d- Non anginal pain

39. A patient with congestive heart failure and severe peripheral edema has a nursing diagnosis of fluid volume excess What are the two MOST important interventions for the nurse to initiate?

- a- Diuretic therapy and intake and output
- b- Nutritional education and low-sodium diet
- c- Daily weights and intake and output**
- d- Low-sodium diet and elevate legs when in bed

40. A 26 year- old female patient's radial pulse is assessed for a full minute. The nurse notes the pulse is difficult to palpate and count. What should the nurse chart?

Blood pressure 82/48mmHg

Respiratory rate 20/min

Temperature 38.8°C

0	Absent pulse
+1	Weak and thread pulse
+2	Normal
+3	Bounding pulse

- a- 0
- b- 1**
- c- 2
- d- 3

41. A patient recently underwent coronary artery graft (CABG) surgery.

Which of the following nursing diagnoses is the PRIORITY?

- a- Anxiety
- b- Impaired gas exchange**
- c- Acute pain
- d- Sleep deprivation

42. A patient presents to the clinic for a routine visit and has the following vital signs: temperature 37.00 C (98.60 F), heart rate 82, respiration rate 18 and blood pressure level of 130/94 mmHg. Which vital sign is abnormal?

- a. Temperature
- b. Pulse
- c. Respiration
- d. Blood pressure**

43. A 20 year-old woman takes cyclosporine 100 mg by mouth twice per day. She had a heart transplant two months ago and complains of dizziness, shortness of breath, light-headedness and fatigue. Capillary refill >3.

Blood pressure 90/50 mmHg

Heart rate, irregular 100/min

Respiratory rate 28/min

Temperature 38.0°C


What is the most likely underlying cause of the patient's problem?

- a- Cyclosporine toxicity
- b- Hyperkalemia
- c- Organ rejection**
- d- Infection

44. A 65 year-old woman with a history of unstable angina and hypertension presents to the Emergency Department with a dull chest pain that she describes as similar to heartburn. The pain radiates down the left arm. She had taken sublingual nitroglycerin tablets without any relief. An electrocardiograph is performed and shows elevated S-T segments. The nurse is to administer a thrombolytic by intravenous infusion. Which factor places this candidate at high risk for bleeding?

- a- Unstable angina
- b- Hypertension**
- c- Age

d- Elevated S- T segments



45. nurse changes the dressing of a patient who has a chronic lower leg ulcer and a history of leg swelling. The wound bed is 3 cm diameter, red in color and is covered with fibrous yellow tissue (see image) What type of ulcer is most likely?

- a- Arterial
- b- Venous**
- c- Decubitus
- d- Neuropathic

46. The nurse is preparing a discharge plan for a 65 year-old man with a new diagnosis of congestive heart failure. The discharge orders include furosemide 40 milligrams by mouth twice per day. The nurse recommends foods to reduce unwanted medication side effects. Which of the following foods would be most appropriate to include in the teaching plan?

- a- Green and leafy vegetables
- b- Bananas and oranges**
- c- Chicken and Fish

47. A patient present to the emergency department with complaint of chest pain on exertion, shortness of breath, and dizziness. Upon examination the nurse notes palpitation and a systolic ejection murmur at the right upper sterna border. Skin color is normal. The nurse suspect.

- a- Aortic stenosis**
- b- Aortic regurgitation
- c- Mitral stenosis
- d- Mitral valve prolapse

48. A 64 year-old man present with a weight gain of four kilograms in the past week. A physical examination shows an enlarged liver, an enlarged abdomen with a fluid wave, and jugular venous distension. The nurse creates a care plan based on the following nursing diagnosis. Decreased cardiac output related to ventricular damage, ischemia and restriction secondary to fluid overload. Which would be an appropriate defining characteristic?

- a-Altered preload
- b-Altered afterload
- c-Poor ventricular contractility**
- d-Increased pulmonary congestion

49. After the nurse has administered a heparin injection. The patient is observed rubbing the area. The nurse should instruct the patient to not rub the area, as it may

- a- Increase the risk of bleeding.**
- b- Prevent proper absorption of the medication.
- c- Cause increased pain.
- d- Result in tissue death.

50. A home nurse attends an obese patient, with decreased mobility. Who has developed pain in her left calf muscle on examination, the leg appears swollen and red with large superficial veins. The nurse refers the patient for investigation and treatment (see lab results) :

Test	Result	Normal Values
Platelet count	49 1	40-400 x 10 ⁹ / L
INR	1.5	0.7 -1.2

What type of treatment is most appropriate for this patient?

- a-Unfractured heparin
- b-Anti-embolic stocking**
- c-Warm compresses and heating pads
- d-Anti-inflammatory medications

51. Which of the following medications would the patient be instructed to stop 2 weeks prior the surgery?

- a- Antiarrhythmic
- b- Antifungal
- c-Ant diabetic
- d-antiplatelet**

52. A patient is admitted to the emergency department with stable angina. The physician writes an order for nitroglycerin 1/150

grains sub lingually (SL). Which is the correct dose in milligrams (mg)?

a. 0.04 mg

b. 0.4 mg

c. 4 mg

d. 40 mg

53. The nurse is assisting a patient to ambulate in hall. The patient has a history of coronary artery disease (CAD) and had coronary artery bypass graft surgery (CABG) 3 days ago. The patient reports chest pain rated 3 on a scale of 0 (no pain) to 10 (severe pain). The nurse should first:

a- Determine how long it has been since the patient's last dose of aspirin

b- Obtain a chair for the patient to sit down

c- Assess the patient's radial pulse

d- Ask the patient to take several slow, deep breaths.

54. A 52 year-old man with congestive heart failure presents to the Emergency Department with rapid and irregular heartbeats, and feeling dizzy and light-headed. The attending physician verbally calls out an order to the nurse to administer digoxin 0.25 milligrams by intravenous injection. How will the nurse complete the paperwork for this order?


a- Write, sign and repeat order back to the physician

b- Ask the physician to write and sign the order

c- Write the order and ask the physician to sign

d- Verbally repeat the order and administer drug


55.



What is the most probable type of arrhythmia in the following ECG (lead II):

- a- Atrial fibrillation
- b- Atrial flutter
- c- Ventricular fibrillation
- d- Supra-ventricular fibrillation


56.



What is the most probable type of arrhythmia in the following ECG (lead II):

- a- Atrial fibrillation
- b- Atrial flutter
- c- Ventricular fibrillation
- d- Supra-ventricular fibrillation


57.



What is the most probable type of arrhythmia in the following ECG (lead II):

- a- Atrial fibrillation
- b- Atrial flutter
- c- Ventricular fibrillation
- d- Supra-ventricular fibrillation**

58.



What is the most probable type of arrhythmia in the following ECG (lead II):

- a- Atrial fibrillation
- b- Atrial flutter
- c- Ventricular Tachycardia**
- d- Supra-ventricular fibrillation

59.



What is the most probable type of arrhythmia in the following ECG (lead II):

- a- Atrial fibrillation
- b- Atrial flutter**
- c- Ventricular fibrillation
- d- Supra-ventricular fibrillation

60. Which of the following is the best position to insert central line catheter:

- a- Prone
- b- Supine

c-Semi- Fowler's

d-Trendelenburg

61.Toxicity from which of the following medications may cause a client to see a green-yellow halo around lights?

a- Digoxin

b- Furosemide (Lasix)

c- Metoprolol (Lopressor)

d- Enalapril (Vasotec)

62. A client who has been hospitalized after experiencing a heart attack will most likely receive a diet consisting of:

a-Low fat, low sodium, and high carbohydrates

b-Low fat, high protein, and carbohydrates

c-Low fat, low sodium, and low carbohydrates

63.which of the following site nurse should avoid to take temperature if the patient has heart attack?

a.oral

b.rectal

c.tympanic

d.axillary

64.Prior to initiating therapy with unfractionated heparin for a patient hospitalized with a deep vein thrombosis, the nurse should plan to:

a. Weigh the patient

b. Administer aspirin

c. Limit fluid intake

d. Undress the patient

65.After cardiac surgery, a patient has been prescribed low-sodium, low cholesterol diet. Which of the following menus is BEST?

a. Salami, rye bread, sauerkraut

b. Baked chicken thigh, iceberg lettuce, sliced tomatoes

c. Pasta with canned tomato sauce, peas, wheat bread

d. Bacon, lettuce and tomato sandwich with mayonnaise dressing

Nervous System

- 1. When conducting discharge teaching for a patient who recently suffered a Cerebrovascular accident (CVA). The nurse would MOST likely implement which of the following interventions to prevent muscle and ligament deformities?**
 - a. Daily moist heat and isometric exercises
 - b. Daily balance training and routine medications for pain
 - c. Instruct patient to use non-affected side to perform activities of daily living
 - d. Daily range of motion exercises**

- 2. patient admitted with a cerebrovascular accident (CVA), is unable to chew or swallowed. The patient is a risk for aspiration. The nurse would anticipate receiving which of the following orders for this patient?**
 - a. Give no food by mouth and start intravenous hydration**
 - b. Start a pureed diet with thickened liquids
 - c. Refer the patient to a psychiatrist for depression related to the CVA
 - d. Refer the patient to physical therapy for muscle strengthening

- 3. A patient had a craniotomy with resection of a nonmalignant neoplasm for the temporal lobe. The patient's vital signs are within the base line normal range. The nurse observes that the patient has developed bilateral per orbital edema. Which of the following actions would be appropriate for the nurse to take?**
 - a. Apply cold compresses to the patient's eyes
 - b. Apply warm compresses to the patient's eyes
 - c. Elevate the head of the patient's bed to 60 degrees
 - d. Elevate the head of the patient's bed to 45 degrees**

- 4. A patient had a Vitrectomy and is about to be transported to the post anesthesia care unit (PACU). The patient should be placed in which of the following positions before transport to the PACU?**
 - a. Semi-fowler's
 - b. Prone**

- c. Dorsal recumbent
- d. Sim's

5. A patient presents to the clinic with “pins and needles” sensations of the left foot and complains that objects appear “Shimmering”. The patient is diagnosed with optic neuritis and referred for further testing. The patient is MOST likely to be tested for:

- a. Glaucoma
- b. Multiple sclerosis**
- c. Lesion of brain stem
- d. Psychosis

6. The nurse is assessing a patient with a history of a seizure disorder. While checking the patient's vital signs, the patient develops rhythmic, jerking movements of the arms and legs. The nurse should IMMEDIATELY place the patient in which of the following positions?

- a. Prone
- b. Supine
- c. Semi-fowler's
- d. Lateral**

7. A nurse is assessing to care for a child with a seizure disorder. The nurse observes the child becomes stiff and lose consciousness, following by jerking movements for 1 minute after which the child becomes very sleepy, which of the following types of seizures occurred?

- a. Absence (petit mal)
- b. Generalized (tonic-clonic)**
- c. Partial Psychomotor(temporal lobe)
- d. Status epileptics

8. A community health nurse visits a patient who has suffered a stroke. The patient's spouse explains to the nurse that the patient chokes on foods at times. Which of the following referral ordered would the nurse anticipate needing for this patient?

- a. Speech therapist**

- b. Dietician
- c. Physician therapist
- d. Neurologist

9. As per of a neurological assessment, which of the following is associated with the higher score on the Glasgow coma scale?

- a. Eye opening to pain, no verbalization
- b. Confused, obey commands**
- c. Localized pain, abnormal extension
- d. Eye opening to speech confused

10. A patient with an unnecessary gait and a history of falls has a care plan intervention that includes keeping the walker in reach and pathway free of obstacle. On evaluation after 1 week, the patient has had no falls, but the gait remains unsteady. The nurse should:

- a. Continue the plan of care as written**
- b. Allow the patient to replace the walker with a cane
- c. Allow the patient to ambulate short distance without the walker
- d. Have the patient practice stepping over small objects

11. A 2-years-old child in the emergency department exhibits symptoms of bacterial meningitis. Which of the following tests confirm or rule out this diagnosed?

- a. Magnetic resonance imaging (MRI)
- b. Magneto encephalogram
- c. Computed tomography scan(CT)
- d. Lumbar puncture (LP)**

12. The nurse assists with a lumbar puncture on a child with suspect bacterial meningitis. If the diagnosis is correct, the cerebrospinal fluid, should have which of the following qualities?

- a. High glucose level
- b. Low protein level
- c. Cloudy or turbid appearance**
- d. Pink or blood-tinged appearance

13. A child in the postictal state of a seizure should show which of the following signs or symptoms?

- a. Feeling sleepy or exhausted
- b. Stiffness over entire body
- c. Verbalizes having an aura
- d. Eyes fixed in one position

14. The nurse is entering the room of a patient who is blind. The nurse should:

- a. Speak before touching the patient
- b. Talk to the patient using aloud tone of voice
- c. Ask then patient questions that can be answered “yes “or “no”
- d. Stand directly in front of the patient while talking

15. A patient is receiving from surgery using spinal anesthesia. The patient develops a spinal headache. Which of the following nursing actions would be MOST appropriate?

- a. Elevate the head of the bed 30 degrees
- b. Keep the patient well hydrated
- c. Limit intake of salty food
- d. Lower the temperature of the room

16. A nurse is caring for a patient who had rhinoplasty 2-weeks ago. Which of the following is an expected outcome?

- a. Oral mucus membranes dry ,but pink and intact
- b. Face and nose free from swelling
- c. Able to make needs know, speech therapy started
- d. Demonstrate throat clearing while eating

17. A patient presents to the emergency room with complaints of eye and drainage. In planning for the examination of the patients complaints, which of the following instruction would the nurse MOST likely select?

- a. Sphygmomanometer
- b. Thermometer
- c. Ophthalmoscope

18. A patient with conjunctivitis reports the presence of photophobia and moderate eye drainage. The nurse should teach patient to

- a. Avoid touching the eye
- b. Use sterile gauze to remove the drainage
- c. Darken the room**
- d. rest in the prone position

19. A patient visiting the clinic 10 days after sinus surgery for checkup complains of having a bad taste in the mouth. When the nurse smells a foul odor while examining the patient's mouth, the nurse suspects the patient has an:

- a. Pulmonary decompensation
- b. Hemorrhage
- c. Aspiration
- d. Infection**

20. A patient with diabetic retinopathy is experiencing an episode of unresolved hemorrhage in the eye. The nurse identifies the MOST likely procedure to benefit this patient would be:

- a. Enucleation
- b. Radial keratotomy
- c. Vitrectomy**
- d. Peripheral Iridectomy

21. A nurse is assigned to care for a patient with a diagnosis of thrombotic stroke. The nurse knows that this type of stroke is most likely caused by:

- a. Blockage of large vessels as a result of atherosclerosis**
- b. Emboli produced from Valvular heart disease
- c. Decreased cerebral blood flow due to circulatory failure
- d. A temporary disruption in oxygenation of the brain

22.A 34 year old quadriplegia patient resides at home with his wife. In order to prevent contractures of all extremities, the community care nurse will instruct the patient's wife in the performance of

- a. Active range of motion exercise
- b. Passive range of motion exercise**
- c. Active assistive range of motion exercise
- d. Resistive range of motion exercise

23.A child is treated for bacterial meningitis with an intravenous antimicrobial agent. Which of the following BEST indicates effectiveness of the treatment?

- a. Increased appetite
- b. Temperature 37.2 C(99 F)**
- c. Episodes of apnea
- d. Increased intra cranial pressure

24.The nurse is caring for a patient with Parkinson's disease. Which of the following is an expected outcome related to the nursing diagnosis of constipation related to diminished motor function, inactivity and medications?

- a. The patient will use a laxative every other day
- b. The patient will have a soft bowel movement daily**
- c. The patient will report minimal pain with bowel movements
- d. The patient will limit the intake of complex carbohydrates

25.Instrument which is used to examine the eye structure is called:

- a. Ophthalmoscope**
- b. Laryngoscope
- c. Otoscope
- d. Bronchoscope

26.While reviewing stress management techniques with a patient diagnosed with multiple sclerosis, what would the nurse identify as MOST appropriate?

- a- Relaxing in a warm bubble bath
- b- Yoga in a cool room**
- c- Sunbathing

d- Cross-country running

27. A 25 years-old male patient suffered a spinal cord injury at the T -4 level and is being cared for in hospital. The nurse enters the patient's room and finds the patient sitting upright and looking anxious and restless. He complains of sudden headache and nausea. Sweat forms on his forehead yet his feet are cool to touch

Blood pressure 150/100 mmhg

Heart rate 55/min

Respiratory rate 28/min

Temperature 37.1C

What nursing intervention is initially most appropriate?

a- Assess for a full bladder

b- Lower the head of the bed 30 degrees

c- Loosen clothing and bed sheets

d- Apply heating pad to lower extremities

28. A 24 year-old man was admitted to the hospital after receiving a closed head injury in a motor vehicle accident. He had a Glasgow Coma Score of 6 on admission and was being mechanically ventilated. The nurse observes the intracranial pressure has slowly and steadily increased over the past hour.

What is the most appropriate initial nursing action?

a- Administer mannitol

b- Administer Phenobarbital

c- Increase respiratory rate

d- Reposition while maintaining neck alignment

29. A 40 year-old man present to the clinic with a weight loss of three kilograms over the past month. His health problem produces moderate spastic body movements, hyper-reflexia, muscular contractions followed by relaxation, and a positive Babinski reflex. He has difficulty speaking due to poor muscle tone and has recently developed swallowing difficulties. The nurse creates a care plan to address his developing problem. Which is the priority nursing intervention?

a- Increase caloric intake

b- Aspiration prevention

c- Infection control

d- Improved muscle control

30. A brick-layer has been diagnosed with multiple sclerosis. The nurse knows that construction work places has increased risk of aggravated symptoms because exposure to.

- a- Dust**
- b- Sunlight
- c- Heat
- d- Heights

31. Patient experienced (CVA) one week ago. The doctor order the patient discharged from the hospital to home health care. The visiting nurse evaluates the patient and notes the patient has difficulty initiating speech and forming words. Which of the following types of aphasia is most likely in this patient?

- a- Receptive
- b- Expressive**
- c- Fluent
- d- Conduction

32. A patient had a retinal detachment surgically repaired the nurse identifies that the detachment would most likely be corrected and unlikely to reoccur if the retina remains attached at least:

- a- Three days
- b- Two weeks**
- c- Two months
- d- Three months

33. A patient complains of incoordination, weakness and fatigue, double vision, and slurred speech laboratory test results are normal except for magnetic resonance imaging (MRI) scan which shows small plaques scattered throughout the white matter. The nurse would anticipate the diagnosis of:

- a- Multiple sclerosis**
- b- Brain metastasis
- c- Amyotrophic lateral sclerosis (Lou Gehrig disease)
- d- Polyradiculoneuritis(Guillain-Barre syndrome)

34. An 82 year-old patient has Parkinson's disease. During the assessment, the nurse would expect which of the following actions to produce the MOST tremor activity of the hands?

- a- Eating with a fork
- b- Resting hands in lap**
- c- Standing with hands loose at sides
- d- Rolling a small pill between the fingers

35. Which of the following is the best position to increase the brain perfusion (cerebral perfusion):

- a- Prone
- b- Supine
- c- Semi- Fowler's
- d- Trendelenburg's**

36. A 34 year- old quadriplegia patient resides at home with his wife. In order to prevent contractures of all extremities, the community care nurse will instruct the patient's wife in performance of:

- A. Active range of motion exercises .
- B. Passive range of motion exercises .**
- C. Active- assistive range of motion exercises .
- D. Resistive range of motion exercises .

37. Following a tympanoplasty, the nurse should maintain the client in which position:

- a- Semi-Fowler's with the operative ear facing down
- b- Low Trendelenburg's with the head in neutral position
- c- Flat with the head turned to the side**
with the operative ear facing up
- d- Supine with a small neck roll to allow for drainage

38. A nurse is providing care for a client following surgery to remove a cataract from the right eye. In which position should the nurse place the client?

- a- Right-side lying
- b- Supine**
- c- Prone
- d- Trendelenburg's

39. An adult is admitted for a neurological workup and is scheduled for a spinal tap. When preparing the client for the procedure, the nurse should position the client in which position:

a-Prone

b-On the side with knees

drawn up to chest

c-Lithotomy

d-Semi-sitting

40. If a patient develops autonomic hyperreflexia the first action his caring nurse should take is to:

a-Elevate head of bed 90 degree

(move from supine to sitting)

b-Make the bed in flat position

c-Apply ice on the axillary and groin

d-But the patient in trendelenburg position

41. When caring for a patient admitted post-stroke (CVA) who has altered consciousness, the nurse should place the patient in which position:

a-Side-lying

b-Prone

c-Supine

d-Semi-fowler's

42. A nurse is providing instructions to a client and the family regarding home care after right eye cataract removal. Which statement by the client would indicate an understanding of the instructions:

a-“I should not sleep on my left side.”

b-“I should not sleep on my right side.”

c-“I should not sleep with my head elevated.”

d-“I should not wear my glasses at any time.”

43. When assessing for therapeutic effects of mannitol, the nurse would expect to see:

a- Decreased intracranial pressure

b- Decreased excretion of therapeutic medications

c- Increased urine osmolality

d- Decreased serum osmolality

44. Which drug would be used to treat a patient who has increased intracranial pressure (ICP) resulting from head trauma after an accident:

a- Mannitol

b- Atropine sulfate

c- epinephrine hydrochloride

d- Sodium bicarbonate

45. A client with myasthenia gravis is suspected of having cholinergic crisis. Which of the following indicate that this crisis exists:

a- Ataxia

b- Hypotension

c- Mouth sores

d- Hypertension

46.36 years old male patient complaining of fever and headache for 3 days now, when examining this patient which of the following signs would indicate that this patient may has meningitis:

a-Positive Kernig's sign

b-Negative Brudzinski's sign

c-Positive homan's sign

d-Negative Kernig's sign

47.A patient presented with high fever, headache, vomiting and neck stiffness for the past 3 days, which of the following is the first diagnostic intervention for this patient:

a-Urine and stool analysis

b-Lumber puncture with CSF aspiration

c-Complete blood count

d-Chest and abdomen x-ray

48.The nurse recognizes that the most common type of brain attack (CVA) is related to which of the following:

a-Ischemia

b- Headache

c-Hemorrhage

d-Vomiting

49. A patient visits the clinic for a 2- week checkup after a corneal transplantation (keratoplasty). The nurse observes the patient's sclera is red and the patient complains of the eye feeling irritated. The nurse suspects the patient may have:

- a-Infection
- b-Hemorrhage
- c-Graft rejection**
- d-Postoperative glaucoma

50. A 32 year- old woman with a motor neuron disease has progressive weakness of cranial nerves V, IX, XII. She has recently been experiencing sudden outbursts of crying and laughing. Both stroke and myasthenia gravis have been ruled out. Which voluntary muscle activity would be most significantly impacted?

- a-Walking
- b-Swallowing**
- c-Breathing
- d-Smelling

51. Adult patient suffers from hoarseness of voice, and difficulty in speech may be suffering from an injury of which of the following cranial nerves:

- a-IX
- b-X**
- c-XI
- d-XII

52.The nurse is performing an assessment in a client with a suspected diagnosis of cataract. The chief clinical manifestation that the nurse would expect to note in the early stages of cataract formation is:

- a-Eye pain
- b-Blurred vision**
- c-Floating spots
- d-Diplopia

53.A nurse examines a client's level of responsiveness. She finds that the patient opens his eyes to verbal commands, obeys verbal commands, and is oriented to time, place, and person. What's the client's Glasgow Coma Scale:

- a-11/15
- b-12/15
- c-13/15
- d-14/15**

54.A 73 year old patient looks at you when you speak to her. When you ask her the date, she says "blue." You note left-sided weakness when she grips your fingers. What is her Glasgow Coma Score?

- a-11/15
- b-12/15**
- c-13/15**
- d-14/15

55.A patient during examination opens his eyes in response to pain, makes no verbal response, but withdraws from pain. What is the Glasgow Coma Score (GCS) for this patient?

a-3

b-7

c-5

d-11

56. Which nursing diagnosis takes priority for a newly diagnosed patient with a left- sided stroke?

a-Risk of impaired swallowing related to
absent gag reflex

b-Risk for impaired skin integrity related
to immobility

c-Risk for infection related to
invasive line placement

d-Risk for impaired speech related to
left sided stroke

57. Changes in personality and judgment are often associated with a lesion in which of the following:

a-Frontal lobe

b-Parietal lobe

c-Broca's area

d-Wernicke's area

58. Anterior lobe of the brain is responsible for which of the following:

a- Personality

b- Movement

c- Speech

d- Memory

59. The nurse is observing a detailed neurological assessment on a client with a suspected brain tumor. When performing the Romberg test, the client sways when the eyes are both open and closed. What does this indicate?

a- The problem is probably in the cerebellum

b- It is a position sense abnormality

c- This is not an abnormal test result

d- The client has lost proprioception

60. A client with multiple sclerosis has been prescribed the drug baclofen (Lioresal). What is the action of this drug?

a- Reduces spasticity

b- Skeletal muscle relaxation

c- Immune suppression

d- Prevents viral infections

61. After X-rays are taken of the head, neck, and spine, the client is diagnosed with a head injury and admitted for inpatient care. When assessing the client with a head injury, which of the following should receive priority attention?

- a- Lung sound
- b- Clarity of speech
- c- Mobility of fingers

d- Pupillary responses

62. An elderly client is experiencing an alteration in his equilibrium and coordinated muscle movements. The nurse realizes that these functions are controlled by which area of the nervous system?

- a- Brain stem
- b- Cerebrum
- c- Diencephalon

d- Cerebellum

63. A nurse evaluates a patient for signs of rebleeding from ruptured intracranial aneurysm that required surgical ligation. The highest risk for aneurysm rebleed is within:

- a. 6 hours
- b. 24 hours
- c. 48 hours

d. 72 hours

Musculoskeletal & hematology & heredity

1. In evaluating the appropriateness of various exercises enjoyed by a patient with osteoporosis, the nurse would recommend:

- a. Walking**
- b. Bowling
- c. Sit-ups
- d. Golf

2. A hospitalized patient has fallen from bed. The nurse notes shortening of the left leg . Pain upon movement of the left leg, and rapid, shallow respirations. What action should the nurse take FIRST?

- a. Call for help
- b. Immobilize the left leg**
- c. Obtain blood pressure
- d. Evaluate lung sounds

3. While caring for a patient prior to surgery to amputate the leg. What is the MOST effective measure to prevent phantom limb sensation after the amputation?

- a. Control pain prior to the surgery
- b. Make sure the patient understands the procedure
- c. Elevate the limb on two pillows
- d. Help the patient grieve for the limb**

4. A marathon runner experiences a sudden onset of sharp pain in calf immediately after a workout. The nurse in the clinic notes mild swelling of the calf and tenderness to touch. Which of the following would the nurse suspect the patient is experiencing?

- a. Bursitis
- b. Tendonitis
- c. Plantar fasciitis**
- d. Joint dislocation

5. A nurse is evaluating a patient 5 days after a right total hip replacement. Which of the following goals is appropriate for the patient?

- a. Maintain hip abduction without dislocation**
- b. Rest with legs elevated while sitting

- c. Tie shoes and put on undergarments without assistive devices
- d. Perform scissors-like leg exercise daily

6. A patient with bowlegs due to abnormal bone formations and deformities has a calcium level of 7.5 mg/100 ml. Which of the following foods would the nurse MOST likely instruct the patients to add to a diet?

- a. Organ meats
- b. Whole grains
- c. Egg yolks
- d. Lean meats

7. A nurse is caring for a 3 year-old child with a fractured arm. Which of the following interventions is the MOST appropriate for pain management?

- a. Administer analgesics when necessary
- b. Assess pain once a shift
- c. Anticipate pain and intervene early
- d. Encourage the use of self-quieting techniques

8. A patient has a history of severe, uncontrolled epistaxis . The patient's blood pressure and patient count are normal. The nurse should teach the patient to:

- a. Sleep with the head elevated on at least two three pillows
- b. Apply firm pressure to the nostrils four times a day
- c. Use a cotton-filled applicator to apply a water-soluble lubricant to the nasal septum twice daily
- d. Minimize the intake of caffeine while increasing the intake of fluids rich in vitamin K

9. The following pain medications are ordered for a patient who had a right leg debridement . Oxycodone 5 mg every 4 hours as needed and morphine 5 mg every 4 hours as needed. The nurse administered oxycodone 2 hours ago, but the patient report pain Rated 8 on a scale of 0 (no pain) to 10 (Severe pain) as the dressing change begins .Vital signs are: blood pressure level, 169/98 mmHg; heart rate, 112; Respiration rate 22; temperature 36.7 C (98.1 F).After evaluating the effectiveness of the pain Medication, what action should the nurse take?

- a. Administer additional oxycodone 5 mg
- b. Administer morphine 5 mg**
- c. Change the dressing quickly
- d. Encourage deep breathing

10. A patient in a long-term care facility is in persistent vegetative state with a right contracture of the right arm and hand. What is the BEST goal over the next 90 days for this patient related to the nursing diagnosis of impaired mobility?

- a. Develop no further contractures**
- b. Wear an arm and hand splint
- c. Have no pain related to the contractures

11. A patient is scheduled for a total hip arthroplasty. The preoperative nurse reviews the chest and notes the following: serum potassium level of 2.8 mEq/l, AB positive blood type, and elevated ST Segments on the electrocardiogram (ECG). Which of the following would be the MOST appropriate action for the nurse to do next?

- a. Report abnormal diagnostic results to the surgeon**
- b. Review the patient consent for the surgical procedure
- c. Educate the patient on the risk factors and side-effects of the surgery
- d. Ensure that the patient has a post-surgery physical therapy order

12. A 45-year-old patient is in a lower body cast following a motor vehicle accident. In order to minimize muscle strength loss while in the cast, the nurse will instruct the patient in the performance of:

- a. Isometric exercises**
- b. Passive range of motion exercises
- c. Active-assistive range of motion exercises
- d. Resistive range of motion exercises

13. After total knee replacement a patient being discharged to have after which he will ambulate with for-prong cane. When providing patient teaching regarding giving up and down stairs with the cane, the first step in going up stairs is to :

- a. Place the cane and the affected extremities upon the step
- b. Place the cane and the unaffected extremity upon the step
- c. Step up on the affected extremity
- d. Step up on the unaffected extremity**

14. A patient with SLE (systemic lupus erythematosus) report decreased urinary output during the past 2-4 days and chest pain that is aggravated by breathing and coughing. The patient vital signs remain within the baseline normal range s1 and s2 are present with audible friction rub. Which of the following statement would be appropriate for the nurse to make?

- a. It sounds like SLE is being well controlled
- b. I need to get some nitroglycerine for your chest pain
- c. There may be some inflammation surrounding your heart**
- d. Your symptoms may be due to a urinary tract infection

15. A diabetic patient comes to the office for follow-up six weeks undergoing below the knee amputation of the right leg for gangrene. The nurse observes that the patient is progressing well with the use of prosthesis and that the skin is intact. The patient reports being generally pain free but occasionally feels severe pain and itching of the right ankle. What should the nurse do?

- a. Notify the doctor that there appears to be nerve damage of the right leg
- b. Refer to pain management specialist for long term management
- c. Refer to psychiatrist for evaluation since the patient has no right ankle
- d. Explain the phenomena of phantom pain and phantom sensation to the patient**

16. A 40 year- old woman presented with right hip pain. Palpation of the pelvic girdle is normal. An X- ray shows bone deformities, with osteolytic lesions and bone enlargement. The patient has not suffered any trauma and has been generally healthy. Which serum laboratory analysis would be most useful?

- a. Prothrombin time
- b. Alkaline phosphatase (if this high, calcium will be low and opposite)**
- c. Acid phosphatase
- d. Parathyroid hormone

17. A patient who is 4 days postoperative after a total hip replacement surgery, is obese and has not been able to ambulate since the surgery. The patient is now diaphoretic, has chills, and complains of pain in the thigh. There is tenderness over the anteromedial surface of the thigh. The MOST likely cause is:

- a. Wound infection**

- b. Deep vein thrombosis (DVT)
- c. Pulmonary edema
- d. Dehydration

18. A 22 year-old man suffered major injuries to the right leg in a road traffic accident that resulted in an above -the- knee amputation with a rigid dressing. Twelve hours later the patient complains of increasing pressure from the dressing and the feeling that something is crawling around the stump.

- a- Notify the physician
- b- Remove and reapply the dressing
- c- Remind patient phantom limb pain is normal**
- d- Elevate and apply ice to the residual limb

19. A nursing home resident with chronic osteoarthritis complains of knee pain. A routine order of acetaminophen (Tylenol) 500 mg every 6 hours was administered 2 hours ago with no relief. The patient also has an order for ibuprofen (Motrin) 400 mg every 6 hours as needed. The patient reports some relief, but is still bothered by mild pain.

The nurse should

- a- Administer ibuprofen (Motrin) 400mg**
- b- Educate the patient that mild pain is expected with osteoarthritis
- c- Call the physician for additional pain medication
- d- Administer acetaminophen (Tylenol) 500 mg

20. A 40 year-old woman present with painful leg cramps and muscle twitching, which wakens her at night. Her menstrual cycles are irregular and cause painful abdominal cramping. The hair, skin and nails appear dry and brittle (see lab results)

Test Result Normal Values

Calcium 2.09 2.15-2.62mmol/L

Phosphate, inorganic 1.69 0.82-1.51mmol/L

Magnesium 0.55 0.7-1.0mmol/L

What type of lifelong diet would be most beneficial for this patient?

- a- Increased magnesium and vitamin C
- b- Increased calcium and vitamin D**
- c- Increased dairy and iron
- d- High protein and high calorie

21. A 35 year-old man twisted his right ankle. Since then, he has had swelling and stiffness of the ankle in the morning hours. He presented to the clinic six months after the injury and complains that he is in too much pain to participate in sports. An arthroscopy

reveals thickening and inflammation of the lining of the ankle joint. Which treatment is most likely to be ordered for this client?

- a- Aspiration of fluids
- b- Surgical intervention
- c- Physical therapy
- d- NSAID administration

22. A 67 year-old man was admitted to the hospital following a closed bone fracture. An intramedullary nail is inserted and the patient is placed in balanced skeletal traction. The following day, the patient becomes restless, drowsy and confused, he has difficulty breathing and appears very tired. Which additional sign or symptom would require immediate intervention?

- a- Anxiety
- b- Cold skin
- c- Constipation
- d- Petechiae on chest

23. A 12 year- old patient had a cast removed from the left leg after wearing it for eight weeks. The patient wants to resume sports as soon as possible. In order to regain muscle strength lost while wearing cast, the nurse will instruct the patient in performance of:

- a- Resistive range of motion exercises to left leg
- b- Passive range of motion exercises to right leg
- c- Active- assistive range of motion exercises to the right leg
- d- Active range of motion exercises to both legs

24. RTA. The patient appears restless confused and disoriented. He reports that he had hit his head against the steering wheel of the car when it had collided with the car directly in front of him.

Blood pressure 110/68 mmHg

Heart rate 100/min


Respiratory rate 22/min

Body Temperature 37.0 coral

Oxygen saturation 98 % on room air

What is the most important next step in management?

- a- Immobilize head and neck
- b- Administer oxygen.
- c- Establish an intravenous line.
- d- Arrange for an MRI scan.



25. which of the following anatomical terms best describes the motion in the picture above:

a-Adduction

b- Abduction

c- Supination

d- Pronation

26. The following picture represents an auto immune disease known as:

a-Psoriasis

b-Systemic Lupus Erythematosus

c- Molluscum Contagiosum

d-Eczema

27. A client has just returned to a nursing unit after an above-knee amputation of the right leg. A nurse places the client in which position:

a-Prone

b-Reverse Trendelenburg's

c-Supine, with the amputated limb flat on the bed

d-Supine, with the amputated limb supported with pillows

28. After the client undergoes a total hip replacement, how should the nurse position the affected hip?

a-Adduct the hip

b-Flex the hip

c-Abduct the hip

d- Extend the hip

29. A client has an open reduction and internal fixation for a fractured hip. Postoperatively the nurse should position the client's affected extremity in:

a-External rotation

b-Slight hip flexion

c-Moderate abduction

d-Anatomical body alignment

30. The nurse is caring for a client with an above-the-knee amputation (AKA). To prevent contractures, the nurse should:

a-Place the client in a prone position 15–30 minutes twice a day

b- Keep the foot of the bed elevated on shock blocks

c-Place trochanter rolls on either side of the affected leg

d-Keep the client's leg elevated on two pillows

31. The nurse is caring for a client who had a right below-the-knee amputation three days ago. The client complains of pain in the right foot and asks for pain medication. What nursing action is appropriate initially?

a- Elevate the stump

b- Administer a placebo

c- Administer ordered medications

d- Encourage the client to discuss his feelings

32. During the assessment of an injury for a possible fracture, which of the following manifestations would make you suspicious that a fracture has occurred:

a-Impaired sensation

b-Loss of function

c-Hotness

d- Pointed tenderness

33.How you will interfere to relieve swelling of the lower extremity in a cast for 3 days now:

a- Change the cast

b- **Elevate the extremity**

c- Call the doctor

d- Give massage

34.Which of the following interventions should be taken to help an older client to prevent osteoporosis:

a-Decrease dietary calcium intake

b- Increase sedentary lifestyles

c- Increase dietary protein intake

d- Encourage regular exercise

35.Which type of medications is commonly used to treat rheumatoid arthritis?

a-Glucocorticoids

b-Non- steroidal antiinflammatory drugs (NSAIDs)

c- Antimalarial drugs

d- Gold salt

36.Clients who have casts applied to an extremity must be monitored for complications. The most significant complication for which the nurse should assess the client's extremity is:

a. Warmth

b. **Numbness**

c. Skin desquamation

d. Generalized discomfort

37. A 55 year- old woman presented with pain in the joints of the hands. She feels generally tired and has had a low grade fever for the past week. On examination, the joints of the fingers on both hands are symmetrically swollen.

Blood pressure	120/70 mmhg
Heart rate	82/min
Respiratory rate	18/min
Temperature	38.0 C
Oxygen saturation	99% on room air

Which type of arthritis is most likely?

- a- Rheumatoid**
- b- Septic
- c- Psoriatic
- d- Osteoarthritis

38. A seven year- old child presented to the emergency room with a fracture of the right arm.

What would be the long- term goal for this patient?

- a- Patient verbalizes decreased pain level
- b- Nurse notes neurovascular checks are within normal limits
- c- Nurse notes decreased swelling of the right
- d- Patient performs activities of daily living without difficulty**

39. A 35-years-old female has an inherited gene mutation for achondroplasia, an autosomal dominant genetic disorder. Her husband does not have gene mutation. In planning genetic counseling for this patient, the nurse would be MOST correct in including which of the following statements regarding the risk of their children inheriting the genetic mutation?

- a. Each child has a 50% chance of inheriting the gene mutation**
- b. Female children have 50% chance of inheriting the gene mutation
- c. Male children will not inherit the gene mutation
- d. All female children will inherit the gene mutation

41. If a patient develops a complication during a blood transfusion, the nurse's first action should do to:

- a. Stop the transfusion**
- b. Notify the practitioner
- c. Administer an antihistamine
- d. Administer an anti-inflammatory medication.

42. A child recently diagnosed with sickle cell anemia is being prepared for discharge. Which of the following statement by one of the parents would require ADDITIONAL teaching by the nurse?

- a. High altitudes can be beneficial**
- b. Blood transfusion may be necessary in the future
- c. Strenuous physical activity should be avoided
- d. Increased fluid intake minimize pain

43. A patient is being evaluated due to onset of paleness, shortness of breath and sensations of heart palpitations. Which of the following component of complete blood count (CBC) should the nurse review to determine if the patient has anemia

- a. Leukocytes
- b. Platelets
- c. Erythrocytes**
- d. Thrombocytes

44. A home care nurse visits a patient who is discharged from a hospital after a treatment of urosepsis. Which of the following post discharge normal laboratory result BEST indicates desired outcome?

a. WBC count

- b. Hematocrit
- c. Platelet level
- d. Potassium level

45. A home care nurse reviews the laboratory results for a postpartum patient who had a caesarean section . Which of the following indicates possible wound infection ?

a. Increased WBC

- b. Decreased hematocrit level
- c. Increased hemoglobin
- d. Decreased platelet

46. Three days ago a patient underwent an invasive surgery with an open wound. The patient is febrile with drop in blood pressure. Laboratory test results shows elevated WBC count. This could be possible presentation of :

a. Sepsis

- b. Atelectasis
- c. Internal hemorrhaging
- d. Excess fluid volume

47. A patient who is receiving chemotherapy has a platelet count of 49,000/mm³ (normal value 150,000 to 400,000/ mm³). Which of the following nursing action is necessary?

a. Minimize invasive procedure

- b. Crush oral medications
- c. Limit intake of vitamin K rich foods
- d. Monitor the temperature every 4 hours

48. A 55 year-old man presented to the clinic with complaints of numbness and tingling in his feet for the past 3 months. He appears thin and his skin is pale. Examination confirms the loss of vibration sense. He also has altered proprioception. He had a subtotal gastrectomy two years ago (see lab results) Test

result	normal values	
MCH	2.72	1.45-2.01 fmol/cell
HCT	0.31	0.41-0.50
MCV	105	81-97 fl
Reticulocyte count	1.5	0.5%-2.5%

Which supplement would most likely improve his condition?

- a-Iron
- b- Folate
- c-B12**
- d-Calcium

49. Best diagnostic test for suspected leukemia is:

- a-Bone marrow aspiration**
- b-Blood chemistry
- c-CBC
- d-PT AND PTT

50. The primary nursing goal when caring of a child with leukemia is to:

- a-Take vital signs
- b-Provide good nutrition
- c-Give analgesics
- d- Prevent infection**

51. The nurse is discussing dietary sources of iron with a client who has iron deficiency anemia. Which menu, if selected by the client, indicates the best understanding of the diet?

- a- Milkshake, hot dog, and beets
- b- Beef steak, spinach, and grape juice**
- c- Chicken salad, green peas, and coffee
- d- Macaroni and cheese, coleslaw, and lemonade

52. The Nurse examines a newborn immediately following delivery and finds single palmar crease on the right hand, upward eye squinting, loos skin at the base of the neck and a heart murmur, the baby appears lethargic and is unable to successfully latch to the breast to feed.

Which genetic abnormality is most likely?

- A. Absent genetic material
- B. excess genetic material**
- C. X-linked recessive gene
- D. Autosomal recessive gene

53. The patient provides genetic counseling to the patients of a newborn infant girl who has an X-linked recessive disorder in which there a disruption of the normal intrinsic coagulation. The nurse the parents of the chances of this infant giving birth to a child with disorder when she becomes on adult. If the husband is not a carrier what re the chances of the boys having disorder?

- A. 0%
- B. 25%**
- C. 75%
- D. 100%

Endocrine system & Integumentary system

- 1. While evaluating the nutritional intake of a bedridden patient with multiple pressure sores, the nurse should make sure the patient INCREASES the intake of which of the following ?**
 - a. Protein-rich foods**
 - b. Water
 - c. Food rich in vitamin A
 - d. A Fiber rich foods

- 2. Which of the following can be used to determine if a prescribed bed sore ulcer for a bedridden patient?**
 - a. Papanicolaou test
 - b. Faces rating scale
 - c. Braden's scale**
 - d. Apgar assessment tool

- 3. A child is treated for superficial (first-degree) thermal burns to the thigh. The child is in great discomfort and does not eat. Which of the following diagnosis should receive PRIORITY?**
 - a. Altered nutrition
 - b. Impaired skin integrity
 - c. Risk for infection
 - d. Acute pain**

- 4. A nurse is providing care to a patient with a new skin graft on left leg. The patient is upset and the nurse notes copious red drainage oozing around the dressing. The nurse should immediately:**
 - a. Lift the dressing to assess the area
 - b. Ask if the patient is having any pain
 - c. Apply firm pressure for 10 to 15 minutes**
 - d. Assess the apical pulse

- 5. A 70 year-old quadriplegic man has ulceration on the left buttocks with full thickness skin loss. Although several interventions have**

been used to promote healing, the ulcer does not improve Which dietary insufficiency is most likely?

- a- Vitamin D
- b- Calcium
- c- Protein**
- d- Vitamin C

6. A patient who sustained extensive abdominal injuries in a motor vehicle accident has developed a large stage II pressure ulcer on the coccyx. A new diagnosis of alteration in skin integrity is added to the care plan. What is the BEST short-term goal for the patient?

- a- Show evidence of healing within one week
- b- Have no discomfort from the pressure ulcer
- c- Eat at least 50% of each meal
- d- Verbalize strategies to prevent further skin breakdown**

7. While providing care to a hospitalized child with third-degree burns over 20% of the body, the child rapidly becomes disoriented. The nurse knows this is an indicator of:

- a- Septic shock
- b- Separation anxiety
- c- Dehydration**
- d- Depression

8. A nurse is caring for a client with a severe burn who is scheduled for an autograft to be placed on the lower extremity. The nurse develops a postoperative plan of care for the client and includes which of the following in the plan:

- a-Maintain the client in a prone position
- b-Elevate and immobilize the grafted extremity**
- c-Maintain the surgical extremity in a flat position
- d-Keep the surgical extremity covered with a blanket

9. the patient have an itchy, erythematous, vesicular, weeping, and crusting patches on his skin which skin

disease is this considered:

- a-Psoriasis
- b-Impetigo
- c-Eczema**
- d-Urticaria

10. Which vitamin deficiency is most likely to be a long-term consequence of a full-thickness burn injury?

- a-Vitamin A
- b- Vitamin C
- c-Vitamin B
- d- Vitamin D**

11. Patient obtained second degree burn to his abdomen and his entire back, according to rule of nine what is the percentage of burn of total body surface area:

- a- 9%
- b- 18%
- c- 27%**
- d- 36%

12. The type of burn in which all the dermis and epidermis, is destroyed and there is involvement of underlying structures is called:

- a-Superficial or first degree burn
- b-Partial thickness or second degree burn
- c-Full-thickness or third degree burn**
- d- Fourth degree burn

13. Your patient's finger stick blood sugar at 07:30 is 300 mg/dl, according to the following sliding scale; how many unites you would give your patient:

- a. 6 i.u
- b. 8 i.u
- c. 14 i.u
- d. 18 i.u

FSBS Glucose level	Amount of regular insulin
0 - 150 mg/dl	0 units
151 - 250 mg/dl	8 units
251 - 350 mg/dl	14 units
351 - 400 mg/dl	18 units
> 400 mg/dl	Call MD

14. The following condition may develop in females due to an increase in the production of which of the following hormones:

- a-Androgens
- b-Progesterone
- c-Estrogen
- d-Growth



15. A client receiving long term treatment with high dose corticosteroids which of the following would the nurse expected the client to exhibit:

- a-Weight loss
- b-Pale thick skin
- c-Hypotension
- d-Moon face

16. While assessing a client with Hypoparathyroidism, the nurse taps the client's facial nerve and observes twitching of the mouth and tightening of the jaw. The nurse would document this finding as which of the following:

- a-Positive Trousseau's sign
- b-Positive Chvostek's sign**
- c-Tetany
- d-Hyperactive deep tendon reflex

17. Which of the following clinical manifestations of type 2 diabetes occurs if glucose levels are very high:

- a-Hyperactivity
- b-Blurred vision**
- c-Oliguria
- d-Increased energy

18. Which of the following would the nurse expect to find in a client with severe hyperthyroidism:

- a-Tetany
- b-Buffalo hump
- c-Exophthalmos**
- d-Striae

19. After undergoing a thyroidectomy, a client develops hypocalcemia and tetany. Which electrolyte should the nurse anticipate administering:

- a-Sodium phosphorus
- b-Sodium bicarbonate
- c-Calcium gluconate**
- d-Potassium chloride

20. The nurse who elicits a positive Chvostek's sign would suspect that the patient has which condition:

- a-Hyperkalemia
- b-Hypocalcemia**
- c-Hypercalcemia

d-Hyponatremia

- 21. Which of the following diets would be most appropriate for a patient has Cushing syndrome:**
- a-High protein diet
 - b-High potassium diet**
 - c-High calcium diet
 - d-High iron diet
- 22. When developing a plan of care for a patient with SIADH (syndrome of inappropriate antidiuretic hormone secretion), which interventions will the nurse include:**
- a-Encourage fluids to 2000 ml/day
 - b- Long-term fluid restriction**
 - c-Monitor for increased peripheral edema
 - d-Keep head of bed elevated to 30 degrees
- 23. A client is admitted for treatment of the syndrome of inappropriate antidiuretic hormone (SIADH). Which nursing intervention is appropriate?**
- a-Infusing I.V. fluids rapidly as ordered
 - b-Administering glucose-containing I.V. fluids as ordered
 - c-Restricting fluid**
 - d-Encouraging increased oral intake
- 24. A hospitalized client is found to be comatose and hypoglycemic with a blood sugar of 50 mg/dL.**

Which of the following would the nurse do first?

- a-Infuse 1000 mL D5W over a 12-hour period
- b-Encourage the client to drink orange juice with added sugar
- c-Check the client's urine for the presence of sugar and acetone
- d-Administer 50% glucose intravenously**

25. A woman comes to the clinic and states, "I've been sick for so long! My eyes have gotten so puffy, and my eyebrows and hair have become coarse and dry." The nurse will assess for other signs and symptoms of:

- a-Cachexia
- b-Parkinson's syndrome
- c- Myxedema**
- d-Scleroderma

26. Causes of primary hypothyroidism in adults include:

- a-Malignant or benign thyroid nodules
- b-Surgical removal or failure of the pituitary gland
- c-Surgical removal or radiation of thyroid gland
- d-Autoimmune-induced atrophy of the gland**

27. A 72-year-old patient is diagnosed with hypothyroidism, and levothyroxine (Synthroid) is prescribed. During initiation of thyroid replacement for the patient, it is most important for the nurse to assess:

- a-Mental status
- b-Nutritional level
- c-Cardiac function**
- d-Fluid balance

28. A client presents with Hypocalcemia, hyperphosphatemia,

muscle cramps, and positive Trousseau's sign. What diagnosis does this support?

a-Diabetes insipidus

b-Conn's syndrome

c-Hypoparathyroidism

d-Acromegaly

29. The nurse caring for a 54-year-old patient hospitalized with diabetes mellitus would look for which of the following laboratory test results to obtain information on the patient's past glucose control:

a-Prealbumin level

b-Urine ketone level

c-Fasting glucose level

d-Glycosylated hemoglobin level

30. Which laboratory test is most important for the nurse to monitor to determine how effectively the client's diabetes is being managed?

a-Fasting blood glucose

b-Blood chemistry profile

c-Complete blood count

d-Glycosylated hemoglobin (HbA1c)

31. HbA1c as a measurement instrument for glucose level in the blood, it measures blood glucose level within:

- a-Last 3 months
- b-Last 7 months
- c-Last 5 months
- d-Last 1 year

32. A home care nurse visits a diabetic patient who was started on insulin injections. Upon examination, the nurse observes small lumps and dents on the right upper arm where the patient has injected insulin. What is the BEST nursing intervention?

- a-Refer the patient to a dermatologist for diabetic cellulitis
- b- Instruct the patient to rotate the sites of injection
- c-Refer the patient to an endocrinologist for better control of glucose level
- d-Instruct the patient to inject in the muscular area instead of the subcutaneous area

33. NPH is an example of which type of insulin:

- a-Short-acting
- b-Intermediate-acting
- c-Rapid-acting
- d - Long-acting


34. A nurse is assessing a client with possible Cushing's syndrome. In a client with Cushing's syndrome, the nurse expects to find:

- a-Weight gain in arms and legs

b-Thick, coarse skin

c-Hypotension

d-Deposits of adipose tissue in the trunk and dorsocervical area



35. A patient has a blood glucose level of 60mg/dl. The patient is to receive 15 g of carbohydrate if the blood glucose is less than 70mg/dl. How many mls of orange juice should the patient receive?

a-120 ml

b-90 ml

c-60 ml

d-30 ml

36. A patient has a blood glucose level of 60mg/dl. The patient is to receive 15 g of carbohydrate if the blood glucose is less than 70mg/dl.

How many candies should the patient receive?

a-4candies

b-6candies

c-8candies

d-10candies

37. The nurse cares for 60 years-old women who has a history of hypertension, hypothyroidism and elevated cholesterol levels, she takes tablets daily for each of the health problem. The doctor orders a routine dual-energy X-ray absorptiometry test that shows decreased bone density. Which medication most likely contributed to the test result?

A-Statins

B-Anti-hypertensive

C-Synthetic thyroid hormones

D-Cholesterol absorption inhibitors

38. A 20 years-old woman presents to the emergency department with confusion, agitation, and fecal incontinence. Despite constantly being hungry and eating, she has lost 5 kg in the past two weeks and has a body mass index of 17 (see lab results).

Test	Result	Normal values
Free thyroxine	150	50-140 nmol/L
Free Triiodothyronine T3	5.5	2-4
TSH	0.1	0.4-6.5 mU/L

Which additional finding would indicate she requires immediate care?

A. Polyuria, polydipsia and severe headache

B. Hypotension, bradycardia and cold clammy skin

C. Cold Clammy skin, flushing and tachypnea

D. Fever, tachycardia and hypertension

39. A 30 year-old football player has a health problem which has been caused by the destruction of the posterior pituitary gland. The patient has increased urinary output and increased thirst. Which hormone has most likely been altered?

A. Insulin

B. Vasopressin

C. Growth hormone

D. Follicle stimulating hormone

- 40. The nurse is caring for a patient with stage III pressure ulcer to the coccyx. Three days after initiating the plan of care, the nurse observes that the ulcer has hard black crust covering the center of the ulcer. The nurse should understand that this indicates**
- a. Healing
 - b. Need for debridement**
 - c. Inadequate nutrition
 - d. Infection
- 41. A nurse is caring for a hospitalized diabetic patient with advanced peripheral recovery. Which of the following nursing action is MOST important?**
- a. Moisturizing the skin with lotion each day
 - b. Ensuring that foods are not too hot
 - c. Facing the patient when speaking
 - d. Assessing the heels for breakdown**
- 42. A parent is concerned their 8-year-old child has 23kg (51lb) over the past 2 weeks and has been urination up to 30 times per day. The child also seems to be eating and drinking constantly. Which test would be MOST helpful in evaluating the child's condition?**
- a. Chest X-ray
 - b. Complete blood count
 - c. Body fat analysis
 - d. Blood glucose level**
- 43. A patient with dementia is being treated for dehydration. The patient is confused and has been immobile for the past month. Currently, the patient is incontinent and unable to feed self. The nursing care plan should include**
- a. Coughing and deep breathing every 30 minutes
 - b. Positioning and turning every 2 hours**
 - c. Range of motion exercise to all extremities every hour
 - d. Ambulates at least 20 steps every shift
- 44. A 28 year old male is recovering from a moderate concussion following a motor vehicle accident 2 weeks ago, when he suddenly develops an increased thirst, craving Coldwater. The patient urinates**

very large amount of dilute, water like urine with a specific gravity of 1.001 to 1.005 the patient is MOST likely developing

a. Diabetic mellitus

b. Diabetic insipidus

c. Hypothyroidism

d. Thyroid storm

45. A three year-old has returned to the clinic 4 days after being diagnosed with gastroenteritis and dehydration. A parent reports that the vomiting has stopped, and the child is tolerating liquids, rice, applesauce, and bananas. The diarrhea persists, but seems to be decreasing in volume. When evaluating for signs of dehydration, the nurse will assess the patient's skin turgor by:

a- Grasping the skin over the abdomen with two fingers and raising the skin with two fingers

b- Grasping the skin over the forehead with two fingers and raising the skin with two fingers

c- Holding the patient's mouth open and assessing the tongue for deep creases or furrows

d- Drawing two tubes of blood and running a blood urea nitrogen (BUN) and creatinine (Cr)

The digestive system & renal system

1. The nurse is planning care for several children who were admitted during the shift. Daily weights should be the plan of care for the child who is receiving:

a. Total parenteral nutrition(TPN)

- b. Supplement oxygen
- c. Intravenous anti-ineffective
- d. Chest physiotherapy

2. While caring for a patient with an ileostomy, the nurse would expect the ostomy to be located In Which Quadrant of the abdomen?

a. Right lower

- b. Left lower
- c. Left upper
- d. Right upper

3. A nurse is caring for a patient receiving total parenteral nutrition (TPN). The patient reports the sudden onset of feeling short of breath and anxious. The nurse hears crackles in bilateral lower lobes of the lungs and the patient's O2 saturation is 90 % on room air. The nurse must IMMEDIATELY:

a. Turn off the TPN

- b. Notify the physician
- c. Asses the patient's capillary blood glucose level
- d. Attempt to suction the patient's airway

4. A nurse has just started total parenteral nutrition TPN1 as prescribed for a patient with severe dysphagia and very low pre albumin levels , What should the nurse assess in one to two hours ?

a. Blood glucose level

b. Weight

c. Liver

d. O₂ saturation 1 Total parental Nutrition

5. While the nurse is administering a large volume enema, the patient complains of cramping. The nurse should:

a. Increase the flow rate

b. Lower the fluid container

c. Elevate the head of the bed

d. Gently massage the abdomen

6. A 52-years-old is admitted to the nursing unit from the physician's office with a diagnosis of acute cholecystitis. Physician orders on admission include: monitor vital sign every 4 hours; IV of ringer's lactate 125 ml per hour; 1500 calorie, low-fat liquid diet, morphine sulfate 2 mg IV every 2 hours as needed for pain, notify physician for sudden increase in frequency or intensity of pain, promethazine 12.5 mg IV every 4 hours as needed for nausea or vomiting. Which of the following should the nurse plan to do FIRST?

a. Remove any high-foods from the patient's room

b. Notify the dietitian of the diet order

c. Obtain venous access and start Ringer's lactate infusion

d. Obtain an emesis basin and clean linens for the be side

7. A patient has a 6-year history of inflammatory bowel disease that is resistant to medical therapy. The patient can BEST decreased the like hood of the disease progressing to

a. Consuming only elemental foods

b. Stopping smoking

c. Using effective birth control

d. Avoiding over heating

8. A patient is one day post-operative repair of a large umbilical hernia. The patient complains of abdominal pain and described feeling the sutures give way. Upon assessment of the abdomen the nurse observes an evisceration. The nurse's IMMEDIATE response should be to:

- a. Medicate the patient for pain
- b. Instruct the patient to cough hard
- c. Have the patient perform the valsalvas maneuver
- d. Cover the abdomen with a sterile soaked dressing**

9. A patient with severe diverticulitis had surgery for placement of colostomy. The patient is upset, crying and will not look at the colostomy. Which of the following would be the HIGHEST priority nursing diagnosis at this time?

- a. Knowledge deficit, colostomy care
- b. Distorted body image**
- c. Self-care deficit, toileting
- d. Alteration in comfort

10. When administering an enema to adult patient, how far should the nurse insert the tubing into the rectum?

- a. 2.2 to 4.4cm (1 to 2 inches)
- b. 4.4 to 6.6cm (1 to 3 inches)
- c. 6.6 to 8.8cm (3 to 4 inches)**
- d. 8.8 to 11cm (4 to 5 inches)

11. During the postoperative period, a nurse is assigned to care for a morbidly obese patient with an abdominal incision. The nurse knows that this patient's weight increases the risk of:

- a. Left-sided heart failure
- b. Pressure sores of the coccyx

c. Constipation and ileus

d. Wound dehiscence

12. A nurse is assigned to care for a patient with an ileostomy. The nurse would expect the ostomy discharge to be:

a. Fluid mushy

b. Mushy

c. Liquid

d. Solid

13. The nurse is caring for a patient who had a total proctocolectomy 24 hours ago due to a malignant neoplasm in the rectum. The patient continues to receive intravenous fluids and has started a clear liquid diet. The nurse understands that the patient is at INCREASED risk for which of the following postoperative complications?

a. Dissemination intravascular coagulopathy (DIC)

b. Atelectasis

c. Syndrome of inappropriate anti-diuretics hormone(SIADH)

d. Hypokalemia

14. A Patient is recovering following surgery for placement of a colostomy. The nurse goes to the patient's room to instruct the patient how to care for the colostomy. The patient's roommate has visitors and the patient does not want to participate at this time. What should the nurse do?

a. Document the patient's refusal and add non-compliance to the care plan

b. Tell the patient that this is vital information and may delay discharged

c. Plan a time convenient to both the patient and the nurse

d. Pull the curtain around the bed and speak, ensuring privacy

15. A physician has ordered gavage feeding every 4 hours for a 12-week-old infant with failure to thrive. In order to know how far to insert the feeding tube. The nurse should measure the distance from:

- a. The infant's mouth to the xiphoid process of the sternum
- b. The tip of the infant's nose to the ear and then to the umbilicus
- c. The infant's mouth to the ear and then to the umbilicus
- d. The tip of the infant's nose to the ear and then to the xiphoid process of the sternum

16. A patient has hepatitis B (HBV) and is now a chronic carrier. In planning care, the nurse would explain an HBV carrier would MOST likely be at risk for developing a super infection with which other type of hepatitis?

- a. A
- b. C
- c. E
- d. D

17. A preoperative patient has a large volume cleansing enema ordered. In order to facilitate the flow of the solution into the rectum and colon, the nurse should position the patient in the:

- a. Supine position with legs flexed to chest
- b. Right lateral position with left sharply flexed
- c. Supine position with legs spread
- d. Left lateral position with right leg sharply flexed

18. Following an open-cholecystectomy, the nurse would instruct the patient to expect to resume normal activities in:

- a. 1 to 2 weeks
- b. 2 to 3 weeks
- c. 4 to 6 weeks

d. 6 to 8 weeks

19. A nurse has been visiting a bed-bound patient with decreased bowel mobility in the home for one month. The family tells the nurse that the patient is becoming incontinent of feces. The nurse evaluates the plan of care and notes which of the following intervention would MOST likely beneficial?

- a. An enema two times a week
- b. Increased fiber in the diet
- c. Aroutinebisacodyl (Dulcolax) suppository

d. An enema three times a week

20. During surgery the pt has the following intake and output: IV fluid 650 cc ,IV antibiotic 50 cc , 1 unit of packed red blood cells 350 cc , nasogastric output 120 cc , estimated blood loss 80 cc , and urine in the folyes catheter 240 cc . what is the patient's total intake

- a. 650 cc
- b. 700 cc
- c. 900 cc

d. 1050 cc

21. The nurse is teaching a group of patient about hepatitis A(HAV). The nurse should state that HAV is MAINLY transmitted Via:

- a. Blood contact
- b. Food**
- c. Sexual activity
- d. Saliva

22. Prior to administering an enema, the nurse will assist the patient to assume what position

- a. Prone with pillow under knees
- b. Left-side with right knee flexed**

- c. Right-side with left knee flexed
- d. On back with head of bed flat

23. 60 years age a patient weighed 73 kilograms (161 pounds). During the current clinic visit the nurse note the patient has an unintended weight loss. This weight loss over 6 months would be considered clinically significant as soon as it reaches the point of being more than a:

- a. 5% loss
- b. 8% loss
- c. 10% loss**
- d. 20% loss

24. A nurse is assessing a 4-month-old formula-fed infant. The parent reports the infant has been irritable, crying excessively, not sleeping well, and vomiting. Gastro-esophageal reflux is expected. What nursing intervention should the nurse expect to teach the parent?

- a. Place the infant in an infant seat after eating**
- b. Give large frequent feedings
- c. Position the child in a swing
- d. Thin formula with water

25. A parent brings their teenage child the pediatrician's office. The parent reports that the patient frequently complains of abdominal bloating and stomach pain after eating and also has a chronic sore throat. The patient's labs show hypokalemia . Which of the following diagnosis should the nurse anticipate?

- a. Anorexia nervosa
- b. Bulimia**
- c. Morbid obesity
- d. Impulsive behavior

26. A urinalysis is best evaluated for accurate result if specimen is analyzed within:

- a. 1 hour of collection or refrigerated until analyzed
- b. 1 hour of collection or left at room temperature
- c. 2 hours of collection**
- d. 4 hours of collection

27. A patient admitted to the hospital with acute cholecystitis , is scheduled for surgery in the morning and is NPO. At 8 am the patient develops a fever of 102.4 F (39.1 C).medication orders include acetaminophen 650 mg orally every four hours as needed. The nurse should:

- a. Give the medication as ordered by the physician
- b. Administer the ordered dose rectally
- c. Put moist cool cloths on the patient's forehead and axillae
- d. Notify the physician and request other orders**

28. During surgery, the patient has the following intake and output: intravenous fluid 650 cc, intravenous antibiotic 50 cc, 1 unit of packed red blood cells (PRBC) 350 cc , nasogastric output 120 cc, estimated blood loss 80 cc, and urine in the Foley catheter 240 cc. What is the patient's total output?

- a. 120 cc
- b. 200 cc
- c. 240 cc
- d. 440 cc**

29. The traction and urinary catheter have been discontinued for a patient who was immobilized in traction for 6 weeks . The pt is now having a problem with urinary incontinence . Which of the following interventions would the nurse most likely implement?

- a. Behavioral training

b. Bladder training

- c. Scheduled toileting
- d. Prompted voiding

30. A patient is diagnosed with peptic ulcer. What would be the long term goal for this patient?

a. Patient remains free of signs and symptoms of gastrointestinal bleeding

b. Patient maintains lifestyle alterations to prevent recurrence of ulcer

- c. Patient expresses decreased pain level
- d. Patient performs activities of daily living without difficulty

31. A patient comes to the medical office with complaints of some urinary incontinence. The nurse discovers the incontinence occurs because of an inability to delay voiding long enough to reach a toilet after the patient feels a sensation of bladder fullness. This type of incontinence is:

a. Stress

b. Urge

- c. Overflow
- d. Functional

32. A patient who had abdominal surgery 6 days ago , has been ambulating the halls without much difficulty. However, on day 7 postoperative the patient complains of increased pain at incisional site and is walking hunched over the MOST likely cause of the change is

- a. Over assertion the day before
- b. Pulmonary edema

c. wound infection

- d. deep vein thrombosis

33. A nurse is preparing an assessment of a patient's nutritional status. Which of the following diagnostic test would be the best measure of the patient's recent nutritional status with a half- life of 2-3 days?

- a. Prealbumin**
- b. Hemoglobin
- c. Albumin
- d. 24- urine Creatinine

34. When is the best time to collect urine specimen for routine urinalysis and culture and sensitivity?

- a. Early morning**
- b. Later afternoon
- c. Midnight
- d. Before breakfast

35. Following abdominal surgery, a child has a nasogastric tube connected to suction. Several hours after surgery, the child tells the nurse that he is nauseated and then vomits approximately 200 ml of fluid. Which of these actions should the nurse take first?

- a. Notify the physician
- b. Check if the nasogastric tube can be irrigated
- c. Discontinue the section attached to the nasogastric tube**
- d. Auscultate for bowel sounds

36. Patient is planning to have an elective surgical procedure to repair an umbilical hernia. The patient is 68 years old. Weighs 136 kg (300 lbs.) Has diabetes mellitus. Which of the following approaches would be the MOST beneficial in order to reduce the patients surgical risk?

- A. Monitor blood glucose levels monthly

- B. Avoid fluid overload by restricting fluids
- C. Discourage any changes in routine before surgery

D. Encourage weight reduction

37. A 52 year- old woman is scheduled to undergo an abdomino-perineal resection in three days for removal of a cancer of the rectum. The nurse reviews the care plan with the patient. The patient will receive prophylactic antibiotics and will be given a mechanical bowel preparation the day before. Which additional preparation should the patient undertake at this time?

- a. Wear pressure stockings
- b. Perform leg strengthening exercises
- c. Maintain high- protein, low- residue diet**
- d. Take daily ferrous iron tablets

38. A patient has a central line catheter and is receiving a three-in-one total parenteral nutrition that contains glucose, proteins and lipids. The pump is set to deliver the infusion over a 12-hour period. After how many hours should the intravenous administration set be changed?

- a. 12
- b. 24**
- c. 48
- d. 72

39. Which of the following is Pancreatitis Patients frequently feel constant pain in the

- a. lower right
- b. upper Left**

c. lower Left

d. upper right

40. A nurse is preparing to insert a nasogastric tube into a client. The nurse places the client in which position for insertion:

a.Right side

b.Low Fowler's

c.High Fowler's

d.Supine with the head flat

41. A patient has an acute inflammation of the gallbladder. The physician orders the nurse to schedule the patient for surgery . Which of the following surgical procedures will the physician MOST likely perform?

A. Pancreatectomy .

B. Cholecystectomy

C. Hepatectomy .

D. Cricothoracotomy


42. Which of the following is the incision site for an open cholecystectomy operation:

a- 1

b- 2


c- 3

D- 4




43. Which of the following is the incision site for an open gastrectomy operation:

- a- 1
- b- 2
- c- 3
- d- 4**



44. Which of the following is the incision site for surgical appendectomy operation:

- a- A
- b- B
- c- C**
- d- D



45. A nurse is administering a cleansing enema to a client with a fecal impaction. Before administering the enema, the nurse places the client in which position:

- a- Left Sims' position**
- b- Right Sims' position
- c- On the left side of the body, with the head of the bed elevated 45 degrees
- d- On the right side of the body, with the head of the bed elevated 45 degrees

46. Which of the following instructions is appropriate for the nurse to give to a female client who complains of abdominal upset after cholecystectomy operation:

- a-Increase fluid intake
- b-Avoid fatty meal**
- c-Increase protein intake
- d-Daily exercise

47. Which of the following is the best way for long term feeding for a patient experiencing severe dysphagia:

- a-NGT
- b-Naso-duodenal tube
- c-Gastrostomy**
- d-Parenteral

Kozier & Erb's Fundamentals of
Nursing (9th Edition) chapter 47 / page
1288

48. A patient suffers from diarrhea and muscle weakness, the nurse would assess this patient for which of the following abnormal laboratory blood tests:

- a-Blood calcium
- b-Blood potassium**
- c-Blood magnesium
- d-Blood sodium

49. A nurse has administered approximately half of an enema solution to a preoperative client when the client complains of pain and cramping. Which nursing action is the most appropriate:

- a-Raise the enema bag so that the solution can be instilled quickly.
- b-lamp the tubing for 30 seconds and restart the flow at a slower rate.**
- c-Reassure the client and continue the flow.
- d-Discontinue the enema and notify the physician

50. When caring for a client who is receiving total parenteral nutrition (TPN), which of the following complications would be most important for the nurse to assess:

- a-Chest pain
- b-Hemorrhage and air embolus
- c-Pneumonia and hyperglycemia
- d-Electrolyte imbalance and sepsis**

51. Postoperatively, patient is expected to void after:

- a-6-8 hours**
- b-2-4 hours
- c-12 – 24 hours
- d-10-12 hours

52. A 40 years old female complains of abdominal pain. It is worse after eating, especially if she has a meal that is spicy or high in fat. She has tried over-the-counter antacids, but they have not helped the pain. After examining her abdomen, you strongly suspect cholecystitis. Which sign on examination increases your suspicion for this diagnosis:

a-Psoas sign

b-Rovsing's sign

c-Murphy's sign

d-Grey turner's sign

53. When planning the care of a client who has undergone an abdominal hysterectomy, which nursing measure is most helpful for preventing postoperative complications and facilitating an early discharge?

a-Reestablishing oral fluids and nutrition

b-Promoting ambulation and movement

c-Maintaining accurate intake and output

d-Exploring feelings about altered image

54. Which of the following is considered one of the common complications following abdominal total hysterectomy:

a-DVT

b-Paralytic ileus

c-Constipation

d-Perforation

55. A client who has a history of Crohn's disease is admitted to the hospital with fever, diarrhea, cramping, abdominal pain, and weight loss. The nurse should monitor the client for:

a-Hyperalbuminemia

b-Thrombocytopenia

c-Hypokalemia

d- Hypercalcemia

MARYAM ALMUHANNA

29. To decrease the incidence of aspiration of gastric contents in a child hospitalization with severe burns, the nurse should position the head:

- a. Flat except during meals
- b. Elevates 30-45 degrees during meals**
- c. Elevated 15-30 degrees for 12-hours after meals
- d. Elevated 45 degrees at all times

34. Following lumbar surgery a patient has a 4 millimeter (mm) surgical incision. The incision is clean and the edges are well appropriate. This type of tissue healing is classified as which of the following?

- a. Primary intention**
- b. Secondary intention
- c. Tertiary intention
- d. Superficial epidermal

36. A surgeon instructs a nurse to serve as a witness to an elderly patient's informed consent for surgery. During the explanations to the patient, it becomes clear that the patient is confused and does not understand the procedure, but reluctantly sign the consent form. The nurse should:

- a. Sign the form as a witness, making a notation that the patient did not appear to understand**
- b. Not sign the form as a witness and notify the nurse supervisor
- c. Not sign the form and answer the patient's questions after the surgeon leaves the room
- d. Sign the form and tell surgeon that the patient doesn't understand the procedure

38. A patient scheduled for a major surgery in one hour is very nervous and upset. Which of the following order medications would the nurse administer to relax this patient?

- a. Meperidine Hydrochloride(Demerol)
- b. Scopolamine (Transderm-Scop)
- c. Pentobarbital sodium(Nembutal sodium)**
- d. Trazodone hydrochloride(Trazadone)

45. While caring for a patient in the post-anesthesia care unit (PACU) Who has developed Hypovolemic shock, a nurse should position the patient:

- a. Flat with legs elevated**
- b. In Trendelenburg position
- c. With the head of the bed elevated 45 degrees
- d. Completely flat

150. A patient with long-standing diabetes mellitus (type I) is scheduled for surgical amputation of 4 gangrenous toes on the right foot. Which surgical intervention would this be classified as?

- a. Palliative
- b. Curative**
- c. Reconstructive
- d. Diagnostic

155. A patient is in the preoperative area to lumbar surgery. The patient reports anxiety about being intubated and expresses concern about waking up during the surgery. The nurse MUST discuss the patient's concern with the

- a. Anesthesia provider**
- b. Surgeon

- c. Scrub nurse
- d. Charge nurse

157. A nurse assesses a 3-month-old infant. The patient expresses anxiety and feeling over whelmed. The nurse offer information on available parenting support. This level of child abuse prevention is classified as which of the following?

- a. Intervention
- b. Primary**
- c. Secondary
- d. Tertiary

165. A nurse assessing a 16-month-old child observes bruises scattered over the body that are at different stage of healing. The child also has poor and diaper rash. What the goal of treatment for this child ?

- a. Ensure the physical and emotional safety of the child**
- b. Remove the child from the parents
- c. Admonish the parents of the child
- d. Ensure that the child stays with the biological parents

274. A co-worker informs that the nurse about experiencing increased level of stress associated with daily responsibilities to help cope with professional stress, the nurse should encourage the co-worker to ;

- a. Make a list of unfinished tasks**
- b. complete complex mental task before physical tasks
- c. Acknowledge daily accomplishments

d. Spend time with colleague away from work

334. The responsibility for teaching patients how to take medications safely when they are discharged from the hospital belongs to the:

a. Nurse

b. Physician

c. Dietitian

d. Therapis

354. When administering an intramuscular injection to an infant, which of the following sites appropriate for the nurse to use?

a. Rectus femoris

b. Deltoid

c. Dorsogluteal

d. Ventrogluteal

402. The nurse administered a prescribed intramuscular medication to a patient during a home health visit. How should the nurse dispose of the used needle and syringe?

a. Recap the needle, then place the needle and syringe into a waterproof container until safe disposal can be made

b. Bend the needle back towards the barrel of the syringe before putting the needle and syringe in a metal trash container

c. Wrap the needle and syringe in disposable paper before putting the needle and syringe into the dirty section of the nurse's equipment bag

d. Put the needle and syringe directly into a puncture-resistant plastic container that has a lid

510. Patient with diagnosis of Dilated cardiomyopathy. [700] The Medication order: Glucophage 850mg po. qd Imdur 60mg. po .qd Lasix 80mg. po. qd Zocor 40mg. po. Qd Past medical history: Heart failure, DM type 2, Peripheral vascular disease (pvd) & primary

hyperlipidemia . Based on the patient's history which of the following medication orders should the nurse verify?

a. Glucophage

b. Zocor

c. Lasix

d. Imdur

540. What of the following has propriety to check before start giving patient blood transfusion

a. Blood group

b. Name

c. Expiry date

608. A 16 year-old boy is in the Post-Operative Care Unit two hours after a tonsillectomy. He is alert and oriented but complains of severe throat pain and difficulty swallowing. He rates the pain at a level 7, on a scale of 1-10. The urine output from the folly catheter is 45 ml over the past two hours. Blood pressure 130/74 mmHg Heart rate 64/min Respiratory rate 18/min Oxygen saturation 98 % on room air, Which clinical finding is most important to report to the doctor?

a. Oxygen saturation

b. Difficulty swallowing

c. Urinary output

d. Pain level

Maryam 2

7) A 58 year-old man has been diagnosed with stage Iv prostate cancer with metastasis to the bones. The patient refuses a medical management plan that includes chemotherapy and radiation. Although the nurse believes that every possible intervention should be done to preserve life, the nurse supports the patient's decision.

Which ethical principle underlies the nurse's response?

- A. fidelity
- B. beneficence
- C. non-maleficence
- D. Autonomy

13) A 30 year-old primi-para is admitted to the postpartum unit with a healthy full-time infant boy. She had undergone an emergency concern section after an umbilical prolapse. All of the patient beds are occupied and the nurse considers some steps of the patient's nursing process that could be delegated to the nursing.

Which step would be most likely?

- A. planning
- B. Evaluation
- C. Assessment

D. Implementation

41) A two month-old infant is brought to the clinic by her parents who report that the baby is having breast feeding difficulties. The infant feeds every one-two hours for twenty minutes and seems to be in pain afterwards. She regurgitates milk after each feed. At birth, she weighed 3.5 Kilograms and her current weight is 4.2 Kilograms.

Which initial home intervention would be most appropriate?

- A. offer milk-based formula before breastfeeding
- B. feed with rice cereal mixed with breast milk
- C. Schedule feeds regular every three hours.

D. Position the baby upright after feeding

9) In the summer months, a five year-old girl present with a sore throat and a dairy cough that has slowly become worse over the past three weeks. Her body temperature is 38.0 ° C. on auscultation, there is wheezing and shortness of breath. She lives in an overcrowded house with thee where she attends school.

Which is the greatest risk factor?

- A. Residing in a low-income neighborhood
- B. Exposure to a pathogen in summer season
- C. Attending school
- D. Living in crowded conditions**

Maryam 3

A 65 year-old woman presented to her care provider with complaints of bright red blood in the stool, a loss of appetite, a feeling of fullness and fatigue. She had lost 5 kilograms in the past three weeks without dieting. A faecal occult blood test is positive and the patient is scheduled for additional screening test.

Which screening test is most likely for this patient?

- a- Barium enema
- b- Colonoscopy**
- c- Endoscopy
- d- Computed tomography scan

A 65 year-old woman presented to her care provider with complaints of bright red blood in the stool, a loss of appetite, a feeling of fullness and fatigue. She had lost 5 kilograms in the past three weeks without dieting. A faecal occult blood test is positive and the patient is scheduled for additional screening test.

Which of the following is most likely diagnosis for this patient?

- a- Gastric cancer
- b-duodenal cancer
- c- colon cancer**

Maryam Almuhanha

d- ovarian cancer

Rn. Maryam

Rn. Maryam