

A Partial Solution Manual for: *The Elements of  
Statistical Learning* by Jerome Friedman, Trevor  
Hastie, and Robert Tibshirani

Wenhao Wu

wnhwu@ucdavis.edu

Dept. ECE, UC Davis

February 14, 2016

# Contents

Preface	2
Acknowledgment	3
2 Overview of Supervised Learning	4
3 Linear Methods for Regression	5
4 Linear Methods for Classification	6
5 Basis Expansions and Regularization	7
6 Kernel Smoothing Methods	8
7 Model Assessment and Selection	9
8 Model Inference and Averaging	10
9 Additive Models, Trees, and Related Methods	11
10 Boosting and Additive Trees	12
11 Neural Networks	13
12 Support Vector Machines and Flexible Discriminants	15
13 Prototype Methods and Nearest-Neighbors	23
14 Unsupervised Learning	27
15 Random Forests	37
16 Ensemble Learning	38
17 Undirected Graphical Models	39
18 High-Dimensional Problems	40

# Preface

This work is expected to be used as a supplementary material for Weatherwax and Epstein's solution manual [?], which I found to be very helpful when self-studying this popular textbook. The numbering of chapters and problems are based on the 2nd edition (10th printing with corrections, Jan 2013) available online [?].

The author was not able to solve all the excercises. Even for the solutions included we expect many mistakes and shortcomings. It would be of great help if people could suggest possible solutions or help us find and correct the errors so this solution manual can be continuously improved to benefit more interested readers. We are also open to all comments and criticisms. Our contact information can be found at the website holding this draft [?].

# Acknowledgment

## Chapter 2

### Overview of Supervised Learning

## Chapter 3

### Linear Methods for Regression

## Chapter 4

### Linear Methods for Classification

## Chapter 5

### Basis Expansions and Regularization



## Chapter 6

### Kernel Smoothing Methods

## Chapter 7

### Model Assessment and Selection

## Chapter 8

### Model Inference and Averaging

## Chapter 9

### Additive Models, Trees, and Related Methods

## Chapter 10

### Boosting and Additive Trees

# Chapter 11

## Neural Networks

### Ex. 11.1

In (11.5), set  $K = 1$ ,  $g_1(T) = T$ , we have

$$f_1(X) = \beta_{01} + \beta_1^T Z = \beta_{01} + \sum_{m=1}^M \beta_{m1} \sigma(\alpha_{0m} + \alpha_m^T X) \quad (11.1)$$

The correspondence between (11.1) and (11.5) becomes clearer, as enumerated in Table 11.1

Table 11.1: Correspondence between the project pursuit regression and the neural network

(11.1)	(11.5)
$\omega_m$	$\alpha_m$
$g_m(\cdot)$	$\beta_{01}, \beta_{m1} \sigma(\alpha_{0m} + \alpha_m^T X)$

### Ex. 11.2

$$\frac{\partial f}{\partial X} = \sum_{m=1}^M \beta_m [\sigma(\cdot)(\sigma(\cdot) - 1)] \alpha_m \quad (11.2)$$

$$\frac{\partial^2 f}{\partial X \partial X^T} = \sum_{m=1}^M \beta_m [(2\sigma(\cdot) - 1)(\sigma(\cdot) - 1)\sigma(\cdot)] \alpha_m \alpha_m^T \quad (11.3)$$

Since  $\sigma(\alpha_{0m} + \alpha_m^T X) \approx 1/2$  when  $\alpha_{0m} \approx 0$  and  $\alpha_m \approx 0$ , therefore  $\frac{\partial^2 f}{\partial X \partial X^T} \approx 0$ , i.e. the resulting model is nearly linear.

### Ex. 11.3

$$R(\theta) = - \sum_{i=1}^N R_i(\theta) = - \sum_{i=1}^N \sum_{j=1}^K y_{ij} \log g_j(T) \quad (11.4)$$

Note that different from regression, each softmax function  $g_j(T)$ ,  $j = 1, \dots, K$  is a function

of all  $T_1, \dots, T_K$ .

$$\frac{\partial R_i}{\partial \beta_{km}} = - \sum_{j=1}^K \frac{y_{ij}}{g_j} \frac{\partial g_j}{\partial T_k} z_{mi} = \delta_{ki} z_{mi} \quad (11.5a)$$

$$\begin{aligned} \frac{\partial R_i}{\partial \alpha_{ml}} &= - \sum_{j=1}^K \frac{y_{ij}}{g_j} \sum_{k=1}^K \frac{\partial g_j}{\partial T_k} \beta_{km} \sigma'(\alpha_m^T x_i) x_{il} \\ &= \left[ \sigma'(\alpha_m^T x_i) \sum_{k=1}^K \beta_{km} \delta_{ki} \right] x_{il} = s_{mi} x_{il} \end{aligned} \quad (11.5b)$$

It is noted that

$$\frac{1}{g_j} \frac{\partial g_j}{\partial T_k} = \begin{cases} 1 - g_j & j = k \\ -g_k / \exp(T_j) & j \neq k \end{cases} \quad (11.6)$$

As a result, although  $g_j(T)$  depends on all  $T_1, \dots, T_K$ ,  $(\partial g_j / \partial T_k) / g_j$  can still be locally evaluated and propagated downward over the link  $(T_k, g_j)$ . Consequently, the forward and backward propagation equations are pretty much the same as those for the square error loss function. In the forward pass for record  $x_i$ ,  $i = 1, \dots, N$ , the weights  $\beta_{km}$  and  $\alpha_{ml}$  are fixed and the predicted  $\hat{g}_j(T_i)$  are evaluated. In the backward pass,  $(y_{ij}/g_j)(\partial g_j / \partial T_k)$  are evaluated and propagated to  $T_k$ , where  $\delta_{ki}$  is computed, and then back-propagated to give  $s_{mi}$  at  $Z_m$ . Then the gradients are evaluated as in Eq. (11.5). The gradient descent update is exactly the same as (11.13).

#### Ex. 11.4

If the network has no hidden layer, we have

$$g_j(x) = \frac{\exp(T_j)}{\sum_{k=1}^K \exp(T_k)} = \frac{\exp(\beta_j^T x)}{\sum_{k=1}^K \exp(\beta_k^T x)}, \quad (11.7)$$

exactly the same as the multinomial logistic model.

#### Ex. 11.5 (Program)

#### Ex. 11.6 (Program)

#### Ex. 11.7 (Program)

## Chapter 12

### Support Vector Machines and Flexible Discriminants

#### Ex. 12.1

Firstly, we prove that for (12.8), the optimal solution must satisfy  $\hat{\xi}_i = [1 - y_i(x_i^T \hat{\beta} + \hat{\beta}_0)]_+$ . To see this, from the constraints in (12.8), we have  $\hat{\xi}_i \geq [1 - y_i(x_i^T \hat{\beta} + \hat{\beta}_0)]_+$ . Assume for contradiction that  $\exists i$  such that  $\hat{\xi}_i > [1 - y_i(x_i^T \hat{\beta} + \hat{\beta}_0)]_+$ , then setting  $\hat{\xi}_i \leftarrow [1 - y_i(x_i^T \hat{\beta} + \hat{\beta}_0)]_+$  results in smaller objective in (12.8), which is in contradiction to the fact that  $\hat{\xi}_i$  is from an optimal solution.

On the other hand,  $\xi_i = [1 - y_i(x_i^T \beta + \beta_0)]_+ \Rightarrow \xi_i \geq 0, y_i(x_i^T \beta + \beta_0) \geq 1 - \xi_i$ . Therefore, the solution to (12.8) is the same as

$$\min_{\beta, \beta_0} \frac{1}{2} \|\beta\|^2 + C \sum_{i=1}^N \xi_i \quad (12.1)$$

$$\text{s.t. } \xi_i = [1 - y_i(x_i^T \beta + \beta_0)]_+, \forall i \quad (12.2)$$

which is exactly the same as (12.25).

#### Ex. 12.2

Define kernel  $K(a, b) = \sum_{j=1}^p a_j b_j$ , i.e.  $\psi_j(x) = x_j, \gamma_j = 1$  for  $j = 1, \dots, p$ . Consequently,  $g(x) = \sum_{j=1}^p \beta_j x_j \Leftrightarrow g(x) \in \mathcal{H}_K$ . Consequently,

$$(12.25) \Leftrightarrow \min_{g, \beta_0} \sum_{i=1}^N [1 - y_i(g(x_i) + \beta_0)]_+ + \frac{\lambda}{2} \|g\|_{\mathcal{H}_K}^2 \quad (12.3)$$

Denote  $L(y_i, g(x_i); \beta_0) = [1 - y_i(g(x_i) + \beta_0)]_+ = L_i(\beta_0)$ , then

$$(12.25) \Leftrightarrow \min_{\beta_0} \left\{ \min_g \sum_{i=1}^N L_i(\beta_0) + \frac{\lambda}{2} \|g\|_{\mathcal{H}_K}^2 \right\}. \quad (12.4)$$

where the inner min must have a solution in the form of  $g(x) = \sum_{i=1}^N \alpha_i K(x, x_i)$  as per



(5.50)(5.51), and we have  $\|g\|_{\mathcal{H}_K}^2 = \alpha^T K \alpha$ . Therefore

$$(12.25) \Leftrightarrow \min_{\beta_0} \left\{ \min_{\alpha} \sum_{i=1}^N [1 - y_i (\sum_{j=1}^N \alpha_j K(x_j, x_i) + \beta_0)]_+ + \frac{\lambda}{2} \alpha^T K \alpha \right\} \quad (12.5)$$

$$\Leftrightarrow \min_{\beta_0, \alpha} \sum_{i=1}^N [1 - y_i (\sum_{j=1}^N \alpha_j K(x_j, x_i) + \beta_0)]_+ + \frac{\lambda}{2} \alpha^T K \alpha \quad (12.6)$$

### Ex. 12.3

Similar to Ex. (12.2). Denote  $g(x) = \sum_{m=1}^M \beta_m h_m(x)$ . Without penalizing the constant term, we have

$$H(\beta, \beta_0) = \sum_{i=1}^N V(y_i - \beta_0 - g(x_i)) + \frac{\lambda}{2} \sum_{m=1}^M \beta_m \quad (12.7)$$

Again we break the minimization problem into 2 steps:

$$\min_{\beta_0, \beta} H(\beta, \beta_0) = \min_{\beta_0} \left\{ \min_{\beta | \beta_0} H(\beta, \beta_0) \right\} \quad (12.8)$$

Consider saquare error loss  $V(r) = r^2$ , the inner min problem is in the form of

$$\min_{\beta} \sum_{i=1}^N (y_i - \beta_0 - H\beta)^2 + \frac{\lambda}{2} \beta^T \beta \quad (12.9)$$

$$\Leftrightarrow \min_{\alpha} \|\mathbf{y} - \beta_0 \mathbf{1} - \mathbf{K}\alpha\|_F^2 + \frac{\lambda}{2} \alpha^T \mathbf{K}\alpha \quad (12.10)$$

whose solution is  $\hat{\alpha} = (\mathbf{K} + \lambda \mathbf{I}/2)^{-1} \mathbf{y}_{\beta_0}$ ,  $\mathbf{y}_{\beta_0} = \mathbf{y} - \beta_0 \mathbf{1}$ . Consequently, the outer min problem w.r.t  $\beta_0$  is in the form of

$$\min_{\beta_0} \mathbf{y}_{\beta_0}^T [\mathbf{I} - (\mathbf{K} + \lambda \mathbf{I}/2)^{-1} \mathbf{K}] \mathbf{y}_{\beta_0} \quad (12.11)$$

which is a quadratic problem.

### Ex. 12.4

(a)

$$\text{Left} = (x - \bar{x}_k)^T U U^T (x - \bar{x}_k) - (x - \bar{x}_{k'})^T U U^T (x - \bar{x}_{k'}) \quad (12.12)$$

where  $U = W^{-1/2} V^*$ , the  $L$  columns of  $V^*$  are the eigen vectors of  $B^* = (W^{-1/2})^T B W^{-1/2}$ ,

where  $B$  is the between-class covariance.

$$\text{Right} = (x - \bar{x}_k)^T W^{-1} (x - \bar{x}_k) - (x - \bar{x}_{k'})^T W^{-1} (x - \bar{x}_{k'}) \quad (12.13)$$

Consequently,

$$\begin{aligned} & \text{Left} - \text{Right} \\ &= 2(\bar{x}_k - \bar{x}_{k'})^T (W^{-1} - UU^T)x + (\bar{x}_k - \bar{x}_{k'})^T [W^{-1} - UU^T](\bar{x}_k + \bar{x}_{k'}) \end{aligned} \quad (12.14)$$

$$\begin{aligned} &= 2(\bar{x}_k - \bar{x}_{k'})^T W^{-1/2} (I - V^*(V^*)^T) (W^{-1/2})^T x \\ &+ (\bar{x}_k - \bar{x}_{k'})^T W^{-1/2} (I - V^*(V^*)^T) (W^{-1/2})^T (\bar{x}_k + \bar{x}_{k'}) \end{aligned} \quad (12.15)$$

Since  $(\bar{x}_k - \bar{x}_{k'})^T \in R(M)$  (row space),  $(\bar{x}_k - \bar{x}_{k'})^T W^{-1/2} \in R(M^*)$ , therefore  $(W^{-1/2})^T (\bar{x}_k - \bar{x}_{k'}) \in C(V^*)$  (column space). Therefore

$$(\bar{x}_k - \bar{x}_{k'})^T W^{-1/2} (I - V^*(V^*)^T) = 0 \quad (12.16)$$

thus Left = Right.

(b) ???

### Ex. 12.5 (Program)

### Ex. 12.6

(a) The  $i$ -th row of  $\mathbf{Y}\theta$  is

$$(\mathbf{Y}\theta)_i = \sum_{j=1}^K 1(Y_{ij} = 1)\theta_j = \theta(g_i) \quad (12.17)$$

(since there are exactly one  $j$  where  $Y_{ij} = 1$  for each  $i$ ). The  $i$ -th row of  $\mathbf{H}\beta$  is

$$(\mathbf{H}\beta)_i = \sum_{j=1}^K \beta_j h_j(x_i) \quad (12.18)$$

therefore

$$\sum_{i=1}^N (\theta(g_i) - \beta^T h(x_i))^2 = \|\mathbf{Y}\theta - \mathbf{H}\beta\|^2 \quad (12.19)$$

(b) According to the definition,  $(\mathbf{D}_\pi)_{kk}$  is the empirical frequency of class  $k$ , and  $\theta_k$  is the score for class  $k$ .  $\theta^T \mathbf{D}_\pi \mathbf{1} = 0$  implies that the average score over the  $N$  records is 0;  $\theta^T \mathbf{D}_\pi \theta = 1$  means the variance of the over the  $N$  records is 1.

(c) Fixing  $\theta$  the optimal  $\beta$  is

$$\hat{\beta} = (\mathbf{H}^T \mathbf{H})^{-1} \mathbf{H}^T \mathbf{Y} \theta \quad (12.20)$$

therefore (12.65) can be rewritten as

$$\min_{\theta} \|(\mathbf{I} - \mathbf{S}) \mathbf{Y} \theta\|^2 \Leftrightarrow \min_{\theta} \theta^T \mathbf{Y}^T \mathbf{Y} \theta - \theta^T \mathbf{Y}^T \mathbf{S} \mathbf{Y} \theta \quad (12.21)$$

where  $\mathbf{S} = \mathbf{H}(\mathbf{H}^T \mathbf{H})^{-1} \mathbf{H}^T$ . Since  $\theta^T \mathbf{Y}^T \mathbf{Y} \theta = N$ , this minimization is equivalent to

$$\max_{\theta} \theta^T \mathbf{Y}^T \mathbf{S} \mathbf{Y} \theta \quad (12.22)$$

(d) Suppose that the SVD of  $\mathbf{H} = \mathbf{U} \mathbf{\Sigma} \mathbf{V}^T$ , then  $\mathbf{S} = \mathbf{U} \mathbf{U}^T$  where  $\mathbf{U}$  is a  $N$ -by- $l$  orthonormal matrix. Therefore  $\mathbf{S}$  has  $L$  eigenvalues of 1 and  $N - L$  eigenvalues of 0. Since constant function is included in  $h_j$ ,  $\mathbf{H} \neq 0$ , therefore  $L > 0$ , so the largest eigenvalue is 1.

(e) (12.53) can be rewritten as

$$ASR = \frac{1}{N} \|\mathbf{Y} \mathbf{\Theta} - \mathbf{H} \mathbf{B}\|_F^2 \quad (12.23)$$

Similar to (c) the solution is the same as

$$\max_{\mathbf{\Theta}} \text{tr}\{\mathbf{\Theta}^T \mathbf{Y}^T \mathbf{S} \mathbf{Y} \mathbf{\Theta}\} \quad (12.24)$$

$$\text{s.t. } \mathbf{\Theta}^T \mathbf{Y}^T \mathbf{Y} \mathbf{\Theta} = \mathbf{I} \quad (12.25)$$

Therefore  $\mathbf{Y} \mathbf{\Theta}$  are the  $K$  largest eigenvectors of  $\mathbf{S}$ .

### Ex. 12.7

The penalized optimal scoring problem is in the form of

$$\min_{\mathbf{\Theta}, \mathbf{B}} \|\mathbf{Y} \mathbf{\Theta} - \mathbf{H} \mathbf{B}\|_F^2 + \lambda \text{tr}(\mathbf{B}^T \mathbf{\Omega} \mathbf{B}) \quad (12.26)$$

Given  $\mathbf{\Theta}$ , the optimal  $\mathbf{B}$  is

$$\hat{\mathbf{B}} = (\mathbf{H}^T \mathbf{H} + \lambda \mathbf{\Omega})^{-1} \mathbf{H}^T \mathbf{Y} \mathbf{\Theta} \quad (12.27)$$

Substitute into Eq. (12.26), we have

$$\min_{\mathbf{\Theta}} \text{tr}(\mathbf{\Theta}^T \mathbf{Y}^T \mathbf{Y} \mathbf{\Theta} - \mathbf{\Theta}^T \mathbf{Y}^T \mathbf{S} \mathbf{Y} \mathbf{\Theta}) \quad (12.28)$$

$$\text{s.t. } \mathbf{\Theta}^T \mathbf{D}_{\pi} \mathbf{\Theta} = \mathbf{I} \quad (12.29)$$

where  $\mathbf{S} = \mathbf{H}(\mathbf{H}^T \mathbf{H} + \lambda \mathbf{\Omega})^{-1} \mathbf{H}^T$ . Therefore  $\mathbf{Y}\mathbf{\Theta}$  are still the eigenvectors of  $\mathbf{S}$ .

### Ex. 12.8

I found the proof to this problem on [?]. I am trying to follow it the best I can and here is my interpretation. Assuming that  $\bar{x} = 0$ . We first perform the generalized SVD:

$$(\mathbf{Y}^T \mathbf{Y})^{-1} \mathbf{Y}^T \mathbf{X} (\mathbf{X}^T \mathbf{X})^{-1} = \mathbf{U} \mathbf{D} \mathbf{V}^T \quad (12.30)$$

$$\text{s.t. } \mathbf{U}^T \mathbf{Y}^T \mathbf{Y} \mathbf{U} = \mathbf{I}, \mathbf{V}^T \mathbf{X}^T \mathbf{X} \mathbf{V} = \mathbf{I} \quad (12.31)$$

Later we will show that both  $\beta_l$  and  $v_l$  are proportional to the columns of  $\mathbf{V}$ . From the GSVD,  $\mathbf{U}$  and  $\mathbf{V}$  satisfy the following 2 equations:

$$\mathbf{U}^T \mathbf{Y}^T \mathbf{X} (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y} \mathbf{U} = \mathbf{D}^2 \quad (12.32a)$$

$$\mathbf{V}^T \mathbf{X}^T \mathbf{Y} (\mathbf{Y}^T \mathbf{Y})^{-1} \mathbf{Y}^T \mathbf{X} \mathbf{V} = \mathbf{D}^2 \quad (12.32b)$$

of which the proof is trivial. First we show that the LDA's discriminant directions  $v_l$  are parallel to the columns of  $\mathbf{V}$ :

**Proposition 12.1.** *For the LDA problem (Fisher)*

$$\max_{\mathbf{A}} \text{tr}(\mathbf{A}^T \mathbf{B} \mathbf{A}), \text{ s.t. } \mathbf{A}^T \mathbf{W} \mathbf{A} = \mathbf{I} \quad (12.33)$$

where

$$\mathbf{B} = \mathbf{X}^T \mathbf{Y} (\mathbf{Y}^T \mathbf{Y})^{-1} \mathbf{Y}^T \mathbf{X} \quad (12.34a)$$

$$\mathbf{W} = \mathbf{T} - \mathbf{B} \quad (12.34b)$$

$$\mathbf{T} = \mathbf{X}^T \mathbf{X} \quad (12.34c)$$

are the between-class, within-class and total variance (up to normalization), the solution is

$$\hat{\mathbf{A}} = \mathbf{V}(\mathbf{I} - \mathbf{D}^2)^{-1/2} \quad (12.35)$$

*Proof.* From Eq. (12.32b) and the second constraint of the GSVD, it is easy to see  $\hat{\mathbf{A}}^T \mathbf{W} \hat{\mathbf{A}} = \mathbf{I}$ . On the other hand,  $\hat{\mathbf{A}}$  diagonalizes  $\mathbf{B}$  by  $\hat{\mathbf{A}}^T \mathbf{B} \hat{\mathbf{A}} = (\mathbf{I} - \mathbf{D}^2)^{-1} \mathbf{D}^2$ .  $\square$

Next we show that the  $\beta_l$  from optimal scoring are also parallel to the columns of  $\mathbf{V}$

**Proposition 12.2.** *The optimal scoring problem as in Eq. (12.24) has solution  $\hat{\mathbf{\Theta}} = \mathbf{U}$ .*

*Proof.* From the first constraint of GSVD, obviously  $\mathbf{U}^T \mathbf{Y}^T \mathbf{Y} \mathbf{U} = \mathbf{I}$ . from Eq. (12.32a),  $\mathbf{U}$  diagonalizes  $\mathbf{Y}^T \mathbf{S} \mathbf{Y}$  by  $\mathbf{U}^T \mathbf{Y}^T \mathbf{S} \mathbf{Y} \mathbf{U} = \mathbf{D}^2$ .  $\square$

Consequently, we have  $\hat{\mathbf{B}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y} \mathbf{U} = \mathbf{V} \mathbf{D}$ . We can see that  $v_l$  (columns of  $\hat{\mathbf{A}}$ ) and  $\beta_l$  (columns of  $\hat{\mathbf{B}}$ ) differ by only a diagonal matrix  $(\mathbf{I} - \mathbf{D}^2)^{-1} \mathbf{D}$ .

### Ex. 12.9

The reduced features are simply

$$\mathbf{X}^* = \mathbf{X} \hat{\mathbf{B}} = \mathbf{S} \mathbf{Y} \quad (12.36)$$

therefore the optimal scoring can be computed by

$$\max_{\boldsymbol{\Theta}} \boldsymbol{\Theta}^T \mathbf{Y}^T [\mathbf{S} \mathbf{Y} (\mathbf{Y}^T \mathbf{S} \mathbf{Y})^{-1} \mathbf{Y}^T \mathbf{S}^T] \mathbf{Y} \boldsymbol{\Theta} \quad (12.37)$$

$$\text{s.t. } \boldsymbol{\Theta}^T \mathbf{Y}^T \mathbf{Y} \boldsymbol{\Theta} = N \mathbf{I} \quad (12.38)$$

with trivial manipulations one can see that the objective function is exactly the same as optimal scoring on original features.

### Ex. 12.10

The derivation for the general  $K$ -class GDA can be found in [?]. The kernel LDA (Fisher) is in the form of

$$\max_{\mathbf{a}} \frac{\mathbf{a}^T \mathbf{B} \mathbf{a}}{\mathbf{a}^T \mathbf{W} \mathbf{a}} \quad (12.39)$$

where

$$\mathbf{B} = (\bar{\mathbf{h}}_1 - \bar{\mathbf{h}}_2)(\bar{\mathbf{h}}_1 - \bar{\mathbf{h}}_2)^T \quad (12.40)$$

$$\bar{\mathbf{h}}_1 = \frac{1}{N_1} \sum_{i \in \mathcal{C}_1} \mathbf{h}(x_i) \quad (12.41)$$

$$\bar{\mathbf{h}}_2 = \frac{1}{N_2} \sum_{i \in \mathcal{C}_2} \mathbf{h}(x_i) \quad (12.42)$$

$$(12.43)$$

(up to constant) and  $N_1 = |\mathcal{C}_1|$ ,  $N_2 = |\mathcal{C}_2|$  are the numbers of data points in class 1, 2, respectively. The discriminant vector  $\mathbf{a}$  is a linear combination

$$\mathbf{a} = \sum_{i=1}^N \alpha_i \mathbf{h}(x_i) \quad (12.44)$$

therefore

$$\mathbf{a}^T \mathbf{B} \mathbf{a} = \boldsymbol{\alpha}^T (\mathbf{k}_1 - \mathbf{k}_2)(\mathbf{k}_1 - \mathbf{k}_2)^T \boldsymbol{\alpha} \quad (12.45)$$

where

$$\{\mathbf{k}_1\}_i = \frac{1}{N_1} \sum_{j \in \mathcal{C}_1} K_{ij}, \quad \{\mathbf{k}_2\}_i = \frac{1}{N_2} \sum_{j \in \mathcal{C}_2} K_{ij}, \quad i = 1, \dots, N \quad (12.46)$$

On the other hand,  $\mathbf{W} = \mathbf{W}_h + \gamma \mathbf{I}$ , where

$$\mathbf{W}_h = \sum_{i \in \mathcal{C}_1} (\mathbf{h}(x_i) \mathbf{h}(x_i)^T - \bar{\mathbf{h}}_1 \bar{\mathbf{h}}_1^T) + \sum_{i \in \mathcal{C}_2} (\mathbf{h}(x_i) \mathbf{h}(x_i)^T - \bar{\mathbf{h}}_2 \bar{\mathbf{h}}_2^T) \quad (12.47)$$

(up to constant) therefore

$$\mathbf{a}^T \mathbf{W}_h \mathbf{a} = \boldsymbol{\alpha}^T \mathbf{K}^2 \boldsymbol{\alpha} - N_1 \boldsymbol{\alpha}^T \mathbf{k}_1 \mathbf{k}_1^T \boldsymbol{\alpha} - N_2 \boldsymbol{\alpha}^T \mathbf{k}_2 \mathbf{k}_2^T \boldsymbol{\alpha} \quad (12.48)$$

and  $\mathbf{a}^T \mathbf{a} = \boldsymbol{\alpha}^T \mathbf{K} \boldsymbol{\alpha}$ . Consequently, the model depend on  $h(\cdot)$  only via the  $N$ -by- $N$  matrix  $\mathbf{K}$ .

### Ex. 12.11

(a)

$$\begin{aligned} P(X = x | G = k) &= \frac{P(X = x, G = k)}{\int_x P(X = x, G = k) dx} \\ &= \frac{\sum_{r=1}^R \pi_r P_r(G = k) \phi(x; \mu_r, \Sigma)}{\sum_{r=1}^R \pi_r P_r(G = k) \int_x \phi(x; \mu_r, \Sigma) dx} \\ &= \frac{\sum_{r=1}^R \pi_r P_r(G = k) \phi(x; \mu_r, \Sigma)}{\sum_{r=1}^R \pi_r P_r(G = k)} \end{aligned} \quad (12.49)$$

Compare with (12.59), we can see that, by setting

$$\pi_{kr} = \frac{\sum_{r=1}^R \pi_r P_r(G = k)}{\sum_{r=1}^R \pi_r P_r(G = k)}, \quad R_k = R, \quad \mu_{kr} = \mu_r \quad (12.50)$$

MDA2 is a generalization of MDA.

(b) E-step: compute the responsibility of subclass  $c_{kr}$  within class  $k$  for each class- $k$  observation ( $g_i = k$ ):

$$W(c_{kr} | x_i, g_i) = \frac{\pi_{kr} \phi(x_i; \mu_r, \Sigma)}{\sum_{r=1}^R \pi_{kr} \phi(x_i; \mu_r, \Sigma)} \quad (12.51)$$

M-step: MLE on  $\mu_r$  and  $\Sigma$

$$\mu_r = \frac{\sum_{k=1}^K \sum_{i:g_i=k} W(c_{kr}|x_i, g_i) x_i}{\sum_{k=1}^K \sum_{i:g_i=k} W(c_{kr}|x_i, g_i)} \quad (12.52)$$

$$\Sigma = \frac{\sum_{k=1}^K \sum_{i:g_i=k} \sum_{r=1}^R W(c_{kr}|x_i, g_i) (x_i - \mu_r)(x_i - \mu_r)^T}{\sum_{k=1}^K \sum_{i:g_i=k} \sum_{r=1}^R W(c_{kr}|x_i, g_i)} \quad (12.53)$$

(c) ???

## Chapter 13

### Prototype Methods and Nearest-Neighbors

#### Ex. 13.1

Again  $k = 1, \dots, K$  are the indices of clusters/classes,  $r = 1, \dots, R$  are the indices of cluster centers/Gaussian components.

For each predictor labeled to class  $k$ , namely  $\{x_i | g_i = k\}$ , in terms of the E-step,

- For EM algorithm, its responsibility to component  $r$  is evaluated as

$$\gamma_{kr}(x_i) = \frac{\pi_{kr} \phi(x_i; \mu_{kr}, \Sigma)}{\sum_{s=1}^R \pi_{kr} \phi(x_i; \mu_{ks}, \Sigma)} \quad (13.1)$$

- For k-means algorithm, each predictor belongs to exactly 1 cluster center, therefore its counterpart of responsibility is binary-valued:

$$\gamma_{kr}(x_i) = \begin{cases} 1 & \text{if } \|x_i - \mu_{kr}\| \leq \|x_i - \mu_{ks}\|, s = 1, \dots, R \\ 0 & \text{otherwise} \end{cases} \quad (13.2)$$

In terms of the M-step,

- For EM algorithm, the component means are updated as a weighted average

$$\mu_{kr} = \frac{\sum_{i:g_i=k} \gamma_{kr}(x_i) x_i}{\sum_{i:g_i=k} \gamma_{kr}} \quad (13.3)$$

and the mixing probability is updated as

$$\pi_{kr} = \sum_{i:g_i=k} \frac{\gamma_{kr}(x_i)}{N_k} \quad (13.4)$$

where  $N_k$  is the number of records labeled to class  $k$ .

- For k-means, the cluster center is taken as an unweighted average over all  $x_i$  closest to it. Using the binary-valued responsibility definition,  $\mu_{kr}$  is exactly the same as Eq. (13.3).

To draw a connection between EM and k-means, as  $\sigma \rightarrow 0$ ,  $\phi(x_i; \mu_{kr}, \Sigma) \gg \phi(x_i; \mu_{ks}, \Sigma)$  if  $\|x_i - \mu_{kr}\| < \|x_i - \mu_{ks}\|$ , therefore the responsibility for EM approaches that of k-means. And  $\pi_{kr}$  becomes the proportion of points in  $\{x_i | g_i = k\}$  that are closer to  $r$  than any other components centers.



**Ex. 13.2**

This problem is similar as Ex 2.3. Denote  $P_0(r, N)$  as the probability of the following event: “Among the  $N$  i.i.d uniformly distributed points, there is none within the ball centered at 0 with radius of  $r$ .” thus  $P_0(r, N) = P_0(r)^N$ , where  $P_0(r) = P_0(r, 1)$ . Since the points are uniformly distributed within the  $p$ -dim cube of edge length 1,  $P_0(r, 1)$  simply equals to the ratio between the volume of spaces out of the  $r$ -ball but within the cube, and the volume of the cube. Consequently,

$$P_0(r, N) = (1 - v_p r^p)^N \quad (13.5)$$

The median  $R_{med}$  satisfy  $P_0(R_{med}, N) = 1/2$ , therefore

$$R_{med} = v_p^{-1/p} (1 - 2^{-1/N})^{1/p} \quad (13.6)$$

**Ex. 13.3**

Since  $\sum_{k \neq k^*} p_k(x) = 1 - p_{k^*}(x)$

$$\begin{aligned} \sum_{k=1}^K p_k(x)(1 - p_k(x)) &= p_{k^*}(x)(1 - p_{k^*}(x)) + \sum_{k \neq k^*} p_k(x)(1 - p_k(x)) \\ &= p_{k^*}(x)(1 - p_{k^*}(x)) + (1 - p_{k^*}(x)) - \sum_{k \neq k^*} p_k(x)^2 \\ &\leq p_{k^*}(x)(1 - p_{k^*}(x)) + (1 - p_{k^*}(x)) - \frac{(\sum_{k \neq k^*} p_k(x))^2}{K-1} \\ &= (1 - p_{k^*}(x)) - \frac{K}{K-1} (1 - p_{k^*}(x))^2 \end{aligned} \quad (13.7)$$

where we made use of Cauchy-Schwarz inequality.

**Ex. 13.4**

(a)

$$\begin{aligned} \mathbf{A} &= \mathbf{Q}\mathbf{R} \\ &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \end{aligned} \quad (13.8)$$

where  $\theta$  represents the angle of rotation,  $a$ ,  $b$  represent the scale in  $x$  and  $y$  direction and  $\lambda$  represent the shear.

(b) Denote  $\mathbf{J}$  as the Jacobian (1-by-2) of  $F$  at  $c + x_0 + \mathbf{A}(\mathbf{x} - \mathbf{x}_0)$ , we have

$$\frac{\partial F}{\partial \theta} = \mathbf{J} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \mathbf{A}(\mathbf{x} - \mathbf{x}_0) \quad (13.9a)$$

$$\frac{\partial F}{\partial a} = \mathbf{J} \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \begin{bmatrix} 1 & \lambda \end{bmatrix} (\mathbf{x} - \mathbf{x}_0) \quad (13.9b)$$

$$\frac{\partial F}{\partial b} = \mathbf{J} \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} (\mathbf{x} - \mathbf{x}_0) \quad (13.9c)$$

$$\frac{\partial F}{\partial \lambda} = \mathbf{J} \begin{bmatrix} a \cos \theta \\ a \sin \theta \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix} (\mathbf{x} - \mathbf{x}_0) \quad (13.9d)$$

(c) It seems that the key to this procedure is to evaluate  $\mathbf{J}$  given a coordinate  $\mathbf{x}$ . Denote the 2-D kernel smoother as  $K(\mathbf{u}, \mathbf{v})$ , then we solve the locally weighted regression at  $\mathbf{x}$ :

$$\min_{\alpha(\mathbf{x}), \beta(\mathbf{x})} \sum_{i=1}^{256} K(\mathbf{x}, \mathbf{x}_i) [F(\mathbf{x}_i) - \alpha(\mathbf{x}) - \beta(\mathbf{x})^T \mathbf{x}_i] \quad (13.10)$$

Then we can use  $\beta(\mathbf{x})^T$  as  $\mathbf{J}(\mathbf{x})$  to compute the tangent space.

### Ex. 13.5

Since

$$N \operatorname{tr}(\bar{\mathbf{B}}\bar{\mathbf{B}}^T) = \sum_{i=1}^N \operatorname{tr}(\mathbf{B}_i\bar{\mathbf{B}}^T) = \sum_{i=1}^N \operatorname{tr}(\bar{\mathbf{B}}\mathbf{B}_i^T) \quad (13.11a)$$

$$N \operatorname{tr}(\bar{\mathbf{B}}\mathbf{M}^T) = N \operatorname{tr}(\mathbf{M}\bar{\mathbf{B}}^T) = \sum_{i=1}^N \operatorname{tr}(\mathbf{B}_i\mathbf{M}^T) = \sum_{i=1}^N \operatorname{tr}(\mathbf{M}\mathbf{B}_i^T), \quad (13.11b)$$

it is easy to show that

$$\sum_{i=1}^N \operatorname{tr}[(\mathbf{B}_i - \mathbf{M})^2] = \sum_{i=1}^N \operatorname{tr}[(\mathbf{B}_i - \bar{\mathbf{B}})^2] + N \operatorname{tr}[(\mathbf{M} - \bar{\mathbf{B}})^2] \quad (13.12)$$

Therefore the rank- $L$  approximation of  $\mathbf{B}_i$  is equivalent to the rank- $L$  approximation of  $\bar{\mathbf{B}}$ , namely  $\bar{\mathbf{B}}_{[L]}$ .

### Ex. 13.6

( $\mathbf{L}_j, j = 1, \dots, M$  are the coordinates of the “black” parts of the cursive letter.) As the

optimal  $\mathbf{A}_j = \mathbf{V}^T \mathbf{L}_j$ , we have

$$\begin{aligned}
 \sum_{j=1}^M \min_{\mathbf{A}_j} \|\mathbf{L}_j - \mathbf{V} \mathbf{A}_j\|^2 &= \sum_{j=1}^M \|(\mathbf{I} - \mathbf{V} \mathbf{V}^T) \mathbf{L}_j\|^2 \\
 &= \sum_{j=1}^M \text{tr} [(\mathbf{I} - \mathbf{V} \mathbf{V}^T) \mathbf{L}_j \mathbf{L}_j^T (\mathbf{I} - \mathbf{V} \mathbf{V}^T)] \\
 &= \sum_{j=1}^M \text{tr} [(\mathbf{I} - \mathbf{V} \mathbf{V}^T) \mathbf{L}_j \mathbf{L}_j^T] \\
 &= \sum_{j=1}^M \text{tr} [\mathbf{U}_j \mathbf{\Sigma}_j^2 \mathbf{U}_j^T] - \text{tr} \left[ \mathbf{V}^T \left( \sum_{j=1}^M \mathbf{U}_j \mathbf{\Sigma}_j^2 \mathbf{U}_j^T \right) \mathbf{V} \right] \quad (13.13)
 \end{aligned}$$

where  $\mathbf{L}_j = \mathbf{U}_j \mathbf{\Sigma}_j \mathbf{V}_j^T$  is the SVD of  $\mathbf{L}_j$ . Therefore  $\mathbf{V}$  corresponds to the 2 largest eigenvectors of  $\sum_{j=1}^M \mathbf{U}_j \mathbf{\Sigma}_j^2 \mathbf{U}_j^T$ .

For the alternative approach,

$$\begin{aligned}
 \sum_{j=1}^M \min_{\mathbf{A}_j} \|\mathbf{L}_j \mathbf{A}_j^T - \mathbf{V}\|^2 &= \sum_{j=1}^M \|(\mathbf{I} - \mathbf{S}_j) \mathbf{V}\|^2 \\
 &= \sum_{j=1}^M \text{tr} [\mathbf{V}^T (\mathbf{I} - \mathbf{S}_j) (\mathbf{I} - \mathbf{S}_j)^T \mathbf{V}] \\
 &= \sum_{j=1}^M \text{tr} [\mathbf{V}^T (\mathbf{I} - \mathbf{S}_j) \mathbf{V}] \\
 &= 2M - \text{tr} \left[ \mathbf{V}^T \left( \sum_{j=1}^M \mathbf{S}_j \right) \mathbf{V} \right] \quad (13.14)
 \end{aligned}$$

where  $\mathbf{S}_j = \mathbf{L}_j (\mathbf{L}_j^T \mathbf{L}_j)^{-1} \mathbf{L}_j^T = \mathbf{U}_j \mathbf{U}_j^T$ . Therefore  $\mathbf{V}$  corresponds to the 2 largest eigenvectors of  $\sum_{j=1}^M \mathbf{U}_j \mathbf{U}_j^T$ .

**Ex. 13.7 (Program)**

**Ex. 13.8 (Program)**

# Chapter 14

## Unsupervised Learning

**Ex. 14.1**

$$\begin{aligned}
 d_e(z_i, z_{i'}) &= \sum_{l=1}^p (z_{il} - z_{i'l})^2 \\
 &= \sum_{l=1}^p (z_{il} - z_{i'l})^2 \frac{w_l}{\sum_{j=1}^p w_j} (x_{il} - x_{i'l})^2 \\
 &= \frac{\sum_{l=1}^p w_l (x_{il} - x_{i'l})^2}{\sum_{j=1}^p w_j} \\
 &= d_e^{(w)}(x_i, x_{i'})
 \end{aligned} \tag{14.1}$$

**Ex. 14.2**

(a) The log-likelihood of a given record  $\mathbf{x}_i$  is

$$l(\theta; \mathbf{x}_i) = -\frac{1}{2} \log |\mathbf{L}| - \frac{p}{2} \log 2\pi - \frac{p}{2} \log \sigma^2 + \log \left[ \sum_{k=1}^K \pi_k \exp(\mathbf{x}_i - \boldsymbol{\mu}_k)^T (\sigma^2 \mathbf{L})^{-1} (\mathbf{x}_i - \boldsymbol{\mu}_k) \right], \tag{14.2}$$

and the log-likelihood over the entire data set is simply  $l(\theta; \mathbf{X}) = \sum_{i=1}^N l(\theta; \mathbf{x}_i)$

(b) Suppose we enlarge the dataset with latent variable  $\boldsymbol{\Delta}$  ( $N$ -by- $K$ ) such that  $\Delta_{ik} = 1$  if  $\mathbf{x}_i$  is associated with the  $k$ -th component and 0 otherwise. Each  $\mathbf{x}_i$  is associated with exactly one  $k$ . The the loglikelihood on  $\mathbf{x}_i, \boldsymbol{\Delta}_i$  becomes

$$\begin{aligned}
 l(\theta; \mathbf{x}_i, \boldsymbol{\Delta}_i) &= -\frac{1}{2} \log |\mathbf{L}| - \frac{p}{2} \log 2\pi - \frac{p}{2} \log \sigma^2 \\
 &\quad + \sum_{k=1}^K \Delta_{ik} \left[ \log \pi_k - \frac{1}{2} (\mathbf{x}_i - \boldsymbol{\mu}_k)^T (\sigma^2 \mathbf{L})^{-1} (\mathbf{x}_i - \boldsymbol{\mu}_k) \right]
 \end{aligned} \tag{14.3}$$

therefore

$$l(\theta; \mathbf{X}, \boldsymbol{\Delta}) = C - \frac{Np}{2} \log \sigma^2 + \sum_{i=1}^N \sum_{k=1}^K \Delta_{ik} \left[ \log \pi_k - \frac{1}{2} (\mathbf{x}_i - \boldsymbol{\mu}_k)^T (\sigma^2 \mathbf{L})^{-1} (\mathbf{x}_i - \boldsymbol{\mu}_k) \right] \tag{14.4}$$

Now we can formulate the EM-algorithm. We replace  $\Delta_{ik}$  with responsibility  $\gamma_{ik}$ . For the maximization step, we evaluate the MLE of  $\sigma^2$  and  $\mu_k$ . Since

$$\frac{\partial l(\theta; \mathbf{X}, \Delta)}{\partial \sigma^2} = -\frac{Np}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^N \sum_{k=1}^K \gamma_{ik} (\mathbf{x}_i - \mu_k)^T \mathbf{L}^{-1} (\mathbf{x}_i - \mu_k) \quad (14.5)$$

$$\frac{\partial l(\theta; \mathbf{X}, \Delta)}{\partial \mu_k} = \sum_{i=1}^N \gamma_{ik} (\mu_k - \mathbf{x}_i)^T (\sigma^2 \mathbf{L})^{-1} \quad (14.6)$$

By setting the partial derivative to 0, the MLE are

$$\hat{\mu}_k = \frac{\sum_{i=1}^N \gamma_{ik} \mathbf{x}_i}{\sum_{i=1}^N \gamma_{ik}} \quad (14.7)$$

$$\hat{\sigma}^2 = \frac{1}{Np} \sum_{i=1}^N \sum_{k=1}^K \gamma_{ik} (\mathbf{x}_i - \mu_k)^T \mathbf{L}^{-1} (\mathbf{x}_i - \mu_k) \quad (14.8)$$

and the MLE for  $\pi_k$  are solved by

$$\max_{\pi_k, l=1, \dots, K} \sum_{i=1}^N \sum_{k=1}^K \gamma_{ik} \log \pi_k \quad (14.9)$$

$$\text{s.t. } \sum_{k=1}^K \pi_k = 1 \quad (14.10)$$

therefore  $\hat{\pi}_k = \sum_{i=1}^N \gamma_{ik} / N$ .

For the expectation step, the responsibilities are updated as

$$\hat{\gamma}_{ik} = \frac{\pi_k \phi_k(\mathbf{x}_i)}{\sum_{l=1}^K \pi_l \phi_l(\mathbf{x}_i)} \quad (14.11)$$

where  $\phi_k(\cdot)$  is the PDF of  $\mathcal{N}(\mu_k, \sigma^2 \mathbf{L})$ .

(c) Pretty much the same as Ex 13.1. Now we are not dealing with classification so we don't need to treat  $\mathbf{x}_i$  with different labels separately.

**Ex. 14.3**

???

**Ex. 14.4 (Program)**

**Ex. 14.5 (Program)**

**Ex. 14.6 (Program)****Ex. 14.7**

$$\sum_{i=1}^N \|\mathbf{x}_i - \boldsymbol{\mu} - \mathbf{V}_q \boldsymbol{\lambda}_i\| = \|\mathbf{X} - \mathbf{1}_N \boldsymbol{\mu}^T - \boldsymbol{\Lambda} \mathbf{V}_q^T\|_F^2 \quad (14.12)$$

which is minimized when  $\hat{\boldsymbol{\Lambda}} = (\mathbf{X} - \mathbf{1}_N \boldsymbol{\mu}^T) \mathbf{V}_q$  given  $\mathbf{V}_q$  and  $\boldsymbol{\mu}$ . Denote the null space of  $\mathbf{V}_q$  is represented by  $\tilde{\mathbf{V}}_q$  where  $\tilde{\mathbf{V}}_q^T \tilde{\mathbf{V}}_q = \mathbf{I}_{p-q}$ . Now (14.50) becomes

$$\min_{\boldsymbol{\mu}, \mathbf{V}_q} \|(\mathbf{X} - \mathbf{1}_N \boldsymbol{\mu}^T) \tilde{\mathbf{V}}_q\|_F^2 \quad (14.13)$$

Taking partial derivative w.r.t  $\boldsymbol{\mu}$ , we can see that given  $\mathbf{V}_q$ , the optimal  $\boldsymbol{\mu}$  satisfy

$$\hat{\boldsymbol{\mu}} = \bar{\mathbf{x}} + \mathbf{V}_q \mathbf{b} \quad (14.14)$$

therefore  $\hat{\boldsymbol{\mu}} = \bar{\mathbf{x}}$  is an optimal solution for arbitrary  $\mathbf{V}_q$ .

**Ex. 14.8**

Since

$$\frac{\partial \|\mathbf{X}_2 - (\mathbf{X}_1 \mathbf{R}) + \mathbf{1} \boldsymbol{\mu}^T\|_F^2}{\partial \boldsymbol{\mu}} = 2N \boldsymbol{\mu}^T - 2\mathbf{1}^T \mathbf{X}_2 2\mathbf{1}^T \mathbf{X}_1 \mathbf{R} \quad (14.15)$$

by setting the partial derivative to 0, we have  $\hat{\boldsymbol{\mu}} = \bar{\mathbf{x}}_2 - \mathbf{R} \bar{\mathbf{x}}_1$ . Substitute this result into (14.56) we get

$$\min_{\mathbf{R}} \|\tilde{\mathbf{X}}_2 - \tilde{\mathbf{X}}_1 \mathbf{R}\|_F^2 \quad (14.16)$$

s.t.  $\mathbf{R}$  is orthogonal.

which is a orthogonal procustes problem (wiki). Since

$$\min_{\mathbf{R}} \|\tilde{\mathbf{X}}_2 - \tilde{\mathbf{X}}_1 \mathbf{R}\|_F^2 \Leftrightarrow \max_{\mathbf{R}} \text{tr}(\tilde{\mathbf{X}}_1^T \tilde{\mathbf{X}}_2 \mathbf{R}^T) \Leftrightarrow \max_{\mathbf{R}} \text{tr}(\mathbf{D} \mathbf{V}^T \mathbf{R}^T \mathbf{U}) \quad (14.17)$$

which is maximized when  $\mathbf{R} = \mathbf{U} \mathbf{V}^T$

**Ex. 14.9**

(14.115) should be Procrustes average with scaling

$$\min_{\{\beta_l, \mathbf{R}_l\}_1^L, \mathbf{M}} \sum_{l=1}^L \|\beta_l \mathbf{X}_l \mathbf{R}_l - \mathbf{M}\|_F^2 \quad (14.18)$$

This problem can be solved (sub-optimally) with alternating optimization

(1) Given  $\mathbf{M}$ , the  $L$  pairs of  $(\beta_l, \mathbf{R}_l)$  can be solved independently

$$\min_{\beta_l, \mathbf{R}_l} \|\beta_l \mathbf{X}_l \mathbf{R}_l - \mathbf{M}\|_F^2 \quad (14.19)$$

(a) Given  $\mathbf{R}_l$ ,  $\beta_l$  is optimized as

$$\hat{\beta}_l = \frac{\text{tr}(\mathbf{X}_l \mathbf{R}_l \mathbf{M}^T)}{\text{tr}(\mathbf{X}_l \mathbf{X}_l^T)} \quad (14.20)$$

(b) Given  $\beta_l$ ,  $\mathbf{R}_l$  is optimized (orthogonal procrustes problem) as

$$\hat{\mathbf{R}}_l = \mathbf{U}_l \mathbf{V}_l^T \quad (14.21)$$

where  $\beta_l \mathbf{X}_l^T \mathbf{M} = \mathbf{U}_l \mathbf{D}_l \mathbf{V}_l^T$  is SVD. However, we note that  $\mathbf{U}_l, \mathbf{V}_l$  does not actually depend on  $\beta_l$ . Therefore, we can evaluate  $\hat{\mathbf{R}}_l$  with the SVD of  $\mathbf{X}_l^T \mathbf{M}$  and then evaluate  $\hat{\beta}_l$ .

(2) Given  $\beta_l, \mathbf{R}_l, \mathbf{M}$  is simply optimized as the average

$$\hat{\mathbf{M}} = \frac{1}{L} \sum_{l=1}^L \beta_l \mathbf{X}_l \mathbf{R}_l \quad (14.22)$$

The above 2 steps are taken alternately until convergence.

#### Ex. 14.10

Given  $\mathbf{M}$ ,  $\mathbf{A}_l$  are optimized as  $\hat{\mathbf{A}}_l = (\mathbf{X}_l^T \mathbf{X}_l)^{-1} \mathbf{X}_l^T \mathbf{M}$ . Consequently, (14.60) is equivalent to

$$\min_{\mathbf{M}} \sum_{l=1}^L \|(\mathbf{I} - \mathbf{H}_l) \mathbf{M}\|_F^2 \quad (14.23)$$

s.t.  $\mathbf{M} \mathbf{M}^T = \mathbf{I}$ , where  $\mathbf{H}_l = \mathbf{X}_l (\mathbf{X}_l^T \mathbf{X}_l)^{-1} \mathbf{X}_l^T$ . This in turn is equivalent to

$$\max_{\mathbf{M}} \sum_{l=1}^L \text{tr}(\mathbf{M}^T \mathbf{H}_l \mathbf{M}) \quad (14.24)$$

s.t.  $\mathbf{M} \mathbf{M}^T = \mathbf{I}$ . As a result,  $\hat{\mathbf{M}}$  is the  $p$  largest eigen vectors of  $\sum_{l=1}^L \mathbf{H}_l$ .

**Ex. 14.11**

???

**Ex. 14.12**

(a)

$$\begin{aligned}
& \sum_{i=1}^N \|\mathbf{x}_i - \mathbf{\Theta} \mathbf{V}^T \mathbf{x}_i\|^2 \\
&= \|\mathbf{X}(\mathbf{I} - \mathbf{V} \mathbf{\Theta}^T)\|_F^2 \\
&= \text{tr}[\mathbf{X}(\mathbf{\Theta} - \mathbf{V})(\mathbf{\Theta} - \mathbf{V})^T \mathbf{X}^T] + \text{tr}[\mathbf{X} \mathbf{X}^T - \mathbf{X} \mathbf{\Theta} \mathbf{\Theta}^T \mathbf{X}] \quad (14.25)
\end{aligned}$$

where the second term is not dependent on  $\mathbf{V}$  and the first term equals to  $\sum_{i=1}^N \|\mathbf{\Theta}^T \mathbf{x}_i - \mathbf{V}^T \mathbf{x}_i\|^2$ . Consequently, the minimization of (14.71) w.r.t  $\mathbf{V}$  becomes

$$\min_{\{\mathbf{v}_k\}_{k=1}^K} \sum_{k=1}^K \left[ \sum_{i=1}^N \|\boldsymbol{\theta}_k^T \mathbf{x}_i - \mathbf{v}_k^T \mathbf{x}_i\|_2^2 + \lambda \|\mathbf{v}_k\|_2^2 + \lambda_{1k} \|\mathbf{v}_k\|_1 \right] \quad (14.26)$$

which can be solved as  $K$  separate elastic net regression problems.

(b) We rewrite

$$\begin{aligned}
& \|\mathbf{X}(\mathbf{I} - \mathbf{V} \mathbf{\Theta}^T)\|_F^2 \\
&= \text{tr}[\mathbf{X} \mathbf{X}^T - \mathbf{X} \mathbf{V} \mathbf{V}^T \mathbf{X}^T] - 2 \text{tr}[\mathbf{\Theta}^T \mathbf{X}^T \mathbf{X} \mathbf{V}] \quad (14.27)
\end{aligned}$$

Since the rest part of (14.71) is not dependent on  $\mathbf{\Theta}$ , the minimization of (14.71) w.r.t  $\mathbf{\Theta}$  is equivalent to

$$\begin{aligned}
& \max_{\mathbf{\Theta}} \text{tr}(\mathbf{\Theta}^T \mathbf{M}) \\
& \text{s.t. } \mathbf{\Theta}^T \mathbf{\Theta} = \mathbf{I} \quad (14.28)
\end{aligned}$$

where  $\mathbf{M} = \mathbf{X}^T \mathbf{X} \mathbf{V}$ . This has the same form as the Procrustes problem in Ex 14.8, therefore its solution is  $\mathbf{\Theta} = \mathbf{U} \mathbf{Q}^T$ .

**Ex. 14.13 (Program)****Ex. 14.14**

Denote  $\mathbf{D}_A = \text{diag}(\{\mathbf{a}_j^T \mathbf{a}_j\}_{j=1}^p)$ , then

$$\mathbf{P} = \mathbf{D}_A^{-1/2} \boldsymbol{\Sigma} \mathbf{D}_A^{-1/2} = \mathbf{P}_A + \mathbf{D}_A^{-1} \mathbf{D}_\epsilon \quad (14.29)$$



where  $\mathbf{P}_A$  is the correlation matrix

$$\{\mathbf{P}_A\}_{ij} = \frac{\mathbf{a}_i^T \mathbf{a}_j}{\|\mathbf{a}_i\| \|\mathbf{a}_j\|} \quad (14.30)$$

**Ex. 14.15 (Program)**

**Ex. 14.16**

Since  $\mathbf{Z} = \tilde{\mathbf{K}}\mathbf{U}\mathbf{D}^{-1}$ , therefore

$$z_{im} = \sum_{j=1}^N \tilde{K}(x_i, x_j) u_{jm} d_m^{-1} \quad (14.31)$$

where  $\tilde{K}(x_i, x_j)$  differs from  $K(x_i, x_j)$  only in centering. For a new observation  $x_0$ , its mapping to the  $m$ -th component is

$$\langle \tilde{\phi}(x_0), \sum_{j=1}^N \alpha_{jm} \tilde{\phi}(x_j) \rangle = \sum_{j=1}^N \alpha_{jm} \langle \tilde{\phi}(x_0), \tilde{\phi}(x_j) \rangle = \sum_{j=1}^N \alpha_{jm} \tilde{K}(x_0, x_j) \quad (14.32)$$

which differs from  $\sum_{j=1}^N \alpha_{jm} K(x_0, x_j)$  only in centering.

**Ex. 14.17**

Denote  $\mathbf{c} = [c_1, \dots, c_N]^T$ . First we note

$$\|g_1(x)\|_{\mathcal{H}_K}^2 = \sum_{i=1}^N \sum_{j=1}^N c_i c_j K(x_i, x_j) = \mathbf{c}^T \mathbf{K} \mathbf{c} \quad (14.33)$$

Secondly, we have

$$\begin{aligned} \text{Var}_{\mathcal{T}} g_1(X) &= \frac{1}{N} \sum_{k=1}^N g_1(x_k)^2 \\ &= \frac{1}{N} \sum_{k=1}^N \sum_{i=1}^N \sum_{j=1}^N c_i c_j K(x_k, x_i) K(x_k, x_j) \\ &= \frac{1}{N} \mathbf{c}^T \mathbf{K} \mathbf{K} \mathbf{c} \end{aligned} \quad (14.34)$$

Since  $\mathbf{K} = \mathbf{U}\mathbf{D}^2\mathbf{U}^T$ , (14.66) can be rewritten as

$$\begin{aligned} &\max \mathbf{c}^T \mathbf{U} \mathbf{D}^4 \mathbf{U}^T \mathbf{c} \\ &\text{s.t. } \mathbf{c}^T \mathbf{U} \mathbf{D}^2 \mathbf{U}^T \mathbf{c} = 1 \end{aligned} \quad (14.35)$$

Denote  $\mathbf{a} = \mathbf{D}\mathbf{U}^T\mathbf{c}$ , then the optimal solution must satisfy  $\hat{\mathbf{a}} = [1, 0, \dots, 0]^T$ , therefore  $\hat{c} = \mathbf{u}_1/d_1$ .  $g_2, \dots, g_M$  can be derived in a similar manner.

**Ex. 14.18**

Consider the stationary condition on  $\theta_0$ , we have

$$\frac{\partial l}{\partial \theta_0} = 1 - \int \phi(t) \exp(\theta_0 + \theta_1 t + \theta_2 t^2) dt = 0 \quad (14.36)$$

also since  $\phi(t) \exp(\theta_0 + \theta_1 t + \theta_2 t^2) > 0$ , it is a probability density function.

Consider the stationary condition on  $\theta_1$ , we have

$$\frac{\partial l}{\partial \theta_1} = \frac{1}{N} \sum_{i=1}^N s_i - \int t \phi(t) \exp(\theta_0 + \theta_1 t + \theta_2 t^2) dt = 0 \quad (14.37)$$

since  $\sum_{i=1}^N s_i = 0$ , this condition suggest that  $\phi(t) \exp(\theta_0 + \theta_1 t + \theta_2 t^2)$  has zero mean.

Consider the stationary condition on  $\theta_2$ , we have

$$\frac{\partial l}{\partial \theta_2} = \frac{1}{N} \sum_{i=1}^N s_i^2 - \int t^2 \phi(t) \exp(\theta_0 + \theta_1 t + \theta_2 t^2) dt = 0 \quad (14.38)$$

since  $\sum_{i=1}^N s_i^2/N = 1$ , this condition suggest that  $\phi(t) \exp(\theta_0 + \theta_1 t + \theta_2 t^2)$  has unit variance.  
(???)

**Ex. 14.19**

$$\sum_{j=1}^p \sum_{i=1}^N \log \phi(\mathbf{a}_j^T \mathbf{x}_i) = -\frac{pN}{2} \log 2\pi - \frac{1}{2} \|\mathbf{A}\mathbf{X}\|_F^2. \quad (14.39)$$

Since  $\|\mathbf{A}\mathbf{X}\|_F^2 = \|\mathbf{X}\|_F^2$  for any orthogonal  $\mathbf{A}$ , this term does not depend on  $\mathbf{A}$ .

**Ex. 14.20**

Since

$$\frac{\partial g}{\partial a} = \mathbb{E}[X g'(a^T X)] \quad (14.40)$$

$$\frac{\partial^2 g}{\partial a \partial a^T} = \mathbb{E}[X X^T g''(a^T X)] \approx \mathbb{E}[g''(a^T X)] I \quad (14.41)$$

the Newton update is

$$a \leftarrow a - (\mathbb{E}[g''(a^T X)])^{-1} \mathbb{E}[X g'(a^T X)] \quad (14.42)$$

Since  $a$  needs to be normalized to ensure  $\|a\| = 1$  anyway, the right hand side of the above equation can be multiplied with a positive constant  $-\mathbb{E}[g''(a^T X)]$  (followed by a normalization), resulting in

$$a \leftarrow \mathbb{E}[Xg'(a^T X)] - \mathbb{E}[g''(a^T X)]a \quad (14.43)$$

**Ex. 14.21**

Since there are  $m$  connected components in the graph,  $\mathbf{L}$  can be transformed into a block-diagonal matrix

$$\mathbf{L} = \text{diag}(\mathbf{L}_1, \dots, \mathbf{L}_m) \quad (14.44)$$

where  $\mathbf{L}_j = \mathbf{G}_j - \mathbf{W}_j$ . Since  $\mathbf{L}_m \mathbf{1} = \mathbf{0}$ ,  $\mathbf{L}$  has  $m$  eigenvectors corresponding to eigenvalue of 0, which are the same as the permuted indicator vectors  $I_{A_1}, \dots, I_{A_m}$ .

**Ex. 14.22**

(a)

$$\begin{aligned} \mathbf{1}^T \mathbf{p} &= (1-d)\mathbf{1}^T \mathbf{e} + d\mathbf{1}^T \mathbf{L} \mathbf{D}_c^{-1} \mathbf{p} \\ &= (1-d)N + d\mathbf{c}^T \mathbf{D}_c^{-1} \mathbf{p} \\ &= (1-d)N + d\mathbf{1}^T \mathbf{p} \end{aligned} \quad (14.45)$$

therefore  $\mathbf{1}^T \mathbf{p} = N$ .

(b) (Program)

**Ex. 14.23**

(a) Since  $\log(\cdot)$  is concave, according to Jensen's inequality

$$\begin{aligned} \sum_{k=1}^r c_k \log(y_k/c_k) &= \frac{1}{\sum_{k=1}^r c_k} \sum_{k=1}^r c_k \log(y_k/c_k) \\ &\leq \log\left(\frac{\sum_{k=1}^r y_k}{\sum_{k=1}^r c_k}\right) \\ &= \log\left(\sum_{k=1}^r y_k\right) \end{aligned} \quad (14.46)$$

where equality holds iff  $c_k = 1/r$ .

(b)

$$g(\mathbf{W}, \mathbf{H} | \mathbf{W}^s, \mathbf{H}^s) = \sum_{i=1}^N \sum_{j=1}^p \sum_{k=1}^r x_{ij} \frac{a_{ikj}^s}{b_{ij}^s} \log \left( \frac{b_{ij}^s}{a_{ikj}^s} w_{ik} h_{kj} \right) - \sum_{i=1}^N \sum_{j=1}^p \sum_{k=1}^r w_{ik} h_{kj} \quad (14.47)$$

minorizes  $L(\mathbf{W}, \mathbf{H})$

(c) The stationary conditions are

$$\frac{\partial g}{\partial w_{ik}} = \sum_{j=1}^p x_{ij} \frac{a_{ikj}^s}{b_{ij}^s} - \sum_{j=1}^p h_{kj} = 0 \quad (14.48)$$

$$\frac{\partial g}{\partial h_{kj}} = \sum_{i=1}^N x_{ij} \frac{a_{ikj}^s}{b_{ij}^s} - \sum_{i=1}^N w_{ik} = 0 \quad (14.49)$$

which are equivalent to

$$w_{ik} = \frac{\sum_{j=1}^p x_{ij} a_{ikj}^s / b_{ij}^s}{\sum_{j=1}^p h_{kj}} \quad (14.50)$$

$$h_{kj} = \frac{\sum_{i=1}^N x_{ij} a_{ikj}^s / b_{ij}^s}{\sum_{i=1}^N w_{ik}} \quad (14.51)$$

which are exactly the updating steps (14.74).

**Ex. 14.24**(a) When  $r = 1$ , we have  $a_{ikj}^s / b_{ij}^s = 1$ , therefore the updating steps are simplified as

$$w_i \leftarrow \frac{\sum_{j=1}^p x_{ij}}{\sum_{j=1}^p h_j}, \quad h_j \leftarrow \frac{\sum_{i=1}^N x_{ij}}{\sum_{i=1}^N w_i} \quad (14.52)$$

(b) From Eq. (14.52), for every two steps, the updating becomes

$$h_k \leftarrow \frac{\sum_{i=1}^N x_{ik}}{\sum_{i=1}^N w_i} \leftarrow \frac{\sum_{i=1}^N x_{ik}}{\sum_{i=1}^N \sum_{j=1}^p x_{ij}} \sum_{j=1}^p h_j \quad (14.53)$$

$$w_l \leftarrow \frac{\sum_{j=1}^p x_{lj}}{\sum_{j=1}^p h_j} \leftarrow \frac{\sum_{j=1}^p x_{lj}}{\sum_{i=1}^N \sum_{j=1}^p x_{ij}} \sum_{i=1}^N w_i \quad (14.54)$$

It is easy to see that throughout the updating,  $\sum_{j=1}^p h_j$  and  $\sum_{i=1}^N w_i$  remains constant, thus  $h_k$  and  $w_l$  remain constant. Consequently, the iteration is completely stationary. By enforcing  $\sum_{j=1}^p h_j \sum_{i=1}^N w_i = 1$ , the iteration has the explicit form as (14.122) for any  $c$ .

**Ex. 14.25 (Program)**



## Chapter 15

### Random Forests

## Chapter 16

### Ensemble Learning

## Chapter 17

### Undirected Graphical Models



## Chapter 18

### High-Dimensional Problems