

CS & IT ENGINEERING

DISCRETE MATHS
SET THEORY



Different Types of Functions

Lecture No. 4



By- SATISH YADAV SIR



TOPICS TO BE
COVERED

01 ONTO Function

02 BIJECTIVE FUNCTION

03 INVERTIBLE FUNCTION

04 TOTAL NUMBER OF
BIJECTIVE FUNCTIONS

05 IDENTITY FUNCTION

06 Different Types of Functions



Functions

$$f: A \rightarrow B$$

$$|A| = m \quad |B| = n$$


$$|A| \geq |B|$$

$$A > B \quad \text{OR} \quad A = B$$



$$Q.1. f: A \rightarrow B \quad |A| = 2 \quad |B| = 3$$

Total no. of onto functions = 0.



Functions

$$|A| = m \quad |B| = n$$

$$\sum_{i=0}^n (-1)^i * {}^nC_i * (n-i)^m$$

Q.2: $|A| = 7 \quad |B| = 4$

GATE: $|A| = n \quad |B| = 2$
 $f: A \rightarrow B$

$$\sum_{i=0}^2 (-1)^i * {}^2C_i * (2-i)^n$$

$${}^2C_0 (2-0)^n - {}^2C_1 (2-1)^n + {}^2C_2 (2-2)^n$$

$$1 \cdot 2^n - 2 \cdot 1^n = \underline{\underline{2^n - 2}}$$


Functions

$$f: A \rightarrow B$$

$$|A| = n \quad |B| = 2$$

$$\begin{aligned} \text{onto} &= \text{Total} - \text{non onto} \\ &= \text{r.s.}^{\text{L.S}} - 2 \end{aligned}$$

$$= 2^n - 2$$



Functions



$$m = |A| = 7 \quad |B| = 4 = n$$

$$\sum_{i=0}^4 (-1)^i * nC_i * (n-i)^m$$

$$nC_0(4-0)^7 - nC_1(4-1)^7 + nC_2(4-2)^7 - nC_3(4-3)^7 + \underline{nC_4(4-4)^7}$$

$$4^7 - 4 \cdot 3^7 + 6 \cdot 2^7 - 4 \cdot 1^7 + 0$$

$$\begin{aligned} 8400 &= 4^7 - 4 \cdot 3^7 + 6 \cdot 2^7 - 4 \\ &= 4(4^6 - 3^7 + 6 \cdot 2^6 - 1) \end{aligned}$$

Functions

How many ways we
 can arrange 7 people
 to 4 room such that
 none of the rooms are empty?

Ans: 8400

onto



rooms contains
 at least 1 person.

Functions


$$|A| = 4 \quad |B| = 3$$

$$3c_0(3-0)^4 - 3c_1(3-1)^4 + 3c_2(3-2)^4 - 3c_3(3-3)^4$$

$$= 3^4 - 3 \cdot 2^4 + 3 \cdot 1^4 - 0$$


$$= 81 - 3 \cdot 16 + 3 = 81 - 48 = \underline{36}$$

onto



$[ab]_1 [c]_2 [d]_3$

$[ab] [c] [d]$



$[ab]_2 [c]_1 [d]_3$

$[ab] [c] [d]$

Functions

onto:

objects \rightarrow distinguishable boxes. (36)

4 people \rightarrow 3 diff rooms. (onto)

4 people \rightarrow 3 identical rooms

onto
3!

Functions



a

b

c

$[\]_1$

$[\]_2$

onto = 6

diff boxes

$[ab]_1 [c]_2$

$[ab]_2 [c]_1$

identical boxes

$[ab] [c]$

$\frac{6}{2!}$

= 3



$[ac]_1 [b]_2$

$[ac]_2 [b]_1$

$[ac] [b]$

$[bc]_1 [a]_2$

$[bc]_2 [a]_1$

$[bc] [a]$

Functions



Bijective function (1:1 correspondance)

1:1 correspondance = 1:1 + onto

$$A = B$$

$$\underline{A \leq B \wedge A \geq B} \longrightarrow A = B$$

Functions



2^3 {
T T T \rightarrow 1 1 1
T F T \rightarrow 1 0 1
F F T \rightarrow 0 0 1
}

2^3

2^3 {

a	b	c	Subset
0	0	0	ϕ
1	0	0	{a}
0	1	0	{b}
0	0	1	{c}

}

2^3

Functions

$$|A| = |B| = n$$

Total no. of $1:1 \subset \rightarrow n!$

$$\{abc\} \quad \{123\}$$

$$f: A \rightarrow B$$


$$1:1 \rightarrow B^n$$

onto $\sum_{i=0}^n (-1)^i \cdot nC_i \cdot (n-i)^n$

$$1:1 \subset \rightarrow m=n \quad (n!)$$

Functions


Invertible function.
Inverse.



$$f: A \rightarrow B$$

$$\left\{ \begin{array}{l} f^{-1}: B \rightarrow A \\ f^{-1}(b) = a. \end{array} \right.$$

Functions



$$f(a) = 1$$

$$f(b) = 2$$

$$f(c) = 3$$

$$f^{-1}(1) = a$$

$$f^{-1}(2) = b$$

$$f^{-1}(3) = c$$

$$f^{-1}: B \rightarrow A$$




$$1 \rightarrow a$$

$$2 \rightarrow b$$


$$3 \rightarrow c$$




onto




Functions




not 1:1.
not onto.



1:1
not onto



not 1:1
but onto.



1:1 ✓
onto ✓

1:1 C.

Functions




$$f: A \rightarrow B$$

$$A_1, A_2 \subseteq A$$

$$a) \quad f(A_1 \cup A_2) = f(A_1) \cup f(A_2)$$

$$b) \quad f(\underbrace{A_1 \cap A_2}_{\emptyset \subseteq}) \subseteq f(A_1) \cap f(A_2)$$

Functions



$$f(A_1) = f(a, b) = 1$$

$$f(A_2) = f(c) = 2$$


$$\underline{f(A_1)} \cup \underline{f(A_2)}$$

$$\{1\} \cup \{2\} = \{1, 2\}$$

$$f(A_1 \cup A_2) = f(a, b, c) = \{1, 2\}$$


$$f(A_1 \cup A_2) = f(A_1) \cup f(A_2)$$

Functions



$A_1 = ab$
 $A_2 = c$

$\textcircled{=}$



$A_1 = ab$
 $A_2 = c$

$$f(A_1 \cap A_2) = f(\emptyset) = \emptyset$$

$$f(A_1) = f(ab) = \{1, 2\}$$

$$f(A_2) = f(c) = \{2\}$$

$$f(A_1) \cap f(A_2) = \{1, 2\} \cap \{2\} = \{2\}$$

$$\emptyset \subseteq \{2\}$$

Functions

