## **Breed Descriptor of Bhakarwal (Sheep)**

Male	Female				
Breeding Tract					
State: Jammu & Kashmir	Rajouri				
	Jammu				
-	Udhampur				
Location					
Longitude (deg.)	74 75.				
Latitude (deg.)	32.35 - 34.				
General Information					
Synonyms					
Origin	The name is derived from the nomadic tribe Bakerwals, which rears these sheep				
Major utility	Food - Meat; Fibre - Wool				
Comments on utility					
Comments on breeding tract  Adaptability to environment	Entirely migratory. Native tract constitutes the Shivalik formation consisting of undulating badly cut low lying to mountainous hills constituting very weak and fragile geographic formation. The Places where the breed is available at present are Jammu (Samba, Gogwal, Jajerkotli, Dhansal, Jindra, Arnia, Kakwakota, Sai, Kool Kalan); Udampur (Riasi, Thenol, Anji nala, Pouni, Bidda, Rud Patina, Thakrakot, Sansui nalla, Touth, Serout, Sanjantharan, Lamsara, Bagdhar); and Rajouri (Kalakot, Nowshera, Sunderbani Mogla, Pouni park, Jasota, Seot, treyat, Buddi).  Sturdy and excellent climber.				
	Sturdy and excer	Tent chimber.			
Population	V	D1-4	Outrania fana di		
	Year 2002	Population 2596	Other information 924 males and 1672 females of all age groups. Data collected through survey under NATP (MM) project by SKUAS&T, Srinagar		

	2004	2600	Source: Final Report, NATP (MM),		
			NBAGR, Karnal		
	2013	82334	Source: Estimated		
			Livestock Population		
			Breed Wise Based on		
			Breed Survey 2013.		
			Department of		
			Animal Husbandry,		
			Dairying & Fisheries,		
			Government of India,		
D.M.			New Delhi		
Management	E-4				
Management system	Extensive				
Mobility	Nomadic				
Feeding of adults  Comments on Management	Grazing				
Comments on Management	Animals are maintained on open grazing in the natural grasslands with no supplementary feeding. During winter, tree loppings and jungle hay is fed to animals. A large				
	11 0		<u> </u>		
	number of flocks migrate to Kashmir valley in the areas of Pahalgam, Shopian, Sonmarg, Minimarg, Baltal, Gurej, Sokhnai and Toss Maidan in summer for grazing. Flocks move down to low lying hills in winter. Flocks are				
	gathered in the ev				
			e open throughout the		
	year.	iais are kept in til	e open unoughout the		
Morphology	year.				
Colour	Generally white. Head is black in colour. Few black				
	coloured animals (about 5%) are also available.				
Horn shape and size	Horns are curved backward and downwards. Whitish grey to brown in colour and medium in size. Ewes are				
	generally polled b				
Visible characteristics	Medium sized animal, with a typical roman nose and black head. Ears are horizontal and directed backwards.				
	Body coat is wooly. Head, face and legs are devoid of				
	wool.				
	Male	Female	Overall		
Height (avg. cm.)	77.0	68.0			
Body Length (avg. cm.)	81.0	67.0			
Heart girth (avg. cm.)	108.0	88.0			
Body weight (avg. kg.)	42.0	33.0	2.0		
Birth weight (avg. kg.)			3.0		
Performance	A	Minimo	Mariana		
Litter size born	Average	Minimum	Maximum		
	1.0	23.0	29.0		
Age at first parturition		23.0	29.0		
(months)  Parturition interval	12.0				
(months)	12.0				
Fleece weight (kg/year)	1.5	0.8	1.8		
1 ICCC WEIGHT (Kg/year)	1.0	0.0	1.0		

Staple length (cm)	7.75	
Fibre Diameter (µ)	30.0	
Dressing percentage	55.0	
Any Peculiarity of the breed		