


Breed Descriptor of Kashmir Favorolla (Poultry-Fowl)



Breeding Tract			
State: Jammu & Kashmir	Pulwama		
	Kupwara		
	Budgam		
	Anantnag		
	Baramulla		
	Srinagar		
Location			
Longitude (deg.)	74. - 76.		
Latitude (deg.)	33. - 34.50		
General Information			
Synonyms	Kashir		
Origin	Basically an inter-breeding group of many distinct genotypes. The three important are normally feathered, nacked neck and gul hapuch (bottle jawed).		
Major utility	Food-egg and meat		
Comments on utility			
Comments on breeding tract	Native tract of distribution is Kashmir Valley		
Adaptability to environment	Adapted to adverse climatic conditions of freezing mercury levels, meager or no supplemntal ration. Disease resistant.		
Population			
	Year	Population	Other information
Management			
Management system	Backyard		
Mobility	Stationary		
Feeding of adults	Free Ranging		
Comments on Management	Scavenging with supplemental feeding. After harvesting the birds are let free in the fields. Housed only during night.		
Morphology			
	Male	Female	Overall
Birth weight (avg. kg.)			30.43
Adult weight(avg. kg)	1.19		

Plumage type	Normal		
Plumage pattern	Solid, stripped, Patchy, Spotted, Barred		
Plumage colour	Mixed shades of Black, Red, Green, Gold		
Comb type	Single		
Skin colour	White		
Shank colour	Yellow		
Egg shell colour	Light Brown		
Visible character	Feathered cap on head.		
Performance			
	Average	Minimum	Maximum
Age at first egg (months)	6.91		
Annual eggs production (avg. no.)		60.0	85.0
Eggs weight (avg. gm)	45.76		
Fertility (%)	76.72		
Hatching (%)	64.27		
Any Peculiarity of the breed	The typical features are feathered cap- tuft of feathers on head.The other physical traits at varied frequencies are: feathered shanks, tuft of feathers over earlobes, naked neck, silky frizzle, multiple spurs.		