Breed Descriptor of Punganur (Cattle)





Breeding Tract					
State: Andhra Pradesh	Chittoor				
Location					
Longitude (deg.)	78.40 - 79.10				
Latitude (deg.)	13.10 - 13.40				
General Information					
Synonyms					
Origin	This breed was developed by Rulers of Punganur. Hence				
	this breed is known as Punganur after the name of area.				
Major utility	Work - Draught and Transportation; Food - Milk				
Comments on utility	These animals are used for agricultural operations on				
	light soil. Bulloc	light soil. Bullocks are used for carts like tongas and for			
	special races.				
Comments on breeding	The breed tract is confined to the taluks of Punganur				
tract	and adjacent taluks of Vayalpad, Madnapalli and				
	Palamaner in Chittoor district of Andhra Pradesh.				
Adaptability to environment					
Population					
	Year	Population	Other information		
	1997	0	ANGRAU, Palmner		
			conducted survey in		
			the breeding tract		
			covering 150		
			villaf\ges and could		
			locate only 21 cows.		
	2013	2772	Source: Estimated		
			Livestock Population		
			Breed Wise Based on		
			Breed Survey 2013.		
			Department of		
			Animal Husbandry,		
			Dairying & Fisheries,		
			Government of India,		
			New Delhi		
Management		ı	1		

Management system	Semi-Intensive			
Mobility	Stationary			
Feeding of adults	Grazing, Fodder and Concentrate			
Comments on Management	Women and children of the family look after the animals.			
	Animals are housed in a small shed made with the help of			
	bamboos and stone pillars. Stacks of paddy straw and			
	groundnut form the roof of cattle shed. Animals are			
	housed only during the night and are grazed in the			
	wastelands in the nearby forests. Soon after the harvest			
	of paddy, animals are allowed to graze over paddy			
	stubbles. Paddy straw forms the major component of			
	fodder. In addition, sugarcane tops, ground nut halms,			
	fingermillet straw and mulberry leaves are fed to cattle.			
	Milking cows are fed concentrates at the time of milking.			
Morphology				
Colour	Punganur cattle are white, grey or light brown to dark			
	brown or red. White mixed with red or black colour			
	animals are also available. Combination of white with			
	brown or black patches is not uncommon.			
Horn shape and size	Horns are crescent shaped and often loose curving			
	backward and forward in males and lateral and forward			
	in females. Short to medium in size.			
Visible characteristics	Short stature			
	Male	Female	Overall	
Height (avg. cm.)	107.0	97.0		
Body Length (avg. cm.)	113.0	108.0		
Heart girth (avg. cm.)	151.0	128.0		
Body weight (avg. kg.)	240.0	170.0		
Birth weight (avg. kg.)	12.8	11.4	12.3	
Performance	г.	Ta et :	Is a	
	Average	Minimum	Maximum	
Age at first parturition	37.0			
(months)	1.4.00	100	25.0	
Parturition interval	14.88	10.0	27.0	
(months)	7 460	1010	11000	
Milk yield per lactation (kg)	546.0	194.0	1100.0	
Milk Fat (%)	5.0	3.2	10.0	
Any Peculiarity of the breed	Dwarf cattle			