


## Breed Descriptor of Gangatiri (Cattle)



Breeding Tract			
State: Bihar	Bhojpur		
State: Uttar Pradesh	Ballia		
	Ghazipur		
	Mirzapur		
	Varanasi		
Location			
Longitude (deg.)	83.38 - 84.39		
Latitude (deg.)	25.19 - 27.11		
General Information			
Synonyms	Eastern Haryana or Shahabadi		
Origin	Derived the name from its habitat, in the area nearby river Ganga		
Major utility	Draught, milk and manure		
Comments on utility	Medium milk producer and possess good draftability		
Comments on breeding tract	Ghazipur, Ballia, Varanasi, Mau, Mirzapur and adjoining districts of Uttar Pradesh and Bhabhua, Buxar and Bhojpur districts of Bihar		
Adaptability to environment	Gangatiri cattle are well adapted in low/medium input production system		
Population			
	Year	Population	Other information
	2007	375154	3,64,806 in Uttar Pradesh.Livestock Census 2007
Management			
Management system	Semi-Intensive		
Mobility	Stationary		
Feeding of adults	Grazing,Fodder and Concentrate		
Comments on Management	Developed mostly by small and marginal farmers of several communities like Yadav, Bhumihar, Brahmin, Rajbhar, Kshatriya, Patel, etc		
Morphology			
Colour	Complete white (Dhawar), Grey (Sokan)		

Horn shape and size	Medium sized. Emerges from side of the poll behind and above eyes in outward and curving upwards and inwards ending with pointed tips		
Visible characteristics	Forehead is Prominent, straight and broad with shallow groove in the middle. Eyelids, muzzle, hooves and tail switch are generally black		
	Male	Female	Overall
Height (avg. cm.)	141.81	123.52	
Body Length (avg. cm.)	119.37	110.12	
Heart girth (avg. cm.)	175.64	152.05	
Body weight (avg. kg.)	340.0	235.0	
Birth weight (avg. kg.)	21.0	20.5	
<b>Performance</b>			
	Average	Minimum	Maximum
Litter size born	1.0		
Age at first parturition (months)	46.98	27.0	82.0
Parturition interval (months)	15.59	11.0	16.0
Milk yield per lactation (kg)	1049.24	900.0	1200.0
Milk Fat (%)	4.9	4.1	5.2
Any Peculiarity of the breed			