Breed Descriptor of Gohilwadi (Goat)





Breeding Tract						
State: Gujarat	Porbandar	Porbandar				
	Rajkot					
	Junagadh					
	Bhavnagar					
	Amreli					
Location						
Longitude (deg.)	70.36 - 72.11	70.36 - 72.11				
Latitude (deg.)	21.31 - 21.46	21.31 - 21.46				
General Information						
Synonyms						
Origin	Originated fro	Originated from Gohilwad, a part of Kathiawad region of				
	Gujarat					
Major utility	Food - Meat and Milk					
Comments on utility						
Comments on breeding						
tract						
Adaptability to environmen	t Adapted to ho	Adapted to hot semi-arid climate, and medium and deep				
-	black soil prevalent in South Saurashtra zone of					
	Kathiawar Peninsulal.					
Population						
	Year	Population	Other information			
	2003	230000	Source: Indian J.			
			Animal Sciences,			
			2008, 78(5):567.			
	2013	168917	Source: Estimated			
			Livestock Population			
			Breed Wise Based on			
			Breed Survey 2013.			
			Department of			
			Animal Husbandry,			
			Dairying & Fisheries,			
			Government of India,			
			New Delhi			
Management						
Management system	Semi-Intensive Semi-Intensive					

Mobility	Stationary				
Feeding of adults	Grazing and Fodder				
Comments on Management	Animals are maintained on grazing. Milking does are				
	given feed supplement.				
Morphology					
Colour	Black				
Horn shape and size	Horns are slightly twisted and turned upward, outward				
	and backward. Horns are 15-25cm long.				
Visible characteristics	Nose line is slightly convex. Ears are tubular and				
	drooping. Body covered with coarse long hairs.				
	Male	Female	Overall		
Height (avg. cm.)	88.4	80.4			
Body Length (avg. cm.)	85.9	79.3			
Heart girth (avg. cm.)	83.9	78.8			
Body weight (avg. kg.)	52.04	41.67			
Performance					
	Average	Minimum	Maximum		
Litter size born	1.5				
Age at first parturition	18.0				
(months)					
Parturition interval	9.86				
(months)					
Milk yield per lactation (kg)	240.0				
Fleece weight (kg/year)	3.17				
Any Peculiarity of the breed					