


Breed Descriptor of Miri (Poultry-Fowl)



Breeding Tract			
State: Assam	Dhemaji		
	Lakimpur		
	Sibsagar		
Location			
Longitude (deg.)	93.45 - 95.30		
Latitude (deg.)	26.45 - 28.		
General Information			
Synonyms	Porog		
Origin	Derived its name after the name of tribe "Miri"		
Major utility	Food - Meat and Egg		
Comments on utility			
Comments on breeding tract			
Adaptability to environment			
Population			
	Year	Population	Other information
	2001	1331000	AAU, Khanapara under NATP project on Animal Genetic Resource Biodiversity
Management			
Management system	Backyard		
Mobility	Stationary		
Feeding of adults	Scavenging		
Comments on Management	Birds are kept in cages only during night. Cages are made of cane and bamboo and kept inside their dwelling. Birds are also provided with supplemental feedind		
Morphology			
	Male	Female	Overall
Birth weight (avg. kg.)			29.11
Adult weight(avg. kg)	1.52		
Plumage type	Normal		
Plumage pattern	Solid		
Plumage colour	White and Brown		

Comb type	Single		
Skin colour	White		
Shank colour	White		
Egg shell colour	Light Brown		
Performance			
	Average	Minimum	Maximum
Age at first egg (months)	6.99	6.58	7.57
Annual eggs production (avg. no.)	62.23	54.0	67.0
Eggs weight (avg. gm)	42.06	40.0	44.0
Fertility (%)	89.93	88.0	91.0
Hatching (%)	79.32	73.0	83.0
Dressing (%)	81.46	79.0	83.0
Any Peculiarity of the breed	Naked neck and frizzle birds have also been observed in very few cases.		