

Estrogen flows in front of nauseous/vomiting pregnant women

6

hormonal correlations of nausea/vomiting of pregnancy = $\uparrow$ β -hCG levels (seen w/

$\uparrow$ placental tissue=multifetal gestations, molar pregnancies), $\uparrow$ estradiol,
 $\uparrow$ progesterone (sm.

muscle relaxation $\rightarrow$ decreased sphincter tone $\rightarrow$ $\uparrow$ reflux)

7. Protective dome over flowers in front of nauseous/vomiting pregnant women

$\rightarrow$ hormonal correlations of nausea/vomiting of pregnancy = $\uparrow$ β -hCG levels
(seen w/

$\uparrow$ placental tissue=multifetal gestations, molar pregnancies), $\uparrow$ estradiol,
 $\uparrow$ progesterone (sm.

muscle relaxation $\rightarrow$ decreased sphincter tone $\rightarrow$ $\uparrow$ reflux)

8. Young nauseous woman w/ pigtales

$\rightarrow$ demographics of those most likely to have nausea/vomiting during pregnancy
= young,
first pregnancy, carrying a female fetus

9. "My First Cruise!" T-shirt on nauseous woman

$\rightarrow$ demographics of those most likely to have nausea/vomiting during pregnancy
= young,
first pregnancy, carrying a female fetus

10. "It's a Girl!" balloon next to nauseous woman

$\rightarrow$ demographics of those most likely to have nausea/vomiting during pregnancy
= young,
first pregnancy, carrying a female fetus

11. Nauseous woman next to crossed out hamburger sign

$\rightarrow$ nausea/vomiting of pregnancy chief complaint = n/v, PO intolerance

12. Nauseous woman now vomiting

$\rightarrow$ RF for nausea/vomiting of pregnancy = h/o hyperemesis gravidarum, h/o
migraines, h/o
motion sickness, FHx of hyperemesis gravidarum

13. Tight headband on vomiting woman

$\rightarrow$ RF for nausea/vomiting of pregnancy = h/o hyperemesis gravidarum, h/o
migraines, h/o
motion sickness, FHx of hyperemesis gravidarum

14. Vomiting woman being splashed by a wave

RF for nausea/vomiting of pregnancy = h/o hyperemesis gravidarum, h/o migraines, h/o

motion sickness, FHx of hyperemesis gravidarum

15. Family photo covered in vomit

→ RF for nausea/vomiting of pregnancy = h/o hyperemesis gravidarum, h/o migraines, h/o motion sickness, FHx of hyperemesis gravidarum

16. Man diving down off of a diving board

→ nausea/vomiting of pregnancy vitals to watch out for = orthostatic hypotension, tachycardia, weight loss >5% of prepregnancy weight (qualifies for hyperemesis dx)

17. Diver with raised arm + heart watch

→ nausea/vomiting of pregnancy vitals to watch out for = orthostatic hypotension, tachycardia, weight loss >5% of prepregnancy weight (qualifies for hyperemesis dx)

18. >5% belt buckle

→ nausea/vomiting of pregnancy vitals to watch out for = orthostatic hypotension, tachycardia, weight loss >5% of prepregnancy weight (qualifies for hyperemesis dx)

19. Low pool water volume

→ nausea/vomiting of pregnancy PE findings = dry mucous membranes (2/2 dehydration)

20. Cluster of red beach balls in the low volume pool

→ severe nausea/vomiting of pregnancy labs = CBC → hemoconcentration 2/2 volume depletion, hypochloremic metabolic (contraction) alkalosis, ↓K⁺, ↓Ca²⁺, ↑AST/ALT, ↑amylase/lipase, UA → ketonuria (dx hyperemesis)

21. Sunken chlorine buoy releasing “OH” bubble

→ severe nausea/vomiting of pregnancy labs = CBC → hemoconcentration 2/2 volume depletion, hypochloremic metabolic (contraction) alkalosis, ↓K⁺, ↓Ca²⁺, ↑AST/ALT, ↑amylase/lipase, UA → ketonuria (dx hyperemesis)

22. Bananas falling off of food stand

→ severe nausea/vomiting of pregnancy labs = CBC → hemoconcentration 2/2 volume depletion, hypochloremic metabolic (contraction) alkalosis, ↓K⁺, ↓Ca²⁺, ↑AST/ALT, ↑amylase/lipase, UA → ketonuria (dx hyperemesis)

23. Milk spilling off of food stand
→ severe nausea/vomiting of pregnancy labs = CBC → hemoconcentration 2/2 volume
depletion, hypochloremic metabolic (contraction) alkalosis, ↓K+, ↓Ca²⁺,
↑AST/ALT,
↑amylase/lipase, UA → ketonuria (dx hyperemesis)
24. Unspilled mALT beverage bottles on food stand
→ severe nausea/vomiting of pregnancy labs = CBC → hemoconcentration 2/2 volume
depletion, hypochloremic metabolic (contraction) alkalosis, ↓K+, ↓Ca²⁺,
↑AST/ALT,
↑amylase/lipase, UA → ketonuria (dx hyperemesis)
25. Bitten apples + fatty potato chips hung up high on the food stand
→ severe nausea/vomiting of pregnancy labs = CBC → hemoconcentration 2/2 volume
depletion, hypochloremic metabolic (contraction) alkalosis, ↓K+, ↓Ca²⁺,
↑AST/ALT,
↑amylase/lipase, UA → ketonuria (dx hyperemesis)
26. Candy key snacks in spilled yellow drink
→ severe nausea/vomiting of pregnancy labs = CBC → hemoconcentration 2/2 volume
depletion, hypochloremic metabolic (contraction) alkalosis, ↓K+, ↓Ca²⁺,
↑AST/ALT,
↑amylase/lipase, UA → ketonuria (dx hyperemesis)
27. Twin dolphins with ultrasound sonar
→ severe nausea/vomiting of pregnancy ultrasound w/u → check for multifetal gestation,
molar pregnancy (snow-storm pattern)
28. Snow-storm bubbles from ultrasound dolphins
→ severe nausea/vomiting of pregnancy ultrasound w/u → check for multifetal gestation,
molar pregnancy (snow-storm pattern)
29. DDx sign on S.S. Hyperemesis boat
→ DDx of nausea & vomiting in pregnancy: Pregnancy = pre-eclampsia, acute fatty liver of pregnancy. GI = gastroenteritis, peptic ulcer disease, h.pylori, gastroparesis, pancreatitis, hepatitis, cholelithiasis, cholangitis, appendicitis. GU = kidney stones, pyelonephritis, ovarian torsion. Endo = DKA, Hyperthyroidism, Hyperparathyroidism. Neuro = migraine, pseudotumor cerebri, tumor. Tox = drug toxicity

workup for nausea/vomiting of pregnancy = assess 1. volume status 2. nutritional status

(ketonuria) 3. weight loss (if >5% pre-pregnancy weight 4. PO intolerance

31. Candy key life preserver in spilled yellow fluid

→ workup for nausea/vomiting of pregnancy = assess 1. volume status 2. nutritional status

(ketonuria) 3. weight loss (if >5% pre-pregnancy weight 4. PO intolerance

32. Vomiting woman wearing >5% belt

→ workup for nausea/vomiting of pregnancy = assess 1. volume status 2. nutritional status

(ketonuria) 3. weight loss (if >5% pre-pregnancy weight 4. PO intolerance

33. Pill bottle with baby label

→ conservative tx for nausea/vomiting of pregnancy (for minimal sx) = prevention (prenatal

vitamins x 1mo. before pregnancy), small high protein meals q1-2h, avoid fatty foods,

avoid triggers (odors/noise/heat)

34. Woman relaxing with eye mask and ear plugs

→ conservative tx for nausea/vomiting of pregnancy (for minimal sx) = prevention (prenatal

vitamins x 1mo. before pregnancy), small high protein meals q1-2h, avoid fatty foods,

avoid triggers (odors/noise/heat)

35. 2 plates with chicken breasts

→ conservative tx for nausea/vomiting of pregnancy (for minimal sx) = prevention (prenatal

vitamins x 1mo. before pregnancy), small high protein meals q1-2h, avoid fatty foods,

avoid triggers (odors/noise/heat)

36. Pair of ducks flying by

→ pharmacologic tx for nausea/vomiting of pregnancy (mild → severe sx) = 1. Vitamin B6

(pyridoxine) +/- doxylamine (1st line tx) 2. Antihistamines 3. DA antagonists (metoclopramide, promethazine, prochlorperazine) → SE tardive dyskinesia 4. 5-HT₃

receptor antagonists (ondansetron) → avoid <10wks GA 5. Corticosteroids (methylprednisolone)

37. Woman at spa with a bee swatter

→ pharmacologic tx for nausea/vomiting of pregnancy (mild → severe sx) = 1. Vitamin B6

(pyridoxine) +/- doxylamine (1st line tx) 2. Antihistamines 3. DA antagonists

(metoclopramide, promethazine, prochlorperazine) → SE tardive dyskinesia 4. 5-HT₃

receptor antagonists (ondansetron) → avoid <10wks GA 5. Corticosteroids (methylprednisolone)

38. Lifeboat rope

→ pharmacologic tx for nausea/vomiting of pregnancy (mild → severe sx) = 1.

Vitamin B₆

(pyridoxine) +/- doxylamine (1st line tx) 2. Antihistamines 3. DA antagonists (metoclopramide, promethazine, prochlorperazine) → SE tardive dyskinesia 4. 5-HT₃

receptor antagonists (ondansetron) → avoid <10wks GA 5. Corticosteroids (methylprednisolone)

39. Woman next to rope sticking her tongue out

→ pharmacologic tx for nausea/vomiting of pregnancy (mild → severe sx) = 1.

Vitamin B₆

(pyridoxine) +/- doxylamine (1st line tx) 2. Antihistamines 3. DA antagonists (metoclopramide, promethazine, prochlorperazine) → SE tardive dyskinesia 4. 5-HT₃

receptor antagonists (ondansetron) → avoid <10wks GA 5. Corticosteroids (methylprednisolone)

40. Ribbon dancer swirling "10" with her ribbon

→ pharmacologic tx for nausea/vomiting of pregnancy (mild → severe sx) = 1.

Vitamin B₆

(pyridoxine) +/- doxylamine (1st line tx) 2. Antihistamines 3. DA antagonists (metoclopramide, promethazine, prochlorperazine) → SE tardive dyskinesia 4. 5-HT₃

receptor antagonists (ondansetron) → avoid <10wks GA 5. Corticosteroids (methylprednisolone)

41. Moon ball

→ pharmacologic tx for nausea/vomiting of pregnancy (mild → severe sx) = 1.

Vitamin B₆

(pyridoxine) +/- doxylamine (1st line tx) 2. Antihistamines 3. DA antagonists (metoclopramide, promethazine, prochlorperazine) → SE tardive dyskinesia 4. 5-HT₃

receptor antagonists (ondansetron) → avoid <10wks GA 5. Corticosteroids (methylprednisolone)

42. Ivy waterski rope

→ fluid/nutrition tx for nausea/vomiting of pregnancy = if PO intolerance → IVF, if losing weight → NG tube, if can't tolerate → TPN (high morbidity)

43. Skinny man with straw in his nose

fluid/nutrition tx for nausea/vomiting of pregnancy = if PO intolerance → IVF, if losing

weight → NG tube, if can't tolerate → TPN (high morbidity)

44. TPN lunchbox

→ fluid/nutrition tx for nausea/vomiting of pregnancy = if PO intolerance → IVF, if losing

weight → NG tube, if can't tolerate → TPN (high morbidity)

45. Partially fixed hole on waterslide

→ complications of severe nausea/vomiting of pregnancy (hyperemesis gravidarum) = 1.

Malnutrition → poor wound healing, immunosuppression, Wernicke encephalopathy 2/2

thiamine deficiency (gait ataxia, oculomotor dysfxn, encephalopathy) 2.

AKI/ATN 3.

Mallory-Weiss tears 4. Esophageal rupture 5. Pneumothorax 6. Diaphragmatic tears 7.

Splenic avulsion

46. Drunk guy drinking Wernicke lager

→ complications of severe nausea/vomiting of pregnancy (hyperemesis gravidarum) = 1.

Malnutrition → poor wound healing, immunosuppression, Wernicke encephalopathy 2/2

thiamine deficiency (gait ataxia, oculomotor dysfxn, encephalopathy) 2.

AKI/ATN 3.

Mallory-Weiss tears 4. Esophageal rupture 5. Pneumothorax 6. Diaphragmatic tears 7.

Splenic avulsion

47. Cracked kidney cup

→ complications of severe nausea/vomiting of pregnancy (hyperemesis gravidarum) = 1.

Malnutrition → poor wound healing, immunosuppression, Wernicke encephalopathy 2/2

thiamine deficiency (gait ataxia, oculomotor dysfxn, encephalopathy) 2.

AKI/ATN 3.

Mallory-Weiss tears 4. Esophageal rupture 5. Pneumothorax 6. Diaphragmatic tears 7.

Splenic avulsion

48. Tears along waterslide

→ complications of severe nausea/vomiting of pregnancy (hyperemesis gravidarum) = 1.

Malnutrition → poor wound healing, immunosuppression, Wernicke encephalopathy 2/2

thiamine deficiency (gait ataxia, oculomotor dysfxn, encephalopathy) 2.

AKI/ATN 3.

Mallory-Weiss tears 4. Esophageal rupture 5. Pneumothorax 6. Diaphragmatic tears 7.

Splenic avulsion

49. Ruptured hole on waterslide

→ complications of severe nausea/vomiting of pregnancy (hyperemesis gravidarum) = 1.

Malnutrition → poor wound healing, immunosuppression, Wernicke encephalopathy 2/2

thiamine deficiency (gait ataxia, oculomotor dysfxn, encephalopathy) 2.
AKI/ATN 3.

Mallory-Weiss tears 4. Esophageal rupture 5. Pneumothorax 6. Diaphragmatic tears 7.

Splenic avulsion

50. Little girl holding small doll

→ fetal complications of severe nausea/vomiting of pregnancy (hyperemesis gravidarum) =
low birth weight