
1
8
9
10
7
11
4
12
5
CONTROL PANEL
6
2
A
3
23
REMODELING
14
16
17
15
18
13
B
A
19
J3
20
21
22
Angel of New Orleans
37
41
48
KEEP OUT OF WATER!
25
40
39
38
42
44
43
CONTROL PANEL
45
29
BNP
CARNIVAL
28
32
33
27
30
47
36
49
C
D
24
46
50
HYDRO BOAT
31
NITRO
NITRO
26
34
35

2.2 – Heart Failure P (Management)

1) Floppy heart balloon:
Heart failure

2) American flag:
Based on history, exam findings, and echo results, place patient into the appropriate **ACCF/AHA heart failure stage (1)**

ACCF/AHA **Stage A** HF

3) “A” deck:
ACCF/AHA **Stage A** = At high risk for HF, but no structural heart disease or symptoms **(2)**

4) Red dice:
ACCF/AHA Stage A HF = At **high risk** for HF, but no structural heart disease or symptoms **(2)**

6) Control panel:
Management of Stage A HF = **control HF risk factors** (e.g., HTN, obesity, hyperlipidemia, diabetes, cardiotoxic agents, tobacco) (3)

7) Steam coming out of ears:
Management of Stage A HF = control HF risk factors (e.g., **HTN**, obesity, hyperlipidemia, diabetes, tobacco, cardiotoxic agents) **(3)**

8) Obese man:
Management of Stage A HF = control HF risk factors (e.g., HTN,**obesity**, hyperlipidemia, diabetes, tobacco, cardiotoxic agents) **(3)**

9) Stick of butter:
Management of Stage A HF = control HF risk factors (e.g., HTN, obesity, **hyperlipidemia**, diabetes, tobacco, cardiotoxic agents) **(3)**

10) Diasweeties candy:
Management of Stage A HF = control HF risk factors (e.g., HTN, obesity, hyperlipidemia, **diabetes**, tobacco, cardiotoxic agents) **(3)**

11) Cigarette:
Management of Stage A HF = control HF risk factors (e.g., HTN, obesity, hyperlipidemia, diabetes, **tobacco**, cardiotoxic agents) **(3)**

12) Heart-shaped box of rubies:
Management of Stage A HF = control HF risk factors (e.g., HTN, hyperlipidemia, diabetes, obesity, tobacco, **cardiotoxic agents**) **(3)**

ACCF/AHA **Stage B** HF

13) “B” deck:
ACCF/AHA **Stage B** HF = structural heart disease WITHOUT signs or symptoms of HF **(4)**

14) REMODELING sign:
ACCF/AHA **Stage B** HF = **structural heart disease** WITHOUT signs or symptoms of HF **(4)**

15) Happy guy on B deck:
ACCF/AHA Stage B HF = structural heart disease **WITHOUT signs or symptoms** of HF **(4)**

16) Broken string on heart-shaped guitar:
Structural heart disease includes LV remodeling, **history of MI**, & valvular disease

17) Broken saxophone valve:
Structural heart disease includes LV remodeling, **history of MI**, & valvular disease

18) Falling cake fraction on B deck:
In **HFrEF patients**, ACE inhibitors (e.g., lisinopril, captopril, enalapril) prevent symptomatic HF & reduce mortality **(5)**

19) Ace card:
In HFrEF patients, **ACE inhibitors (e.g., lisinopril, captopril, enalapril)**prevent symptomatic HF & reduce mortality **(5)**

20) Muted beta bugle:
In HFrEF patients, **beta blockers** (e.g., carvedilol, metoprolol, bisoprolol) prevent symptomatic HF **(6)**

21) Carved, metal shield with image of bison:
In HFrEF patients, beta blockers **(e.g., carvedilol, metoprolol, bisoprolol)** prevent symptomatic HF **(6)**

22) Angel playing heart-shaped harp with broken string:
In HFrEF patients with **history of ACS**, ACE inhibitors & beta blockers have been shown to **reduce mortality (7)**

23) Steampunk pirate next to broken heartstring:
Statins should be used in patients with history of ACS to prevent symptomatic HF **(8)**

24) C Deck:
ACCF/AHA **Stage C** HF = structural heart disease + current or prior symptoms of HF **(9)**

ACCF/AHA **Stage C** HF

25) Unhappy bride holding floppy heart balloon:
ACCF/AHA Stage C HF = structural heart disease + **current or prior symptoms of HF (9)**

26) Falling cake fraction on C deck:
Reduced ejection fraction

27) Arrow pointing to staircase to level B:
ACE inhibitors / ARBs (if unable to tolerate ACEi) & beta blockers should be used in all patients with Stage C HFrEF to decrease morbidity & mortality **(10)**

28) cARNIval banner:
ARNIs (e.g., Valsartan/Sacubitril) should replace ACEi / ARB in patients with Stage C HFrEF to reduce morbidity & mortality **(11)**

29) BNP blimp:
Sacubitril inhibits neprilysin, an enzyme that degrades natriuretic peptides (e.g., **BNP**)

30) Spiral notebook with picture of apple:
Aldosterone receptor antagonists (e.g., **spironolactone, eplerenone**) improve morbidity and mortality in patients with Stage C HFrEF & symptoms limiting activity (i.e., NYHA class II-IV) (12)

31) Hydro Boat loaded with Nitro:
Hydralazine + Isosorbide Dinitrate reduce morbidity & mortality in some African-American patients with Stage C HFrEF (i.e., significant symptoms despite optimal therapy) **(13)**

32) African-American bridesmaid:
Hydralazine + Isosorbide Dinitrate reduce morbidity & mortality in some **African-American patients**with Stage C HFrEF (i.e., significant symptoms despite optimal therapy) (13)

33) Tired angel bridesmaid:
Hydralazine + Isosorbide Dinitrate reduce morbidity & mortality in some African-American patients with Stage C HFrEF (i.e., **significant symptoms despite optimal therapy**) **(13)**

34) Metal heart pin:
Implanted cardiac devices (i.e., ICD, pacemaker) can improve mortality in some patients with Stage C HFrEF (e.g., NYHA II-III and LVEF ≤ 35% on optimal medical therapy) **(14)**

35) Maid of honor suddenly falling into water:
ICDs are recommended for prevention of **sudden cardiac death** in certain patients with Stage C HFrEF & symptoms on optimal medical therapy

36) Wet, angry bride:
Loop diuretics (e.g., furosemide, bumetanide) can **improve symptoms in patients with Stage C HF and evidence of fluid retention (e.g., crackles, elevated JVP, peripheral edema)**, but offer NO mortality benefit **(15)**

---

37) Looped tiara:
Loop diuretics (e.g., furosemide, bumetanide) can improve symptoms in patients with Stage C HF and evidence of fluid retention (e.g., crackles, elevated JVP, peripheral edema), but offer NO mortality benefit **(15)**

38) Pantsless waiter with white chloro-thighs:
Thiazides (e.g., HCTZ, chlorthalidone) are good diuretic options for HF patients with mild fluid retention AND concurrent HTN due to their antihypertensive effects **(16)**

39) Wet waiter:
Thiazides (e.g., HCTZ, chlorthalidone) are good diuretic options for HF patients with **mild fluid retention** AND concurrent HTN due to their antihypertensive effects **(16)**

40) Angry waiter:
Thiazides (e.g., HCTZ, chlorthalidone) are good diuretic options for HF patients with mild fluid retention AND **concurrent HTN** due to their antihypertensive effects **(16)**

41) Foxglove flower bouquet:
Digoxin can decrease hospitalizations in patients with HFrEF, but offers NO mortality benefit **(17)**

42) Dropped salt shaker:
Sodium restriction is reasonable for patients with symptomatic HF to reduce congestive symptoms **(18)**

---

43) Preserved cake fraction:
Management goals for Stage C **HFpEF** = treatment of symptoms & control of underlying conditions

44) Patched steaming pipe:
Blood pressure should be controlled (i.e., ACE inhibitors, ARBs, and/or beta blockers) in patients with Stage C HFpEF to prevent morbidity **(19)**

45) Control panel with Afib on monitor:
AFib should be controlled in patients with Stage C HFpEF to prevent morbidity **(20)**

ACCF/AHA **Stage D** HF

46) “D” buoy floating in water:
Stage D HF = refractory HF requiring specialized interventions **(21)**

47) Soaking wet groom:
Stage D HF = **refractory HF** requiring specialized interventions **(21)**

48) “Keep Out of Water!” sign:
Fluid restriction(~2 L/day) is reasonable for patients in Stage D HF to improve congestive symptoms **(22)**

49) LVAD fanny pack:
LVAD can improve mortality in carefully selected patients with Stage D HF **(23)**

50) Mermaid offering heart locket:
Cardiac transplantation is indicated for carefully selected patients with Stage D HF **(24)**