



## Unit 2 3set1 MTH174 MCQs for practice

Engineering Mathematics (Lovely Professional University)

**Normal ODE, Dependent –Independent functions , Homogeneous Linear ODE:**

- 1.) The differential equation  $y'' + 3y' + \sqrt{x}y = \sin x$  is normal on every subinterval of  
a)  $(-\infty, \infty)$                       b)  $[0, \infty)$                       c)  $(0, \infty)$                       d)  $(-\infty, 1), (-1, 1), (1, \infty)$
- 2.) The differential equation  $(x^2 - 1)y'' + 2xy' + y = x \ln x$  is normal on every subinterval of  
a)  $(-\infty, \infty)$                       b)  $(0, \infty)$                       c)  $(-\infty, 2)$                       d)  $(0, 1), (1, \infty)$
- 3.) Which of the following functions are linearly independent for  $x \in (0, \infty)$  ?  
a)  $1, x, x^2, 1+x$                       b)  $1, x(1-x), x^2, x$                       c)  $2x, 6x+3, 3x+2$                       d)  $1, x, x^2, x^2(1-x)$
- 4.) If  $y_1(x)$  and  $y_2(x)$  be the linearly independent solutions of the equation  $y'' + a(x)y' + b(x)y = 0$  on an interval  $I$  then which of the following is true  
a) both  $y_1(x)$  and  $y_2(x)$  vanishes for some  $x_0 \in I$   
b) both  $y_1(x)$  and  $y_2(x)$  take extreme values for some  $x_0 \in I$   
c) both  $y_1(x)$  and  $y_2(x)$  can not vanishes for some  $x_0 \in I$                       d) None of these
- 5.) The general solution of the differential equation  $y'' + 2\pi y' + \pi^2 y = 0$  is  
a)  $(A + Bx)e^{-\pi x}$                       b)  $(A + Bx)e^{-x}$                       c)  $(A + B)e^{-\pi x}$                       d)  $(A + Bx)e^{\pi x}$
- 6.) The differential equation whose linearly independent solutions are  $e^{2x}, xe^{2x}$  is ?  
a)  $y'' + 4y' + 4y = 0$                       b)  $y'' + 4y' - 4y = 0$                       c)  $y'' - 4y' + 4y = 0$                       d)  $y'' - 4y' - 4y = 0$
- 7.) The lowest possible order of homogeneous linear differential equation whose particular solution is  $3\cos 2x + 5\sinh 3x$  is  
a) 2                      b) 3                      c) 5                      d) 4
- 8.) The roots of characteristic equation of differential equation  $y^{iv} + 8y'' - 9y = 0$   
a)  $\pm 1, \pm 3i$                       b)  $\pm i, \pm 3i$                       c)  $\pm i, \pm 3$                       d)  $\pm 1, \pm 3$
- 9.) The general solution of differential equation  $y''' - 2y'' + y' = 0$  is  
a)  $Ae^x + (Bx + C)$                       b)  $A + (Bx + C)e^x$                       c)  $(Bx + C)e^x$                       d)  $A + (Bx + C)e^{-x}$
- 10.) The lowest possible order of homogeneous linear differential equation whose particular solution is  $1 + x + e^x - 3e^{3x}$  is  
a) 4                      b) 3                      c) 2                      d) 5

**(Operator method ) Exp(ax), cosh(ax) sinh(ax), h(x). exp(ax): : Polynomila , Sin(ax) , cos(ax):**

1. The particular integral  $\frac{1}{D+3} e^{2x}$  is  
(a)  $\frac{1}{5} e^{2x}$  (b)  $\frac{1}{5}$  (c)  $\frac{1}{3} e^{2x}$  (d)  $\frac{1}{2x+3} e^{2x}$
2. The particular integral  $\frac{1}{D^2-9} e^{3x}$  is  
(a)  $\frac{1}{6} e^{3x}$  (b)  $\frac{x e^{3x}}{6}$  (c)  $\frac{x}{3} e^{3x}$  (d) doesn't exist
3. The particular integral  $\frac{1}{f(D)} e^{ax} g(x)$  is  
(a)  $e^{ax} \frac{1}{f(D)} g(x)$  (b)  $g(x) \frac{1}{f(D)} e^{ax}$  (c)  $e^{ax} \frac{1}{f(D+a)} g(x)$  (d)  $\frac{1}{f(a)} e^{ax} g(x)$
4. The particular integral  $\frac{1}{f(D^2)} e^{ax}$  is  
(a)  $\frac{1}{f(-a^2)} e^{ax}$  (c)  $\frac{1}{f(a^2)} e^{ax}$ , provided  $f(a^2) = 0$   
(b)  $\frac{1}{f(a^2)} e^{ax}$ , provided  $f(a^2) \neq 0$  (d)  $\frac{1}{f(a)} e^{ax}$ , provided  $f(a) \neq 0$
5. The particular integral  $\frac{1}{D^3-D^2+4D-4} \sin 3x$  is  
(a)  $-\frac{1}{5} \sin 3x$  (c)  $\frac{1}{9} x \cos 3x$   
(b)  $\frac{1}{50} (\sin 3x + x \cos 3x)$  (d)  $\frac{1}{50} (\sin 3x + 3 \cos 3x)$
6. The particular integral of the differential equation  $y'' + y = 6 \sin x$  is  
(a)  $6 \cos x$  (b)  $3x \sin x$  (c)  $-3x \cos x$  (d)  $6 x \cos x$
7. The particular integral  $\frac{1}{D+5} (2016)^x$  is  
(a)  $\frac{1}{2021} (2016)^x$  (b)  $x (2016)^x$  (c)  $\frac{1}{\ln 2016} (2016)^x$  (d)  $\frac{1}{(\ln 2016)+5} (2016)^x$
8. The particular integral of  $\frac{d^2y}{dx^2} + \frac{dy}{dx} = x^2 + 2x + 4$  is  
(a)  $\frac{x^2}{3} + 4x$  (b)  $\frac{x^3}{3} + 4x$  (c)  $\frac{x^3}{3} + 4$  (d)  $\frac{x^2}{3} + 4x^2$
9. The particular integral of  $\frac{d^4y}{dx^4} - 16 \frac{d^2y}{dx^2} = (8x + 16)$  is  
(a)  $-\frac{x^2}{12} - \frac{x}{2}$  (b)  $\frac{x^3}{6} + 2x$  (c)  $\frac{x^3}{3} + 4$  (d)  $\frac{x^2}{3} + 4x$
10. The particular integral  $\frac{1}{(D+1)^3} (2x + 4)$  is  
(a)  $\frac{x^2}{3} + 4x$  (b)  $2x + 4$  (c)  $x - 2$  (d)  $2x - 2$
11. The particular integral of the differential equation  $y'' + 4y = \sin x \cos x$  is  
(a)  $6 \cos x$  (b)  $3x \sin x \cos x$  (c)  $-3x \cos x$  (d)  $-\frac{x}{8} \cos 2x$
12. The particular integral  $\frac{1}{D^2} \cos 2x$  is  
(a)  $-4 \cos 2x$  (b)  $-4 \sin 2x$  (c)  $\frac{1}{2} \sin 2x$  (d)  $-\frac{1}{4} \cos 2x$
13. The particular integral  $\frac{1}{D-1} \cos 2x$  is  
(a)  $-\frac{1}{5} (\cos 2x - 2 \sin 2x)$  (b)  $\frac{1}{2} \cos 2x$  (c)  $-\sin 2x$  (d)  $-\frac{1}{5} (\sin 2x + 2 \cos 2x)$

### Method of Variation of Parameter, Method of Undetermined co-efficient:

1 The value of parameter  $A(x)$  for LDE  $y'' - 2y' - 3y = e^x$  using method of variation of parameters when  $y_1 = e^{-x}$  and  $y_2 = e^{3x}$  is

- (a)  $-\frac{e^{2x}}{8} + c$  (b)  $\frac{3}{2}e^x + c$  (c)  $\frac{2}{3}e^{3x} + c$  (d)  $-\frac{e^{5x}}{18} + c$

2 Solving by variation of parameter for the equation  $y'' + y = \sec x$ , the value of Wronskian

- a. 1 b. 2 c. 3 d. 4

3 Solving by variation of parameters for the equation  $y'' - 4y' + 3y = e^x$ ,  $x \neq 0$  the value of Wronskian is

- (a)  $2e^x$  (b)  $3e^{4x}$  (c)  $2e^{4x}$  (d)  $4e^x$

4 If by the method of variation of parameter  $y(x) = A(x)\sin x + B(x)\cos x$  is the general solution of  $y'' + y = \sec x$  then  $B(x)$  is

- (A)  $\ln|\sin x| + c$ , (B)  $\ln|\cos x| + c$ , (C)  $X + c$ , (D)  $\ln|x| + c$

5 The choice of particular integral for the equation  $y'' - 9y = 13e^{3x}$  is

- (a)  $ce^{3x}$  (b)  $cxe^{3x}$  (c)  $cx^2e^{3x}$  (d) none of these

6 By the method of undetermined coefficients the choice of particular integral of the ODE

$$y'' + 4y' + 4y = 12e^{-2x}$$

- (A)  $ax^2e^{2x}$ , (B)  $ax^2e^{-2x}$ , (C)  $12ae^{2x}$ , (D)  $12axe^{-2x}$

7 By the method of undetermined coefficients the trial solution for  $y_p$  for the differential equation  $y'' + 3y' + 2y = 12x^2$  is of the form

- (A)  $a + bx + cx^2$  (B)  $a + bx$  (C)  $ax + bx^2 + cx^3$  (D) None of these

8 By the method of undetermined coefficients the trial solution for  $y_p$  for the differential equation  $y'' + 2y' + y = 6e^{-x}$  is of the form

- (a)  $Ae^{-x}$  (b)  $Bxe^{-x}$  (c)  $Cx^2e^{-x}$  (d) None of these

Q19. By the method of variation of parameter if  $A(x)\cos x + B(x)\sin x$  be the particular integral of the differential equation

$$y'' + y = \sec x \text{ then } A(x) \text{ is}$$

- (a)  $-\log|\cos x|$  (b)  $\log|\cos x|$  (c)  $\log|\sin x|$  (d)  $-\log|\sin x|$

Q20. By the method of variation of parameter if  $A(x)\cos 2x + B(x)\sin 2x$  be the particular integral of the differential equation

$$y'' + 4y = \sec 2x \text{ then } B(x) \text{ is}$$

- (a)  $3x/2$  (b)  $x^2/2$  (c)  $-x/2$  (d)  $x/2$

Q21. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' + 9y = \cos 3x \text{ is}$$

- (a)  $a\cos 3x + b\sin 3x$  (b)  $x(a\cos 3x + b\sin 3x)$  (c)  $x^2(A\cos 3x + b\sin 3x)$  (d) None of these

Q22. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' + y = \sin x \text{ is}$$

- (a)  $a\cos x + b\sin x$  (b)  $x(a\cos x)$  (c)  $x^2(A\cos 3x + b\sin 3x)$  (d)  $x(a\cos x + b\sin x)$

Q23. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' - 2y' - 3y = 6e^{-x} - 8e^x \text{ is}$$

- (a)  $a e^{-x} + b e^x$  (b)  $a x e^{-x} + b e^x$  (c)  $a e^{-x} + b x e^x$  (d)  $a x^2 e^{-x} + b e^x$

**Q25.** By method of undetermined coefficients the assumed particular integral of the differential equation

- $y'' - y = \sin x$  is  
 (a)  $a \cos x + b \sin x$  (b)  $x(a \sin x)$  (c)  $x^2(A \cos x + b \sin x)$  (d)  $x(a \cos x + b \sin x)$

**Q26.** By method of undetermined coefficients the assumed particular integral of the differential equation

- $y'' - 4y' + 13y = 12e^{2x} \sin 3x$   
 (a)  $a \cos 3x + b \sin 3x$  (b)  $x e^{2x}(a \sin 3x + b \cos 3x)$   
 (c)  $x^2(A \cos 3x + b \sin 3x)$  (d)  $x^2 e^{2x}(a \cos x + b \sin x)$

### Euler-Cauchy Equation , Simultaneous ODE

**Q28.** If  $D = \frac{d}{dx}$ , then  $\frac{1}{x^2 D^2 + 2} 16x^3$  is equals to

- a)  $\frac{1}{2}x^3$  b)  $2x^3$  c)  $\frac{1}{4}(\log x)^3$  d)  $\frac{1}{4}x^3$

**Q29.** C.F of  $(x^2 D^2 - xD)y = 0$  is

- a)  $a + bx$  b)  $ax + bx^2$  c)  $a \log x + bx^2$  d)  $a + bx^2$

**Q30.** To convert the Euler Cauchy equation into a linear equation with constant coefficient we assume

- a)  $x = e^t$  b)  $x = t$  c)  $x = t^2$  d) None of these

**Q31.** If the system of equations is  $(2D - 4)y_1 + (3D + 5)y_2 = 3t + 2$ ,  $(D - 2)y_1 + (D + 1)y_2 = t$  then  $y_2$  is

- a)  $ae^{-3t} + \frac{1}{9}(3t + 5)$  b)  $ae^{-3t}$  c)  $ae^{-2t} + \frac{1}{9}(3t + 5)$  d) None of these

**Q27.** Which of the following is Euler-Cauchy equation?

- a)  $x^3 y'' + x^2 y' + y = 0$  b)  $x^2 y'' + xy' + y = 0$  c)  $x^2 y'' + xy = 0$  d) None of these

**Q40.** Particular integral of  $x^2 y'' - 2y = 2x + 6$  is

- a)  $-x - 3$  b)  $x - 3$  c)  $-x + 3$  d) None of these

**Q38.** The complimentary function of the differential equation  $x^2 y'' - xy' + y = \log x$  is given by

- a)  $ae^x + be^{2x}$  b)  $ax$  c)  $(a + b \log x)x$  d)  $ax \log x$

**Q36.** For the given system of linear differential equation  $y_1' = 2y_1 + y_2$ ,  $y_2' = y_1 + 2y_2$ , then the second order linear differential equation satisfied by  $y_1$  is

- a)  $y_1'' + 4y_1' + 3y_1 = 0$  b)  $y_1'' - 4y_1' + 3y_1 = 0$  c)  $y_1'' - 4y_1' - 3y_1 = 0$  d) None of these

❖ **Q.5** For the differential equation  $(x^3 D^3 - 3xD + 3)y = 0$  using transformation  $x = e^t$  the roots of its operator notation are:

- a) 1, 1, 3 b) 1, -1, 3 c) 1, 1, -3 d) 1, 2, 3

❖ **Q.25** If  $y_1(t), y_2(t)$  satisfy the equations  $y_1' + 5y_2 = 0$ ,  $y_2' + y_1 = 0$  and  $y_2'' + by_2 = 0$  is the second order differential equation satisfied by  $y_1$  then what is the value of b

- a) 5 b) -5 c) 3 d) -3

❖ **Q.29** If  $D = \frac{d}{dx}$ , then  $\frac{1}{(x^2 D^2 + 2)} 16x^3$  is equal to

- a)  $\frac{1}{2}x^3$  b)  $2x^3$  c)  $\frac{1}{4}(\log x)^3$  d)  $\frac{1}{4}x^3$

❖ For a given system of linear differential equation  $y_1' = 2y_1 + y_2$ ,  $y_2' = y_1 + 2y_2$ , the second order linear differential satisfied by the  $y_1$  is

- (A)  $y_1'' + 4y_1' + 3y_1 = 0$  (B)  $y_1'' - 4y_1' + 3y_1 = 0$  (C)  $y_1'' - 4y_1' - 3y_1 = 0$  (D) none

- ❖ The particular integral of differential equation ( $x > 0$ )  
 $x^3 y''' + 5x^2 y'' + 5xy' + y = x^2$  Using the transformation  $x = e^t$ , we get (in operator notation)  $[\theta^3 + 2\theta^2 + 2\theta + 1]y = e^{2t}$  is
- (A)  $\frac{1}{21}e^{2t}$  (B)  $\frac{1}{31}e^{-2t}$  (C)  $-\frac{1}{51}e^{2t}$  (D)  $\frac{1}{21}e^{7t}$