

JEE MAIN

COURSE
NUCLEUS

TEST CODE				
1	1	3	0	4

MOCK TEST-14

Class: XII

Date: 24-12-2018

Time: 3 Hours.

Max. Marks: 360

IMPORTANT INSTRUCTIONS

1. The question paper consists of '90' objective type questions. There are '30' questions each in Physics , Chemistry and Mathematics respectively. Please fill the OMR answer Sheet accordingly and carefully.
2. Each question has four choices (1), (2), (3) and (4) out of which **ONLY ONE** is correct.
3. You will be **awarded 4 marks** for each question, if you have darkened only the bubble corresponding to the correct answer and zero mark if no bubble are darkened. In all other cases, **minus one (-1) mark** will be awarded.
4. There is only one correct response for each question. Filling up more than one response in each question will be treated as wrong response and marks for wrong response will be deducted accordingly as per instruction 3 above.
5. Use **Black or Blue Ball Point Pen** only for filling particulars.
6. Use of **Calculator, Log Table, Slide Rule and Mobile** is not allowed.
7. Rough work is to be done on the space provided at the bottom and in end of the booklet for this purpose in the Test Booklet only.
8. On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator. However, the candidates are allowed to take away this Test Booklet with them.
9. Do not fold or make any stray marks on the Answer Sheet.


BANSAL CLASSES
 PRIVATE LIMITED
Ideal for Scholars

Corporate Office : A-10, "BANSAL TOWER", Road No.-1, I.P.I.A., Kota-324005 (Raj.) INDIA

Tel.: (0744) 2791000

Helpline: 09571042038 | Email: dlpd@bansal.ac.in, dlpd.care@bansal.ac.in | website : www.bansal.ac.in

USEFUL DATA


Atomic weights: Al = 27, Mg = 24, Cu = 63.5, Mn = 55, Cl = 35.5, O = 16, H = 1, P = 31, Ag = 108, N = 14, Li = 7, I = 127, Cr = 52, K = 39, S = 32, Na = 23, C = 12, Br = 80, Fe = 56, Ca = 40, Zn = 65.5, Ti = 48, Ba = 137, U = 238, Co = 59, B = 11, F = 19, He = 4, Ne = 20, Ar = 40, Mo = 96

[Take : $\ln 2 = 0.693$, $\ln 1.1 = 0.095$, $\ln 3 = 1.09$, $e = 1.6 \times 10^{-19}$, $m_e = 9.1 \times 10^{-31}$ kg]

Take: $\epsilon_0 = 8.85 \times 10^{-12}$ C²/Nm², $g = 10$ m/s², $S_{\text{water}} = 1$ cal/gm °C, $L_{\text{ice}} = 80$ cal/gm., $g = 10$ m/s² unless otherwise stated

PHYSICS

- Q.1 Two particles A and B start moving from rest from $x = 0$ at $t = 0$ along x-axis with variable acceleration as shown in graph.



Then relative velocity of A with respect to B is 21 m/sec along negative x-axis after


- (1) 11 sec (2) 21 sec (3) 31 sec (4) None
- Q.2 Wedge is fixed on horizontal surface block A is pulled upwards by applying a force F as shown in the figure and there is no friction between wedge and the block A while coefficient of friction between A and B is μ . If there is no relative motion between block A and block B then frictional force developed between A and B is.

(1) $\left[\frac{F + (M + m)g \sin \theta}{m + m} \right]$

(2) μmg


(3) $\frac{\mu mg}{2}$

(4) $\left[\frac{F - (M + m)g \sin \theta}{m + M} \right] m \cdot \cos \theta$




SPACE FOR ROUGH WORK

- Q.3 A 20 kg block attached to a spring of force constant $k = 5 \text{ N/m}$ is released from rest at A. The spring at this instant is having an elongation of 1m. The block is allowed to move in smooth horizontal slot with the help of a constant force of 50N as shown. The speed of block as it reaches B is :- [Assume the rope to be light].



- (1) 4 m/s (2) 2 m/s (3) 1 m/s (4) 3 m/s
- Q.4 A flexible chain of mass m and length L is suspended vertically as shown in figure. If upper end of the chain is released so that the chain starts falling and forms a small heap on the table. When h length of chain is still in air, what is the force exerted on the table by the chain ?


- (1) $\frac{3mgh}{L}$
- (2) $3mg\left(1 - \frac{h}{L}\right)$
- (3) $mg\left(1 - \frac{h}{L}\right)$
- (4) $\frac{mgh}{L}$



SPACE FOR ROUGH WORK

- Q.5 Two identical rods are joined as shown. The system is pivoted at point O and is released from rest from the horizontal position. The speed of point A when OB becomes vertical is

- (1) $6\sqrt{\frac{gl}{17}}$
 (2) $3\sqrt{\frac{5gl}{17}}$
 (3) $4\sqrt{\frac{3gl}{17}}$
 (4) $2\sqrt{\frac{15gl}{17}}$




- Q.6 At what speed, the velocity head of water is equal to pressure head of 40 cm of Hg ?
 (1) 10.3 m/s (2) 2.8 m/s (3) 5.6 m/s (4) 8.4 m/s

- Q.7 Three very large plates of same area are kept parallel and close to each other. They are considered as ideal black surfaces and have very high thermal conductivity. The first and third plates are maintained at temperatures $2T$ and $3T$ respectively. The temperature of the middle (i.e., second) plate under steady state condition is :

- (1) $\left(\frac{65}{2}\right)^{\frac{1}{4}} T$ (2) $\left(\frac{97}{4}\right)^{\frac{1}{4}} T$ (3) $\left(\frac{97}{2}\right)^{\frac{1}{4}} T$ (4) $(97)^{\frac{1}{4}} T$

- Q.8 The electric dipole is situated along an electric field. The dipole is rotated about an axis perpendicular to the plane of the paper in clockwise direction with respect to the field. The graph between torque and θ will be represented by :




- (1) A (2) B (3) C (4) D

SPACE FOR ROUGH WORK

- Q.9 A bar magnet has a magnetic moment equal to 5×10^{-5} weber-m. It is suspended in a magnetic field, which has a magnetic induction B equal to $8\pi \times 10^{-4}$ tesla. The magnet vibrates with a period of vibration equal to 15 sec. The moment of inertia of the magnet is :
- (1) $22.5 \times 10^{-7} \text{ kg-m}^2$ (2) $11.25 \times 10^{-7} \text{ kg-m}^2$
 (3) $5.62 \times 10^{-7} \text{ kg-m}^2$ (4) $7.16 \times 10^{-7} \text{ kg-m}^2$


- Q.10 Given is a circuit diagram of an AM demodulator. For good demodulation of AM signal of carrier frequency f , the value of RC should be :



- (1) $RC = 1/f$ (2) $RC < 1/f$ (3) $RC \geq 1/f$ (4) $RC \gg 1/f$
- Q.11 Two moles of helium, four moles of hydrogen and one mole of water vapour forms an ideal gas mixture. The molar specific heat at constant pressure of the mixture is (neglecting vibrational degrees of freedom):
- (1) $\frac{7}{3}R$ (2) $\frac{16}{7}R$ (3) $\frac{10}{3}R$ (4) $\frac{23}{7}R$

- Q.12 A small block of mass m is rigidly attached to P to a ring of mass $3m$ and radius R . The system is released from rest at $\theta = 90^\circ$ and rolls without sliding. The angular acceleration of hoop after release is :

- (1) $\frac{g}{4R}$
 (2) $\frac{g}{8R}$
 (3) $\frac{g}{3R}$
 (4) $\frac{g}{2R}$



SPACE FOR ROUGH WORK

Q.13 Two vibrating tuning forks producing progressive waves given by :

$$y_1 = 4 \sin (500\pi t)$$

$$y_2 = 2 \sin (506\pi t)$$

are held near the ear of a person. The person will hear :

- (1) 3 beats/s with intensity ratio between maxima and minima equal to 2
- (2) 3 beats/s with intensity ratio between maxima and minima equal to 9
- (3) 6 beats/s with intensity ratio between maxima and minima equal to 2
- (4) 6 beats/s with intensity ratio between maxima and minima equal to 9

Q.14 A particle free to move along the x-axis has potential energy given by :

$$U(x) = k [1 - \exp(-x^2)] \text{ for } -\infty \leq x \leq +\infty$$

Where k is positive constant of appropriate dimensions. Then :

- (1) at points away from the origin, the particle is in stable equilibrium
- (2) for any finite non-zero value of x, there is a force directed away from the origin
- (3) if its total mechanical energy is k/2, it has its minimum kinetic energy at the origin
- (4) for small displacements from x = 0, the motion is simple harmonic

Q.15 A planet revolves around the sun in an elliptical orbit. If v_p and v_a are the velocities of the planet at the perigee and apogee respectively, then the eccentricity of elliptical orbit is given by :


$$(1) \frac{v_p}{v_a}$$

$$(2) \frac{v_a - v_p}{v_a + v_p}$$

$$(3) \frac{v_p + v_a}{v_p - v_a}$$

$$(4) \frac{v_p - v_a}{v_p + v_a}$$

Q.16 A small hole is made at the bottom of a symmetrical jar as shown in figure. A liquid is filled into the jar upto a certain height. The rate of fall of liquid is independent of the level of liquid in the jar. Then the surface of jar is a surface of revolution of the curve $y = kx^n$, the value of n is :



(1) 1

(2) 2

(3) 3

(4) 4

Q.17 An ideal gas is expanding such that $PT^2 = \text{constant}$. The coefficient of volume expansion of the gas is A/T where T is temperature in kelvin. Find the value of A.

(1) 1

(2) 2

(3) 3


(4) 4

SPACE FOR ROUGH WORK

- Q.18 The intensity absorbed in a material of depth d with absorption coefficient μ , when I_0 is the incident intensity of X-rays, is :
(1) $(1 - e^{-\mu d})$ (2) $I_0 (1 - e^{-\mu d})$ (3) $\mu I_0 (1 - e^{-\mu d})$ (4) μd
- Q.19 Boron has two isotopes $^{10}_5\text{B}$ and $^{11}_5\text{B}$. If the atomic weight of boron is 10.81, the ratio of $^{10}_5\text{B}$ and $^{11}_5\text{B}$ in nature is :
(1) 19/81 (2) 20/53 (3) 15/16 (4) 10/11
- Q.20 A parallel plate capacitor consists of two circular plates each of radius 2 cm, separated by a distance of 0.1 mm. If voltage across the plates is varying at the rate of 5×10^{13} V/s, then the value of displacement current is :
(1) 5.50 A (2) 5.56×10^2 A (3) 5.56×10^3 A (4) 2.28×10^4 A
- Q.21 A beam of light is incident on a glass plate at an angle of incidence 60° . The reflected ray is completely polarised. What is the angle of refraction when angle of incidence is 45° ?
(1) $\sin^{-1}(\sqrt{3}/2)$ (2) $\cos^{-1}(\sqrt{3}/2)$ (3) $\sin^{-1}(\sqrt{1/6})$ (4) $\sin^{-1}(\sqrt{1/3})$
- Q.22 In Young's experiment, find the distance between two slits that results in the third minimum for 420 nm violet light at an angle 30° . It may be assumed that separation between the slits (d) is much smaller than the separation between slits and the screen (D).
(1) 2.5×10^{-6} m (2) 2.1×10^{-6} m (3) 4.2×10^{-6} m (4) None of these


SPACE FOR ROUGH WORK

- Q.23 In the circuit shown in figure R_3 is a variable resistance. As the value R_3 is changed, current I through the cell varies as shown. Obviously, the variation is asymptotic, i.e., $I \rightarrow 6\text{A}$ as $R_3 \rightarrow \infty$. Resistance R_1 and R_2 are, respectively :



- (1) $4\ \Omega, 2\ \Omega$ (2) $2\ \Omega, 4\ \Omega$ (3) $2\ \Omega, 2\ \Omega$ (4) $1\ \Omega, 4\ \Omega$


- Q.24 Equivalent capacitance of network in figure between x and y is (all capacitances are in μF) :



- (1) $8\ \mu\text{F}$ (2) $6\ \mu\text{F}$ (3) $4\ \mu\text{F}$ (4) $2\ \mu\text{F}$


SPACE FOR ROUGH WORK

- Q.25 Three long wires (1, 2 and 3) of resistances in the ratio 3 : 4 : 5 are connected in parallel to each other as shown in the diagram. If net force on middle wire is zero. Then d_1/d_2 will be :



- (1) 9 : 25 (2) 5 : 3 (3) $\sqrt{5} : \sqrt{3}$ (4) 1 : 1

- Q.26 In the CE amplifier circuit shown in the figure an n-p-n transistor of $\beta = 100$ is used. The output voltage of the amplifier is :



- (1) 1 volt (2) 2 volt (3) 3 volt (4) 4 volt


- Q.27 There is horizontal cylindrical uniform but time varying magnetic field increasing at a constant rate $\frac{dB}{dt}$ as shown. A charged particle having charge q and mass m is kept in equilibrium, at the top of a spring of spring constant K in such a way that it is on the horizontal line passing through the centre of the magnetic field as shown in figure. The compression in the spring will be :

(1) $\frac{1}{K} \left[mg - \frac{qR^2}{2l} \frac{dB}{dt} \right]$

(2) $\frac{1}{K} \left[mg + \frac{qR^2}{2l} \frac{dB}{dt} \right]$


(3) $\frac{1}{K} \left[mg + \frac{2qR^2}{l} \frac{dB}{dt} \right]$

(4) $\frac{1}{K} \left[mg + \frac{qR^2}{l} \frac{dB}{dt} \right]$



SPACE FOR ROUGH WORK

Q.28 In the circuit shown in the below figure, current through source will be : [Given $\cos^{-1}(0.6) = 53^\circ$]



- (1) $\frac{1}{3} + \frac{\sqrt{2}}{5} \sin(100\pi t - 8^\circ)$ (2) $\frac{1}{5} + \frac{\sqrt{2}}{5} \sin(100\pi t - 8^\circ)$
 (3) $\frac{1}{3} + \frac{\sqrt{2}}{5} \sin(100\pi t - 98^\circ)$ (4) $\frac{1}{5} + \frac{\sqrt{2}}{5} \sin(100\pi t - 98^\circ)$

Q.29 A compound microscope is used to enlarge an object kept at a distance of 3 cm from its objective. The objective consists of several convex lenses in contact and has a focal length of 2cm. If a lens of focal length 10 cm is removed from the objective, the eyepiece has to be moved by x cm to refocus the image. The value of x is :

- (1) 5 (2) 9 (3) 4 (4) 15

Q.30 Charge is distributed non-uniformly in the volume of a solid insulating sphere of radius 'a' such that volume charge density (ρ) varies with distance from the centre (r) as,

$$\rho(r) = Cr^2, r \leq a$$

Here C is a constant.

Magnitude of electric field strength outside the sphere, i.e., for $r > a$, is :

- (1) $E = \frac{Ca^5}{5\epsilon_0 r^2}$ (2) $\frac{Cr^3}{5\epsilon_0}$ (3) $\frac{Cr^2}{2\epsilon_0}$ (4) $\frac{Ca^3}{\epsilon_0 r^2}$





SPACE FOR ROUGH WORK

CHEMISTRY

Q.31 Which of the following set of transition metal of 3d-series have maximum and minimum melting point respectively?

- (1) Cr and Mn (2) Fe and Zn (3) Cr and Cu (4) Fe and Hg


Q.32 Which of the following reaction will not give propane ?



Q.33 At the given condition of CH_4-O_2 fuel cell the cell emf is 0.8 V and the enthalpy of combustion of $\text{CH}_4(\text{g})$ is -772 kJ/mol . The maximum efficiency of the given fuel cell in the given condition is :


- (1) 60 % (2) 75 % (3) 80 % (4) 90 %

Q.34 For the following conversion 'x' and 'y' could be respectively.







- (1) $\text{H}_2\text{O}_2/\text{H}^+$ and $\text{H}_2\text{O}_2/\text{OH}^-$
(3) H^+ and OH^-

- (2) Heat and O_3/Δ
(4) Cooling and Heating




Product 'A' is




SPACE FOR ROUGH WORK

- Q.36 Which of the following is NOT true ?
 (1) The catalyst ZSM-5 converts alcohols directly into gasoline (petrol).
 (2) Charge on Lyophilic colloids depends on pH of medium.
 (3) The charged colloidal particles of the sol formed by addition of FeCl_3 in excess NaOH (aq.) move towards cathode during electrophoresis.
 (4) Physisorption is reversible in nature
- Q.37 Select the **incorrect** match for the extraction process involved for the given metal ore.
 (1) Cu_2S : Self reduction
 (2) $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$: Carbon Reduction
 (3) CuFeS_2 : Froth floatation
 (4) FeS_2 : Self Reduction
- Q.38 $\text{Me}-\text{CH}=\text{CH}-\text{CH}_2\text{OH} \xrightarrow[\text{ZnCl}_2]{\text{conc. HCl}}$
 Which statement is correct ?
 (1) Both $\text{Me}-\text{CH}=\text{CH}-\text{CH}_2\text{Cl}$ (I) and $\text{Me}-\text{CH}(\text{Cl})-\text{CH}=\text{CH}_2$ (II) are formed
 (2) Only (I) is formed
 (3) Only (II) is formed
 (4) Both are not formed
- Q.39 What is the pH of the solution obtained by mixing equal volumes of two solutions having pH values 9 and 11? Assume no components of the two solutions reacts. [Given : $\log 5 = 0.7$]
 (1) 3.3
 (2) 10.7
 (3) 11.3
 (4) 10.3
- Q.40 A species 'X' can show reaction with both HCl and NaOH . 'X' cannot be :
 (1) Al_2O_3
 (2) Zn
 (3) PbS
 (4) ZnCO_3

 SPACE FOR ROUGH WORK



Select correct option for reagent A, B, C, D ?

	(A)	(B)	(C)	(D)
(1)	H_2/Pt	HCO_2H	cold alk. KMnO_4	$\text{HgSO}_4/\text{dil. H}_2\text{SO}_4$
(2)	$\text{H}_2/\text{Pd-BaSO}_4$	mCPBA	hot alk. KMnO_4	$\text{Hg}(\text{OAc})_2/\text{H}_2\text{O}, \text{NaBH}_4$
(3)	$\text{B}_2\text{H}_6/\text{THF}, \text{CH}_3\text{COOH}$	PhCO_3H	$\text{OsO}_4/\text{NaHSO}_3$	$\text{B}_2\text{H}_6/\text{THF}, \text{H}_2\text{O}_2/\text{OH}^-$
(4)	$(\text{Sia})_2\text{BH}/\text{THF}, \text{CH}_3\text{COOH}$		Baeyer's Reagent	$\text{Hg}(\text{OCOCH}_3)_2/\text{H}_2\text{O}, \text{NaBH}_4$


Q.42 The molar mass of a gas is 50 g / mol. The density of the gas at critical temperature and critical pressure of 30 atm is 125 g / L. What is the critical temperature of the gas?

[Take : $R = 0.08 \text{ L atm mol}^{-1} \text{ K}^{-1}$]


- (1) 200 K (2) 500 K (3) 300 K (4) 400 K

SPACE FOR ROUGH WORK

- Q.43 The ionic radii (in Å) of N^{3-} , O^{2-} , F^- are respectively
 (1) 1.71, 1.40 and 1.36 (2) 1.71, 1.36 and 1.40
 (3) 1.36, 1.40 and 1.71 (4) 1.36, 1.71 and 1.40




'X' would be




- Q.45 A photon of energy 9.4 eV strikes to the electron present in third excited state of He^+ . What is the wavelength of the electron after absorption of the 9.4 eV energy of the photon ?
 (1) 4 Å (2) 1.8 Å (3) 5 Å (4) 6.65 Å

- Q.46 In which of the following complex, ligands are considered as strong field ligands (SFL) ?
 (1) $[\text{NiCl}_4]^{2-}$ (2) $[\text{PtCl}_4]^{2-}$ (3) $[\text{Fe}(\text{NH}_3)_6]^{2+}$ (4) $[\text{FeF}_6]^{3-}$



Product of reaction will be :




SPACE FOR ROUGH WORK

- Q.48 A mixture of NaHC_2O_4 and $\text{H}_2\text{C}_2\text{O}_4$ requires 50 mL, 0.1 M KMnO_4 (aq.) solution during titration in acidic medium. The same mass of NaHC_2O_4 and $\text{H}_2\text{C}_2\text{O}_4$ requires 50 mL, 0.4 M NaOH (aq.) solution for the complete neutralisation. Calculate the mass of $\text{H}_2\text{C}_2\text{O}_4$ in the initial mixture.
- (1) 1.250 g (2) 0.900 g (3) 0.450 g (4) 0.675 g

- Q.49 Which of the following reactions is not feasible ?
- (1) $[\text{Fe}(\text{en})_3]^{+3} + 6\text{CN}^- \longrightarrow [\text{Fe}(\text{CN})_6]^{3-} + 3 \text{ en}$
 (2) $[\text{Co}(\text{en})_3]^{+3} + \text{EDTA}^{4-} \longrightarrow [\text{Co}(\text{EDTA})]^{-1} + 3 \text{ en}$
 (3) $[\text{Cr}(\text{CO})_6] + 4\text{NO} \longrightarrow [\text{Cr}(\text{NO})_4] + 6\text{CO}$
 (4) $[\text{Fe}(\text{CO})_5] \xrightarrow{\text{Dimerisation}} [\text{Fe}_2(\text{CO})_{10}]$

- Q.50 In which reaction incorrect product is formed?




- Q.51 Identify the **incorrect** statement :

- (1) During an adiabatic expansion of an ideal gas temperature will always decrease.
 (2) During an isothermal expansion of a real gas temperature will always remain constant throughout.
 (3) $\Delta_{\text{rxn}} S$ must be zero for $\text{A(g)} + \text{B(g)} \rightarrow 2\text{AB(g)}$
 (4) During reversible adiabatic expansion of an ideal gas entropy of system will remain constant.





- Q.52 Which of the following would not result in the formation of paramagnetic substance ?



SPACE FOR ROUGH WORK



A, B, C are :




Q.54 In which of the following option the property of the given substance is **wrongly** matched?

- (1) CrO_2 – Ferromagnetic
 (2) MnO – Antiferromagnetic
 (3) C_6H_6 – Ferrimagnetic
 (4) Fe_3O_4 – Ferrimagnetic





Q.55 Hybridisation of underlined atom would not change after the given reaction in :

- (1) $\underline{\text{P}}\text{Cl}_3 + \text{Cl}_2 \longrightarrow$
 (2) $\text{H}_2\underline{\text{S}}\text{O}_4 \xrightarrow{\Delta}$
 (3) $\underline{\text{N}}\text{Cl}_3 + \text{H}_2\text{O} \longrightarrow$
 (4) $\underline{\text{Xe}}\text{F}_4 + \text{F}^- \longrightarrow$

SPACE FOR ROUGH WORK



A will be :



Q.57 Calculate the weight of urea (NH_2CONH_2) which must be dissolved in 490 g water so that the solution obtained has vapour pressure 2% less than vapour pressure of pure water.

- (1) 60 g (2) 30 g (3) 33.33 g (4) 40 g

Q.58 Which of the following option is incorrect about NO_2 and ClO_2 ?

- (1) Both are paramagnetic species (2) Both have bent shape
(3) Both compounds dimerised readily. (4) Both have sp^2 hybridisation

Q.59 Which statement is incorrect ?

- (1) Proline has 2° amine group
(2) D-Glucose and D-Fructose form same product on reduction by red P / HI
(3) D-Glucose and D-Mannose form different product on reaction with 3 eq. of phenyl hydrazine.
(4) Sucrose is non reducing carbohydrate

Q.60 For the reaction : $\text{A} \longrightarrow \text{Product}$ having order of the reaction 2.5. The half life period of the reaction is

given as $t_{1/2} \propto \frac{1}{[\text{A}]_0^m}$.

[Here, $[\text{A}]_0$ is the initial concentration of A]. Then the value of "m" is :

- (1) 0.5 (2) 3.5 (3) 2.5 (4) 1.5

SPACE FOR ROUGH WORK

MATHEMATICS

- Q.61 The value of $\left| (\vec{r} \cdot \vec{a})(\vec{b} \times \vec{c}) + (\vec{r} \cdot \vec{b})(\vec{c} \times \vec{a}) + (\vec{r} \cdot \vec{c})(\vec{a} \times \vec{b}) \right|$, where $|\vec{r}| = 2$ and $\vec{a}, \vec{b}, \vec{c}$ are coterminous sides of a tetrahedron with volume 3, is equal to
 (1) 6 (2) 12 (3) 18 (4) 36
- Q.62 For a twice differentiable non-constant function $f(x)$, if $f''(x) = f'(x)$, then the value of $\lim_{x \rightarrow \infty} \frac{f(x)}{f'(x)}$ is
 (1) 1 (2) -1 (3) 0 (4) Does not exist
- Q.63 If x, y, z satisfy the system of equations $\tan^2 x + \cot^2 x = 2 \cos^2 y$ and $\cos^2 y + \sin^2 z = 1$, then the value of $\int_{\cos^2 y}^{\sec^2 x + \sec^2 z} \frac{t^2}{t^2 - 4t + 8} dt$ is equal to
 (1) 0 (2) 1 (3) 2 (4) 3
- Q.64 The value of ${}^{50}C_0 \cdot {}^{50}C_{20} - {}^{50}C_1 \cdot {}^{50}C_{21} + {}^{50}C_2 \cdot {}^{50}C_{22} - {}^{50}C_3 \cdot {}^{50}C_{23} + \dots + {}^{50}C_{30} \cdot {}^{50}C_{50}$ is equal to
 (1) ${}^{100}C_{40}$ (2) ${}^{50}C_{15}$ (3) ${}^{50}C_{20}$ (4) ${}^{50}C_{10}$
- Q.65 The total number of ways of selecting 5 balls out of 20 identical red balls, 10 identical blue balls and 6 identical green balls is
 (1) ${}^{36}C_5$ (2) ${}^{20}C_5 + {}^{10}C_5 + {}^6C_5$
 (3) 21 (4) 42
- Q.66 A dice is thrown six times, it is being known that each time a different digit is shown. The probability that a sum of 11 will be obtained in the first three throws is
 (1) $\frac{3}{20}$ (2) $\frac{5}{24}$ (3) $\frac{25}{216}$ (4) $\frac{1}{12}$

SPACE FOR ROUGH WORK

- Q.67 The value of 'a' for which the straight line $4x - 3y + 4z - 2 = 0 = 3x - 2y + z - 5$ is parallel to the plane $2x - y + az - 7 = 0$, will be
 (1) 8 (2) -8 (3) 11 (4) -2
- Q.68 Mean of 50 observations is 20. If the sum of 40 observations is 300, then the mean of remaining 10 observation is
 (1) 30 (2) 50 (3) 70 (4) 10
- Q.69 If the equations $ax^2 + bx + c = 0$ and $x^3 + 6x^2 + 12x + 9 = 0$ have two common roots, ($a, b, c \in \mathbb{R}$), then
 (1) $a = 3b = 3c$ (2) $a = b = c$ (3) $a = -3b = c$ (4) $3a = b = c$
- Q.70 Let $f(x)$ be a twice differentiable function such that $f''(x) > 0$ in $x \in [0, 1]$, then
 (1) $f(0) + f(1) = 4f(c)$ for some $c \in (0, 1)$ (2) $f(0) + f(1) = 2f\left(\frac{1}{2}\right)$
 (3) $f(0) + f(1) > 2f\left(\frac{1}{2}\right)$ (4) $f(0) + f(1) < 2f\left(\frac{1}{2}\right)$
- Q.71 The area of the figure bounded by two branches of curve $(y - x)^2 = x^3$ and the line $x = 1$ is
 (1) $\frac{1}{3}$ (2) $\frac{4}{5}$ (3) $\frac{5}{4}$ (4) 3
- Q.72 Let $a > 0$ and $b = \lim_{x \rightarrow 0} \frac{a(1 - 4\sqrt{a^2 - x^2})}{x^2\sqrt{a^2 - x^2}}$ is finite, then the value of ab is
 (1) 1 (2) 2 (3) 4 (4) 8
- Q.73 The fundamental period of the function $\sin\left(\frac{\pi[x]}{12}\right) + \tan\left(\frac{\pi[x]}{3}\right) + \cos\left(\frac{\pi x}{4}\right)$ is
 [Note : where $[\cdot]$ denotes greatest integer function.]
 (1) 12 (2) 24 (3) 36 (4) function is non-periodic

SPACE FOR ROUGH WORK

- Q.74 Let A and B are two sets then $(A \cup B)' \cup (A' \cap B)$ is equal to
 (1) A' (2) A (3) B' (4) None of these
- Q.75 If p, q, r are statements with truth values false, true and false respectively, then the truth value of $(\sim p \vee \sim q) \vee r$ is
 (1) True (2) False (3) False if r is true (4) false if q is false
- Q.76 Consider four 2×2 non singular matrices A, B, C, D such that $ABC = BCD$, then
 (1) $A = D$ (2) $A = BCDB^{-1}C^{-1}$ (3) $A = C^{-1}B^{-1}DCB$ (4) $A = BCDC^{-1}B^{-1}$
- Q.77 If roots of the equation $z^6 = 1$ are 1, z_1, z_2, z_3, z_4 and z_5 , then the value of $\sum_{k=1}^5 z_k^5$ is
 (1) 0 (2) -1 (3) 6 (4) 5
- Q.78 The equation of axis and directrix of parabola $x^2 + 4x + 4y + 16 = 0$ are given by respectively
 (1) $x + 2 = 0, y - 2 = 0$ (2) $x = 2$ & $y = 2$
 (3) $x + 2 = 0$ and $y + 2 = 0$ (4) None of these
- Q.79 Product of length of perpendiculars drawn from foci upon a variable tangent to the hyperbola $\frac{(3x + 4y - 1)^2}{40} - \frac{(4x - 3y + 1)^2}{100} = 1$ is equal to
 (1) 40 (2) 100 (3) 8 (4) 4
- Q.80 If $\int \frac{(\sqrt{x})^5}{(\sqrt{x})^7 + x^6} dx = a \ln \left(\frac{x^k}{x^k + 1} \right) + c$, the value of a & k respectively are
 (1) $\frac{5}{2}$ and $\frac{2}{5}$ (2) $\frac{2}{5}$ and $\frac{5}{2}$ (3) $\frac{5}{2}$ and 2 (4) None of these

SPACE FOR ROUGH WORK

- Q.81 The ratio in which the plane $\vec{r} \cdot (\hat{i} - 2\hat{j} + 3\hat{k}) = 17$ divides the line joining the points $-2\hat{i} + 4\hat{j} + 7\hat{k}$ and $3\hat{i} - 5\hat{j} + 8\hat{k}$ is given by
 (1) 1 : 5 (2) 1 : 10 (3) 3 : 5 (4) 3 : 10
- Q.82 The number of pairs of integers (x, y) that satisfy the following two equations

$$\begin{cases} \cos(xy) = x \\ \tan(xy) = y \end{cases}, \text{ is(are)}$$

 (1) 1 (2) 2 (3) 4 (4) 6
- Q.83 Let $f\left(x + \frac{1}{2}\right) = f\left(x - \frac{1}{2}\right)$ and $f\left(\frac{-1}{2}\right) = \frac{3}{2}$. If $g(x) = \int_0^x f(t+n) dt$ for all $n \in \mathbb{N}$. Then $g'\left(\frac{5}{2}\right)$ is equal to
 (1) $\frac{1}{2}$ (2) $\frac{3}{2}$ (3) $\frac{5}{2}$ (4) none of these
- Q.84 The population $P(t)$ at time t of a certain mouse species follows the differential equation

$$\frac{d(P(t))}{dt} = 0.5P(t) - 450, \text{ if } P(0) = 850. \text{ Then the time at which the population becomes zero is}$$

 (1) $\ln 9$ (2) $\frac{\ln 18}{2}$ (3) $\ln 18$ (4) $2\ln 18$
- Q.85 If the point (2, 3) lies inside the circle $x^2 + y^2 - 6x - 10y + k = 0$ and the circle neither touches nor intersect the line $y = 1$. Then the set of all possible value of k lies in the interval
 (1) no value of k (2) (9, 29) (3) (18, 29) (4) (25, 31)
- Q.86 If $f(x) = \frac{1}{1-x}$ where $x \neq 0, 1$. Then the graph of the function $y = f(f(f(x)))$, $x > 1$ is
 (1) circle (2) ellipse (3) straight line (4) hyperbola

SPACE FOR ROUGH WORK

Q.87 The co-ordinate of a point P on the line $x + 2y + 5 = 0$ such that $|PA - PB|$ is minimum, where A (3, 4) & B (1, 2), is equal to

- (1) $\left(-\frac{7}{3}, -\frac{4}{3}\right)$ (2) $(-5, 0)$ (3) $\left(-\frac{9}{13}, -\frac{28}{13}\right)$ (4) $(15, -10)$

Q.88 The value of $\prod_{r=1}^3 \sin\left(\frac{2r\pi}{7}\right)$ is equal to

- (1) $\frac{1}{8}$ (2) $\frac{7}{64}$ (3) $\frac{3}{4}$ (4) $\frac{\sqrt{7}}{8}$

Q.89 Let $f(x) = \begin{cases} \operatorname{sgn}(\cot^{-1} x - \cot^{-1}(x^2)) & ; x \neq 0 \\ 1 & ; x = 0 \end{cases}$

(where sgn denotes signum function), then

- (1) $f(x)$ is continuous at $x=0$
 (2) $f(x)$ is continuous at $x=0$ but not differentiable at $x=0$
 (3) $f(x)$ is discontinuous at $x=0$
 (4) $f(x)$ is differentiable at $x=0$

Q.90 Consider a variable plane $lx + my + nz = k$ ($k > 0$) and l, m, n are direction cosines of normal of the plane. Let the given plane intersects the co-ordinate axes at A, B and C, then the minimum area of ΔABC is

- (1) $\frac{3\sqrt{3}k^2}{2}$ (2) $\frac{3\sqrt{3}k^2}{4}$ (3) $3\sqrt{3}k^2$ (4) $12\sqrt{3}k^2$

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK