


Chemical Kinetics

- Units of rate constant of first and zero order reactions in terms of molarity M unit are respectively [2002]
 (a) sec^{-1} , Msec^{-1} (b) sec^{-1} , M
 (c) Msec^{-1} , sec^{-1} (d) M, sec^{-1} .
- For the reaction $A + 2B \rightarrow C$, rate is given by $R = [A][B]^2$ then the order of the reaction is [2002]
 (a) 3 (b) 6
 (c) 5 (d) 7.
- The differential rate law for the reaction $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$ is [2002]
 (a) $-\frac{d[\text{H}_2]}{dt} = -\frac{d[\text{I}_2]}{dt} = -\frac{d[\text{HI}]}{dt}$
 (b) $\frac{d[\text{H}_2]}{dt} = \frac{d[\text{I}_2]}{dt} = \frac{1}{2} \frac{d[\text{HI}]}{dt}$
 (c) $\frac{1}{2} \frac{d[\text{H}_2]}{dt} = \frac{1}{2} \frac{d[\text{I}_2]}{dt} = -\frac{d[\text{HI}]}{dt}$
 (d) $-2 \frac{d[\text{H}_2]}{dt} = -2 \frac{d[\text{I}_2]}{dt} = \frac{d[\text{HI}]}{dt}$
- If half-life of a substance is 5 yrs, then the total amount of substance left after 15 years, when initial amount is 64 grams is [2002]
 (a) 16 grams (b) 2 grams
 (c) 32 grams (d) 8 grams.
- The integrated rate equation is [2002]
 $Rt = \log C_0 - \log C_t$
 The straight line graph is obtained by plotting
 (a) time vs $\log C_t$ (b) $\frac{1}{\text{time}}$ vs C_t
 (c) time vs C_t (d) $\frac{1}{\text{time}}$ vs $\frac{1}{C_t}$
- The half-life of a radioactive isotope is three hours. If the initial mass of the isotope were 256 g, the mass of it remaining undecayed after 18 hours would be [2003]
 (a) 8.0 g (b) 12.0 g
 (c) 16.0 g (d) 4.0 g
- In respect of the equation $k = Ae^{-E_a/RT}$ in chemical kinetics, which one of the following statements is correct ? [2003]
 (a) A is adsorption factor
 (b) E_a is energy of activation
 (c) R is Rydberg's constant
 (d) k is equilibrium constant
- For the reaction system : [2003]
 $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{NO}_2(\text{g})$ volume is suddenly reduced to half its value by increasing the pressure on it. If the reaction is of first order with respect to O_2 and second order with respect to NO, the rate of reaction will
 (a) diminish to one-eighth of its initial value
 (b) increase to eight times of its initial value
 (c) increase to four times of its initial value
 (d) diminish to one-fourth of its initial value
- In a first order reaction, the concentration of the reactant, decreases from 0.8 M to 0.4 M in 15 minutes. The time taken for the concentration to change from 0.1 M to 0.025 M is [2004]
 (a) 7.5 minutes (b) 15 minutes
 (c) 30 minutes (d) 60 minutes
- The rate equation for the reaction $2A + B \rightarrow C$ is found to be: $\text{rate} = k[A][B]$. The correct statement in relation to this reaction is that the [2004]
 (a) rate of formation of C is twice the rate of disappearance of A
 (b) $t_{1/2}$ is a constant
 (c) unit of k must be s^{-1}
 (d) value of k is independent of the initial concentrations of A and B

c-86

Chemistry

11. The half-life of a radioisotope is four hours. If the initial mass of the isotope was 200 g, the mass remaining after 24 hours undecayed is [2004]
 (a) 3.125 g (b) 2.084 g
 (c) 1.042 g (d) 4.167 g
12. A reaction involving two different reactants can never be [2005]
 (a) bimolecular reaction
 (b) second order reaction
 (c) first order reaction
 (d) unimolecular reaction
13. A schematic plot of $\ln K_{eq}$ versus inverse of temperature for a reaction is shown below [2005]



- The reaction must be
 (a) highly spontaneous at ordinary temperature
 (b) one with negligible enthalpy change
 (c) endothermic
 (d) exothermic
14. $t_{\frac{1}{4}}$ can be taken as the time taken for the concentration of a reactant to drop to $\frac{3}{4}$ of its initial value. If the rate constant for a first order reaction is K , the $t_{\frac{1}{4}}$ can be written as [2005]
 (a) $0.75/K$ (b) $0.69/K$
 (c) $0.29/K$ (d) $0.10/K$
15. Consider an endothermic reaction $X \rightarrow Y$ with the activation energies E_b and E_f for the backward and forward reactions, respectively. In general [2005]
 (a) there is no definite relation between E_b and E_f
 (b) $E_b = E_f$
 (c) $E_b > E_f$
 (d) $E_b < E_f$
16. A reaction was found to be second order with respect to the concentration of carbon monoxide. If the concentration of carbon monoxide is doubled, with everything else kept the same, the rate of reaction will [2006]
 (a) increase by a factor of 4
 (b) double
 (c) remain unchanged
 (d) triple
17. Rate of a reaction can be expressed by Arrhenius equation as : [2006]
 $k = A e^{-E/RT}$
 In this equation, E represents
 (a) the total energy of the reacting molecules at a temperature, T
 (b) the fraction of molecules with energy greater than the activation energy of the reaction
 (c) the energy above which all the colliding molecules will react
 (d) the energy below which all the colliding molecules will react
18. The following mechanism has been proposed for the reaction of NO with Br_2 to form NOBr :
 $NO(g) + Br_2(g) \rightleftharpoons NOBr_2(g)$
 $NOBr_2(g) + NO(g) \longrightarrow 2NOBr(g)$
 If the second step is the rate determining step, the order of the reaction with respect to NO(g) is [2006]
 (a) 3 (b) 2
 (c) 1 (d) 0
19. The energies of activation for forward and reverse reactions for $A_2 + B_2 \rightleftharpoons 2AB$ are 180 kJ mol^{-1} and 200 kJ mol^{-1} respectively. The presence of a catalyst lowers the activation energy of both (forward and reverse) reactions by 100 kJ mol^{-1} . The enthalpy change of the reaction ($A_2 + B_2 \rightarrow 2AB$) in the presence of a catalyst will be (in kJ mol^{-1}) [2007]
 (a) 20 (b) 300
 (c) 120 (d) 280
20. Consider the reaction, $2A + B \rightarrow \text{products}$. When concentration of B alone was doubled, the half-life did not change. When the concentration of A alone was doubled, the rate

Chemical Kinetics

c-87

- increased by two times. The unit of rate constant for this reaction is [2007]
 (a) s^{-1} (b) $L \text{ mol}^{-1} s^{-1}$
 (c) no unit (d) $\text{mol L}^{-1} s^{-1}$.
21. A radioactive element gets spilled over the floor of a room. Its half-life period is 30 days. If the initial velocity is ten times the permissible value, after how many days will it be safe to enter the room? [2007]
 (a) 100 days (b) 1000 days
 (c) 300 days (d) 10 days.
22. For a reaction $\frac{1}{2}A \rightarrow 2B$, rate of disappearance of 'A' is related to the rate of appearance of 'B' by the expression [2008]
 (a) $-\frac{d[A]}{dt} = \frac{1}{2} \frac{d[B]}{dt}$ (b) $-\frac{d[A]}{dt} = \frac{1}{4} \frac{d[B]}{dt}$
 (c) $-\frac{d[A]}{dt} = \frac{d[B]}{dt}$ (d) $-\frac{d[A]}{dt} = 4 \frac{d[B]}{dt}$
23. The half life period of a first order chemical reaction is 6.93 minutes. The time required for the completion of 99% of the chemical reaction will be ($\log 2 = 0.301$) [2009]
 (a) 23.03 minutes (b) 46.06 minutes
 (c) 460.6 minutes (d) 230.03 minutes
24. The time for half life period of a certain reaction $A \rightarrow \text{Products}$ is 1 hour. When the initial concentration of the reactant 'A', is 2.0 mol L^{-1} , how much time does it take for its concentration to come from 0.50 to 0.25 mol L^{-1} if it is a zero order reaction? [2010]
 (a) 4 h (b) 0.5 h
 (c) 0.25 h (d) 1 h
25. Consider the reaction :
 $\text{Cl}_2(\text{aq}) + \text{H}_2\text{S}(\text{aq}) \rightarrow$
 $\text{S}(\text{s}) + 2\text{H}^+(\text{aq}) + 2\text{Cl}^-(\text{aq})$
 The rate equation for this reaction is
 $\text{rate} = k[\text{Cl}_2][\text{H}_2\text{S}]$
 Which of these mechanisms is/are consistent with this rate equation? [2010]
 A. $\text{Cl}_2 + \text{H}_2\text{S} \rightarrow \text{H}^+ + \text{Cl}^- + \text{Cl}^+ + \text{HS}^-$ (slow)
 $\text{Cl}^+ + \text{HS}^- \rightarrow \text{H}^+ + \text{Cl}^- + \text{S}$ (fast)
 B. $\text{H}_2\text{S} \rightleftharpoons \text{H}^+ + \text{HS}^-$ (fast equilibrium)
 $\text{Cl}_2 + \text{HS}^- \rightarrow 2\text{Cl}^- + \text{H}^+ + \text{S}$ (Slow)
 (a) B only (b) Both A and B
 (c) Neither A nor B (d) A only
26. A reactant (A) forms two products : [2011RS]
 $A \xrightarrow{k_1} B$, Activation Energy Ea_1
 $A \xrightarrow{k_2} C$, Activation Energy Ea_2
 If $Ea_2 = 2 Ea_1$, then k_1 and k_2 are related as :
 (a) $k_2 = k_1 e^{Ea_1/RT}$ (b) $k_2 = k_1 e^{Ea_2/RT}$
 (c) $k_1 = A k_2 e^{Ea_1/RT}$ (d) $k_1 = 2 k_2 e^{Ea_2/RT}$
27. For a first order reaction $A \rightarrow \text{products}$ the concentration of A changes from 0.1 M to 0.025 M in 40 minutes.
 The rate of reaction when the concentration of A is 0.01 M is : [2012]
 (a) $1.73 \times 10^{-5} \text{ M/min}$ (b) $3.47 \times 10^{-4} \text{ M/min}$
 (c) $3.47 \times 10^{-5} \text{ M/min}$ (d) $1.73 \times 10^{-4} \text{ M/min}$
28. The rate of a reaction doubles when its temperature changes from 300 K to 310 K . Activation energy of such a reaction will be : ($R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ and $\log 2 = 0.301$) [2013]
 (a) 53.6 kJ mol^{-1} (b) 48.6 kJ mol^{-1}
 (c) 58.5 kJ mol^{-1} (d) 60.5 kJ mol^{-1}
29. For the non - stoichiometric reaction
 $2A + B \rightarrow C + D$, the following kinetic data were obtained in three separate experiments, all at 298 K .
- | Initial Concentration (A) | Initial Concentration (B) | Initial rate of formation of C ($\text{mol L}^{-1} \text{ s}^{-1}$) |
|---------------------------|---------------------------|---|
| 0.1 M | 0.1 M | 1.2×10^{-3} |
| 0.1 M | 0.2 M | 1.2×10^{-3} |
| 0.2 M | 0.1 M | 2.4×10^{-3} |
- The rate law for the formation of C is: [2014]
 (a) $\frac{dc}{dt} = k[A][B]$ (b) $\frac{dc}{dt} = k[A]^2[B]$
 (c) $\frac{dc}{dt} = k[A][B]^2$ (d) $\frac{dc}{dt} = k[A]$
30. Higher order (>3) reactions are rare due to : [JEE M 2015]
 (a) shifting of equilibrium towards reactants due to elastic collisions
 (b) loss of active species on collision
 (c) low probability of simultaneous collision of all the reacting species
 (d) increase in entropy and activation energy as more molecules are involved

c-88

Chemistry

31. Decomposition of H_2O_2 follows a first order reaction. In fifty minutes the concentration of H_2O_2 decreases from 0.5 to 0.125 M in one such decomposition. When the concentration of H_2O_2 reaches 0.05 M, the rate of formation of O_2 will be: [JEE M 2016]
- (a) 2.66 L min^{-1} at STP
 (b) $1.34 \times 10^{-2} \text{ mol min}^{-1}$
 (c) $6.96 \times 10^{-2} \text{ mol min}^{-1}$
 (d) $6.93 \times 10^{-4} \text{ mol min}^{-1}$
32. Two reactions R_1 and R_2 have identical pre-exponential factors. Activation energy of R_1 exceeds that of R_2 by 10 kJ mol^{-1} . If k_1 and k_2 are rate constants for reactions R_1 and R_2 respectively at 300 K, then $\ln(k_2/k_1)$ is equal to: ($R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$) [JEE M 2017]
- (a) 8 (b) 12
 (c) 6 (d) 4

Answer Key														
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
(a)	(a)	(d)	(d)	(a)	(d)	(b)	(b)	(c)	(d)	(a)	(d)	(d)	(c)	(d)
16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
(a)	(c)	(b)	(a)	(b)	(a)	(b)	(b)	(c)	(d)	(c)	(b)	(a)	(d)	(c)
31	32													
(d)	(d)													

SOLUTIONS

1. (a) For a zero order reaction.
 $\text{rate} = k[A]^0$ i.e. $\text{rate} = k$
 hence unit of $k = \text{M} \cdot \text{sec}^{-1}$
 For a first order reaction.
 $\text{rate} = k[A]$
 $k = \text{M} \cdot \text{sec}^{-1} / \text{M} = \text{sec}^{-1}$
2. (a) **NOTE** Order is the sum of the power of the concentrations terms in rate law expression.
 Hence the order of reaction is $= 1 + 2 = 3$
3. (d) rate of appearance of HI $= \frac{1}{2} \frac{d[\text{HI}]}{dt}$
 rate of formation of $\text{H}_2 = \frac{-d[\text{H}_2]}{dt}$
 rate of formation of $\text{I}_2 = \frac{-d[\text{I}_2]}{dt}$
 hence $\frac{-d[\text{H}_2]}{dt} = -\frac{d[\text{I}_2]}{dt} = \frac{1}{2} \frac{d[\text{HI}]}{dt}$
- or $-\frac{2d[\text{H}_2]}{dt} = -\frac{2d[\text{I}_2]}{dt} = \frac{d[\text{HI}]}{dt}$
4. (d) $t_{1/2} = 5$ years, $T = 15$ years hence total number of half life periods $= \frac{15}{5} = 3$.
 $\therefore \text{Amount left} = \frac{64}{(2)^3} = 8 \text{ g}$
5. (a) $Rt = \log C_0 - \log C_t$
 It is clear from the equation that if we plot a graph between $\log C_t$ and time, a straight line with a slope equal to $-\frac{k}{2.303}$ and intercept equal to $\log [A_0]$ will be obtained.
6. (d) $t_{1/2} = 3 \text{ hrs}$, $T = 18 \text{ hours}$
 $\therefore T = n \times t_{1/2}$
 $\therefore n = \frac{18}{3} = 6$
 Initial mass (C_0) = 256 g

$$\therefore C_n = \frac{C_0}{2^n} = \frac{256}{(2)^6} = \frac{256}{64} = 4g.$$

7. (b) In equation $k = Ae^{-E_a/RT}$;
 A = Frequency factor
 k = velocity constant, R = gas constant
and E_a = energy of activation

8. (b) $r = k [O_2][NO]^2$. When the volume is reduced to $1/2$, the conc. will double

$$\therefore \text{New rate} = k [2O_2][2NO]^2 = 8k [O_2][NO]^2$$

The new rate increases to eight times of its initial.

9. (c) As the concentration of reactant decreases from 0.8 to 0.4 in 15 minutes hence the $t_{1/2}$ is 15 minutes. To fall the concentration from 0.1 to 0.025 we need two half lives i.e., 30 minutes.


10. (d) The velocity constant depends on temperature only. It is independent of concentration of reactants.

11. (a) $N_t = N_0 \left(\frac{1}{2}\right)^n$ where n is number of half life periods.

$$n = \frac{\text{Total time}}{\text{half life}} = \frac{24}{4} = 6$$

$$\therefore N_t = 200 \left(\frac{1}{2}\right)^6 = 3.125g.$$

12. (d) The molecularity of a reaction is the number of reactant molecules taking part in a single step of the reaction.

 **NOTE** The reaction involving two different reactant can never be unimolecular.

13. (d) The graph show that reaction is exothermic.

$$\log k = \frac{-\Delta H}{RT} + 1$$

For exothermic reaction $\Delta H < 0$

$\therefore \log k$ Vs $\frac{1}{T}$ would be negative straight line with positive slope.

14. (c) $t_{1/4} = \frac{2.303}{K} \log \frac{1}{3/4} = \frac{2.303}{K} \log \frac{4}{3}$

$$= \frac{2.303}{K} (\log 4 - \log 3)$$

$$= \frac{2.303}{K} (2 \log 2 - \log 3)$$

$$= \frac{2.303}{K} (2 \times 0.301 - 0.4771) = \frac{0.29}{K}$$

15. (d) Enthalpy of reaction $(\Delta H) = E_{a(f)} - E_{a(b)}$
for an endothermic reaction $\Delta H = +ve$
hence for ΔH to be negative
 $E_{a(b)} < E_{a(f)}$

16. (a) Since the reaction is 2nd order w.r.t CO. Thus, rate law is given as.

$$r = k [CO]^2$$

Let initial concentration of CO is a i.e. $[CO] = a$

$$\therefore r_1 = k (a)^2 = ka^2$$


when concentration becomes doubled, i.e. $[CO] = 2a$

$$\therefore r_2 = k (2a)^2 = 4ka^2 \quad \therefore r_2 = 4r_1$$

So, the rate of reaction becomes 4 times.

17. (c) In Arrhenius equation $k = Ae^{-E/RT}$, E is the energy of activation, which is required by the colliding molecules to react resulting in the formation of products.

18. (b) (i) $NO(g) + Br_2(g) \rightleftharpoons NOBr_2(g)$



Rate law equation $= k[NOBr_2][NO]$

But $NOBr_2$ is intermediate and must not appear in the rate law equation

$$\text{from 1st step } K_C = \frac{[NOBr_2]}{[NO][Br_2]}$$

$$\therefore [NOBr_2] = K_C [NO][Br_2]$$

$$\therefore \text{Rate law equation} = k \cdot K_C [NO]^2 [Br_2]$$

hence order of reaction is 2 w.r.t. NO.

19. (a) $\Delta H_R = E_f - E_b = 180 - 200 = -20 \text{ kJ/mol}$

The nearest correct answer given in choices may be obtained by neglecting sign.

20. (b) For a first order reaction $t_{1/2} = \frac{0.693}{K}$ i.e.

for a first order reaction $t_{1/2}$ does not depend up on the concentration. From the given data, we can say that order of reaction with respect to $B = 1$ because change in concentration of B does not change half life.

Order of reaction with respect to $A = 1$ because rate of reaction doubles when concentration of B is doubled keeping concentration of A constant.

\therefore Order of reaction $= 1 + 1 = 2$ and units of second order reaction are $\text{L mol}^{-1} \text{sec}^{-1}$.

21. (a) Suppose activity of safe working $= A$


Given $A_0 = 10A$

$$\lambda = \frac{0.693}{t_{1/2}} = \frac{0.693}{30}$$

$$t_{1/2} = \frac{2.303}{\lambda} \log \frac{A_0}{A} = \frac{2.303}{0.693/30} \log \frac{10A}{A}$$

$$= \frac{2.303 \times 30}{0.693} \times \log 10 = 100 \text{ days.}$$

22. (b) The rates of reactions for the reaction



can be written either as

$$-2 \frac{d}{dt}[A] \text{ with respect to 'A'}$$

$$\text{or } \frac{1}{2} \frac{d}{dt}[B] \text{ with respect to 'B'}$$

From the above, we have

$$-2 \frac{d}{dt}[A] = \frac{1}{2} \frac{d}{dt}[B]$$

$$\text{or } -\frac{d}{dt}[A] = \frac{1}{4} \frac{d}{dt}[B]$$

i.e., correct answer is (b)

23. (b) For first order reaction,

$$k = \frac{2.303}{t} \log \frac{100}{100 - 99}$$

$$\frac{0.693}{6.93} = \frac{2.303}{t} \log \frac{100}{1}$$

$$\frac{0.693}{6.93} = \frac{2.303 \times 2}{t}$$

$$t = 46.06 \text{ min}$$

24. (c) For the reaction



given $t_{1/2} = 1 \text{ hour}$

for a zero order reaction

$$t_{\text{completion}} = \frac{[A_0]}{k} = \frac{\text{initial conc.}}{\text{rate constant}}$$

$$\therefore t_{1/2} = \frac{[A_0]}{2k}$$

$$\text{or } k = \frac{[A_0]}{2t_{1/2}} = \frac{2}{2 \times 1} = 1 \text{ mol lit}^{-1} \text{ hr}^{-1}$$

Further for a zero order reaction

$$k = \frac{dx}{dt} = \frac{\text{change in concentration}}{\text{time}}$$

$$1 = \frac{0.50 - 0.25}{\text{time}}$$

$$\therefore \text{time} = 0.25 \text{ hr.}$$

25. (d) Since the slow step is the rate determining step hence if we consider option (1) we find

$$\text{Rate} = k[\text{Cl}_2][\text{H}_2\text{S}]$$

Now if we consider option (2) we find

$$\text{Rate} = k[\text{Cl}_2][\text{HS}^-] \quad \dots(1)$$

From equation (i)

$$k = \frac{[\text{H}^+][\text{HS}^-]}{\text{H}_2\text{S}}$$

$$\text{or } [\text{HS}^-] = \frac{k[\text{H}_2\text{S}]}{[\text{H}^+]}$$

Substituting this value in equation (1) we find

$$\text{Rate} = k[\text{Cl}_2]K \frac{[\text{H}_2\text{S}]}{[\text{H}^+]} = k' \frac{[\text{Cl}_2][\text{H}_2\text{S}]}{[\text{H}^+]}$$

hence only, mechanism (1) is consistent with the given rate equation.

26. (c) $k_1 = A_1 e^{-E_{a1}/RT} \quad \dots\dots\dots(i)$

$$k_2 = A_2 e^{-E_{a2}/RT} \quad \dots\dots\dots(ii)$$

Chemical Kinetics

c-91

On dividing eqn (i) from eqn. (ii)

$$\frac{k_1}{k_2} = \frac{A_1}{A_2} (E_{a1} - E_{a2}) / RT$$

.....(iii)

Given $E_{a2} = 2E_{a1}$

On substituting this value in eqn. (iii)

$$k_1 = k_2 A \times e^{E_{a1}/RT}$$

27. (b) For a first order reaction

$$k = \frac{2.303}{t} \log \frac{a}{a-x} = \frac{2.303}{40} \log \frac{0.1}{0.025}$$

$$= \frac{2.303}{40} \log 4 = \frac{2.303 \times 0.6020}{40}$$

$$= 3.47 \times 10^{-2}$$

$$R = k(A)^1 = 3.47 \times 10^{-2} \times 0.01$$

$$= 3.47 \times 10^{-4}$$

28. (a) Activation energy can be calculated from the equation

$$\frac{\log k_2}{\log k_1} = \frac{-E_a}{2.303R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

$$\text{Given } \frac{k_2}{k_1} = 2; T_2 = 310 \text{ K}; T_1 = 300 \text{ K}$$

$$= \log 2 = \frac{-E_a}{2.303 \times 8.314} \left(\frac{1}{310} - \frac{1}{300} \right)$$

$$E_a = 53598.6 \text{ J/mol} = 53.6 \text{ kJ/mol.}$$

29. (d) Let rate of reaction = $\frac{d[C]}{t} = k[A]^x[B]^y$

Now from the given data

$$1.2 \times 10^{-3} = k[0.1]^x[0.1]^y \quad \text{.....(i)}$$

$$1.2 \times 10^{-3} = k[0.1]^x[0.2]^y \quad \text{.....(ii)}$$

$$2.4 \times 10^{-3} = k[0.2]^x[0.1]^y \quad \text{.....(iii)}$$

Dividing equation (i) by (ii)

$$\Rightarrow \frac{1.2 \times 10^{-3}}{1.2 \times 10^{-3}} = \frac{k[0.1]^x[0.1]^y}{k[0.1]^x[0.2]^y}$$

We find, $y = 0$

Now dividing equation (i) by (iii)

$$\Rightarrow \frac{1.2 \times 10^{-3}}{2.4 \times 10^{-3}} = \frac{k[0.1]^x[0.1]^y}{k[0.2]^x[0.1]^y}$$

We find, $x = 1$

$$\text{Hence } \frac{d[C]}{dt} = k[A]^1[B]^0$$

30. (c) Reactions of higher order (>3) are very rare due to very less chances of many molecules to undergo effective collisions.

31. (d) $\text{H}_2\text{O}_2(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{aq}) + \frac{1}{2} \text{O}_2(\text{g})$

For a first order reaction

$$k = \frac{2.303}{t} \log \frac{a}{(a-x)}$$

Given $a = 0.5$, $(a-x) = 0.125$, $t = 50 \text{ min}$

$$\therefore k = \frac{2.303}{50} \log \frac{0.5}{0.125}$$

$$= 2.78 \times 10^{-2} \text{ min}^{-1}$$

$$r = k[\text{H}_2\text{O}_2] = 2.78 \times 10^{-2} \times 0.05$$

$$= 1.386 \times 10^{-3} \text{ mol min}^{-1}$$

Now

$$-\frac{d[\text{H}_2\text{O}_2]}{dt} = \frac{d[\text{H}_2\text{O}]}{dt} = \frac{2d[\text{O}_2]}{dt}$$

$$\therefore \frac{2d[\text{O}_2]}{dt} = -\frac{d[\text{H}_2\text{O}_2]}{dt}$$

$$\therefore \frac{d[\text{O}_2]}{dt} = \frac{1}{2} \times \frac{d[\text{H}_2\text{O}_2]}{dt}$$

$$= \frac{1.386 \times 10^{-3}}{2} = 6.93 \times 10^{-4} \text{ mol min}^{-1}$$

32. (d) From arrhenius equation,

$$k = A.e^{\frac{-E_a}{RT}}$$

$$\text{so, } k_1 = A.e^{-E_{a1}/RT} \quad \text{.....(1)}$$

$$k_2 = A.e^{-E_{a2}/RT} \quad \text{.....(2)}$$

On dividing equation (2) (1)

$$\Rightarrow \frac{k_2}{k_1} = e^{\frac{(E_{a1}-E_{a2})}{RT}}$$

$$\ln \left(\frac{k_2}{k_1} \right) = \frac{E_{a1} - E_{a2}}{RT} = \frac{10,000}{8.314 \times 300} = 4$$