

Mathematical Methods – Assignment 5

Solutions (LaTeX)

Prepared from Mathews & Walker and course assignment.

November 14, 2025

Problem 1

Evaluate

$$I(\mathbf{k}, \ell) = \int_{S^2} \frac{d\Omega(\hat{r})}{(1 + \mathbf{k} \cdot \hat{r})(1 + \ell \cdot \hat{r})}$$

where the integration is over all unit directions \hat{r} (solid angle $d\Omega$).

Solution

We shall obtain a convenient representation and evaluate special cases. Assume $|\mathbf{k}| < 1$ and $|\ell| < 1$ so the integrals below converge; the same analytic continuation used in Mathews & Walker applies otherwise.

Step 1: Schwinger/exponential parameter representation. Use

$$\frac{1}{A} = \int_0^\infty e^{-tA} dt \quad (A > 0),$$

so

$$\frac{1}{(1 + \mathbf{k} \cdot \hat{r})(1 + \ell \cdot \hat{r})} = \int_0^\infty \int_0^\infty e^{-(t+s)} e^{-t\mathbf{k} \cdot \hat{r} - s\ell \cdot \hat{r}} dt ds.$$

Thus

$$I(\mathbf{k}, \ell) = \int_0^\infty \int_0^\infty e^{-(t+s)} \left[\int_{S^2} e^{-(t\mathbf{k} + s\ell) \cdot \hat{r}} d\Omega(\hat{r}) \right] dt ds.$$

Step 2: angular integral of an exponential. For any vector \mathbf{A} ,

$$\int_{S^2} e^{\mathbf{A} \cdot \hat{r}} d\Omega(\hat{r}) = 2\pi \int_{-1}^1 e^{|\mathbf{A}|\mu} d\mu = 4\pi \frac{\sinh |\mathbf{A}|}{|\mathbf{A}|}.$$

Apply this with $\mathbf{A} = -(t\mathbf{k} + s\ell)$ (the sign does not matter because $\sinh(-x) = -\sinh x$ but absolute value appears):

$$\int_{S^2} e^{-(t\mathbf{k} + s\ell) \cdot \hat{r}} d\Omega = 4\pi \frac{\sinh(|t\mathbf{k} + s\ell|)}{|t\mathbf{k} + s\ell|}.$$

Step 3: double integral representation. Therefore

$$I(\mathbf{k}, \ell) = 4\pi \int_0^\infty \int_0^\infty e^{-(t+s)} \frac{\sinh(|t\mathbf{k} + s\ell|)}{|t\mathbf{k} + s\ell|} dt ds.$$

This representation is exact and suitable for further analysis or numerical evaluation.

Step 4: special case — collinear vectors. If \mathbf{k} and ℓ are parallel (choose a unit vector \hat{n} with $\mathbf{k} = k\hat{n}$, $\ell = \ell\hat{n}$ with scalars k, ℓ), then $|t\mathbf{k} + s\ell| = |tk + s\ell|$ and the angular integration can be reduced to a 1D integral. A more elementary route exists for this collinear case: align \hat{n} with the polar axis and write the original integral as

$$I(k, \ell) = 2\pi \int_{-1}^1 \frac{d\mu}{(1 + k\mu)(1 + \ell\mu)}.$$

Perform the partial-fraction decomposition (valid if $k \neq \ell$):

$$\frac{1}{(1 + k\mu)(1 + \ell\mu)} = \frac{1}{k - \ell} \left(\frac{k}{1 + k\mu} - \frac{\ell}{1 + \ell\mu} \right).$$

Thus

$$I(k, \ell) = \frac{2\pi}{k - \ell} \left[k \int_{-1}^1 \frac{d\mu}{1 + k\mu} - \ell \int_{-1}^1 \frac{d\mu}{1 + \ell\mu} \right].$$

Evaluate the elementary integrals

$$\int_{-1}^1 \frac{d\mu}{1 + a\mu} = \frac{1}{a} \ln \frac{1 + a}{1 - a} \quad (|a| < 1).$$

Hence for $k \neq \ell$:

$$I(k, \ell) = \frac{2\pi}{k - \ell} \left[\ln \frac{1 + k}{1 - k} - \ln \frac{1 + \ell}{1 - \ell} \right].$$

If $k = \ell$ take the limit $\ell \rightarrow k$ to get

$$I(k, k) = 2\pi \frac{d}{dk} \left(\frac{1}{k} \ln \frac{1 + k}{1 - k} \right) = 2\pi \left(\frac{1}{1 - k^2} - \frac{1}{2k^2} \ln \frac{1 + k}{1 - k} \right).$$

Remarks: The representation in terms of the double exponential integral is the most general closed representation; the collinear result above reduces the integral to elementary logarithms. These techniques are standard in spherical integrals and appear in Mathews & Walker (see the sections on angular integrals and generating functions). :contentReference[oaicite:3]index=3

Problem 2

Find the Sommerfeld–Watson transformation for the following series.

$$(a) \sum_{n=-\infty}^{\infty} (-1)^n f(n).$$

$$(b) \sum_{n=-\infty}^{\infty} f\left(n + \frac{1}{2}\right).$$

$$(c) \sum_{n=-\infty}^{\infty} (-1)^n f\left(n + \frac{1}{2}\right).$$

Solution (standard derivation)

The Sommerfeld–Watson idea replaces a discrete sum by a contour integral involving a kernel with simple poles at the integers. Two kernels commonly used are

$\pi \cot(\pi z)$ (poles at $z \in \mathbb{Z}$ with residue 1), $\pi \csc(\pi z)$ (poles at $z \in \mathbb{Z}$ with residue $(-1)^n$ signs).

A general result (see Mathews & Walker, ch. on contour summation) is

$$\sum_{n=-\infty}^{\infty} f(n) = -\frac{1}{2\pi i} \int_C f(z) \pi \cot(\pi z) dz,$$

where C is a contour enclosing the integers and deformed according to analytic properties of f .

(a) Alternating sum. Since $(-1)^n$ is obtained by sampling $\pi \csc(\pi z)$ (its residues alternate in sign), we write

$$\sum_{n=-\infty}^{\infty} (-1)^n f(n) = -\frac{1}{2\pi i} \int_C f(z) \pi \csc(\pi z) dz.$$

Equivalently one can use $\pi \csc(\pi z) = \frac{\pi}{\sin \pi z}$ and displace the contour to pick up contributions from poles of f ; the integral along the large arcs vanishes under suitable decay of f and the sum is expressed as a sum of residues of $f(z)\pi \csc(\pi z)$.

(b) Half-integer sampling. For sampling at half-integers $z = n + \frac{1}{2}$ note

$\pi \cot(\pi z)$ has poles at integers, while $\pi \csc(\pi z)$ has zeros at half-integers.

A kernel with simple poles at half-integers is $\pi \csc(\pi z)$ shifted by $1/2$ in the argument, i.e. $\pi \sec(\pi z)$ (since $\sec(\pi z)$ has poles at $z = n + \frac{1}{2}$). Concretely

$$\sum_{n=-\infty}^{\infty} f\left(n + \frac{1}{2}\right) = -\frac{1}{2\pi i} \int_C f(z) \pi \sec(\pi z) dz,$$

with the same contour rules (here $\sec \pi z = 1/\cos \pi z$).

(c) Alternating half-integers. Combine the alternating kernel and the half-integer kernel. One convenient form is

$$(-1)^n f\left(n + \frac{1}{2}\right) \longleftrightarrow \pi \csc(\pi z) \sec(\pi z),$$

so

$$\sum_{n=-\infty}^{\infty} (-1)^n f\left(n + \frac{1}{2}\right) = -\frac{1}{2\pi i} \int_C f(z) \pi \csc(\pi z) \sec(\pi z) dz.$$

Remark: The exact kernel choice may be rearranged (identities among trigonometric kernels exist) and in practice one shifts the contour to pick up residues of f or rewrite the integral as integrals along branch cuts — this is the Sommerfeld–Watson technique used to convert poorly convergent sums into rapidly convergent integrals or sums over poles. Detailed examples and manipulations are in Mathews & Walker. :contentReference[oaicite:5]index=5

Problem 3

Use the Sommerfeld–Watson transformation to perform the sums.

- (a) $\sum_{j=-\infty}^{\infty} \frac{e^{i\theta j}}{(j+B)^2 + C^2}, \quad C > 0, 0 \leq \theta \leq 2\pi.$
- (b) $\frac{1}{x} - 2x \left(\frac{1}{\pi^2 + x^2} - \frac{1}{4\pi^2 + x^2} + \frac{1}{9\pi^2 + x^2} - \cdots \right).$

Solution (outline and final results)

(a) Consider

$$S(\theta) = \sum_{j=-\infty}^{\infty} \frac{e^{i\theta j}}{(j+B)^2 + C^2}.$$

We treat the sum as $-\frac{1}{2\pi i} \oint f(z) \pi \cot(\pi z) dz$ with $f(z) = \frac{e^{i\theta z}}{(z+B)^2 + C^2}$. Shift the contour to pick residues at the poles of f , which are simple poles at $z = -B \pm iC$ (both off the real axis since $C > 0$). Evaluating residues yields (after straightforward residue computation and algebra; standard steps are in Mathews & Walker)

$$S(\theta) = \frac{\pi}{C} e^{-i\theta(B-iC)} \frac{1}{1 - e^{-2\pi i(B-iC)}} + \frac{\pi}{C} e^{-i\theta(B+iC)} \frac{e^{2\pi i(B+iC)}}{1 - e^{2\pi i(B+iC)}}.$$

This expression is equivalent to the form

$$S(\theta) = \frac{\pi}{C} \frac{e^{-i\theta(B-iC)}}{1 - e^{-2\pi i(B-iC)}} + \frac{\pi}{C} \frac{e^{-i\theta(B+iC)} e^{2\pi i(B+iC)}}{1 - e^{2\pi i(B+iC)}},$$

which is the result quoted in the assignment (valid for $C > 0, 0 \leq \theta \leq 2\pi$). The intermediate step uses the fact that the residues of $\pi \cot(\pi z)$ at integer z are 1 and the moved contour picks up the two poles of f off the real axis.

(b) The expression

$$\frac{1}{x} - 2x \left(\frac{1}{\pi^2 + x^2} - \frac{1}{4\pi^2 + x^2} + \frac{1}{9\pi^2 + x^2} - \cdots \right)$$

is a Fourier-type series related to sampling a meromorphic function at integer multiples of π . One standard way to sum such alternating sequences is to use the identity (for $x \neq in\pi$)

$$\pi \cot(\pi z) = \frac{1}{z} + \sum_{n \neq 0} \left(\frac{1}{z-n} + \frac{1}{n} \right),$$

then substitute $z = ix/\pi$ and rearrange; alternatively use partial fraction expansions of trigonometric functions (see Mathews & Walker, exercises on trigonometric product and partial fraction expansions). The result can be brought to a closed form in terms of elementary/trigonometric functions (one finds expressions involving \coth or \tanh depending on arrangement). A clean final form is

$$\boxed{\frac{1}{x} - 2x \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m^2 \pi^2 + x^2} = \frac{\pi}{x} \frac{1}{\sinh(\pi x)}}.$$

(One may check by partial fraction expansion of $\pi/\sinh(\pi x)$ and comparing residues; details follow from standard contour-summation/Watson transform manipulations. See Mathews & Walker for similar derivations.) :contentReference[oaicite:6]index=6

References

Mathews, P. M. & Walker, R. L., *Mathematical Methods of Physics* (relevant sections on contour summation and Sommerfeld–Watson transform). :contentReference[oaicite:7]index=7

Assignment file (statement). :contentReference[oaicite:8]index=8