

Assignment 2 – QT 207
All the problems are from the book
An Introduction to Quantum Computing by Kaye, Laflamme and Mosca

Exercise 3.3.1 Consider the 2-qubit state

$$|\psi\rangle = \frac{1}{\sqrt{2}}|0\rangle|0\rangle + \frac{1}{\sqrt{2}}|1\rangle|1\rangle. \quad (3.3.2)$$

Show that this state is entangled by proving that there are no possible values $\alpha_0, \alpha_1, \beta_0, \beta_1$ such that

$$|\psi\rangle = (\alpha_0|0\rangle + \alpha_1|1\rangle)(\beta_0|0\rangle + \beta_1|1\rangle). \quad (3.3.3)$$

Exercise 3.4.1

- (a) Prove that if the operators P_i satisfy $P_i^\dagger = P_i$ and $P_i^2 = P_i$, then $P_i P_j = 0$ for all $i \neq j$.
- (b) Prove that any pure state $|\psi\rangle$ can be decomposed as $|\psi\rangle = \sum_i \alpha_i |\psi_i\rangle$ where $\alpha_i = \sqrt{p(i)}$, $p(i) = \langle \psi | P_i | \psi \rangle$, and $|\psi_i\rangle = \frac{P_i |\psi\rangle}{\sqrt{p(i)}}$.

Also prove that $\langle \psi_i | \psi_j \rangle = \delta_{i,j}$.

- (c) Prove that any decomposition $I = \sum_i P_i$ of the identity operator on a Hilbert space of dimension N into a sum of nonzero projectors P_i can have at most N terms in the sum.

Exercise 3.4.2 Show that measuring the observable $|1\rangle\langle 1|$ is equivalent to measuring the observable Z up to a relabelling of the measurement outcomes.

Exercise 4.2.3

- (a) Prove $X R_y(\theta) X = R_y(-\theta)$ and $X R_z(\theta) X = R_z(-\theta)$.
- (b) Prove Corollary 4.2.1.

Hint: Using Theorem 4.2.1 we can write

$$U = e^{i\alpha} R_z(\beta) R_y(\gamma) R_z(\delta). \quad (4.2.12)$$

Then take $A \equiv R_z(\beta) R_y(\gamma/2)$, $B \equiv R_y(-\gamma/2) R_z(-(\delta + \beta)/2)$, and $C \equiv R_z((\delta - \beta)/2)$.

Assignment 2 – QT 207

All the problems are from the book

An Introduction to Quantum Computing by Kaye, Laflamme and Mosca

Exercise 4.2.4 Describe the effect of the CNOT gate with respect to the following bases.

$$(a) B_1 = \left\{ |0\rangle \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right), |0\rangle \left(\frac{|0\rangle - |1\rangle}{\sqrt{2}} \right), |1\rangle \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right), |1\rangle \left(\frac{|0\rangle - |1\rangle}{\sqrt{2}} \right) \right\}$$

$$(b) B_2 = \left\{ \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right) \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right), \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right) \left(\frac{|0\rangle - |1\rangle}{\sqrt{2}} \right), \left(\frac{|0\rangle - |1\rangle}{\sqrt{2}} \right) \left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \right), \left(\frac{|0\rangle - |1\rangle}{\sqrt{2}} \right) \left(\frac{|0\rangle - |1\rangle}{\sqrt{2}} \right) \right\}$$

Express your answers both using Dirac notation, and also with matrix notation.

Exercise 5.2.1 Prove that

$$|\psi\rangle|\beta_{00}\rangle = \frac{1}{2}|\beta_{00}\rangle|\psi\rangle + \frac{1}{2}|\beta_{01}\rangle(X|\psi\rangle) + \frac{1}{2}|\beta_{10}\rangle(Z|\psi\rangle) + \frac{1}{2}|\beta_{11}\rangle(XZ|\psi\rangle). \quad (5.2.8)$$

Exercise 6.4.1 Prove that

$$\left(\frac{|0\rangle + (-1)^{x_1}|1\rangle}{\sqrt{2}} \right) \left(\frac{|0\rangle + (-1)^{x_2}|1\rangle}{\sqrt{2}} \right) \dots \left(\frac{|0\rangle + (-1)^{x_n}|1\rangle}{\sqrt{2}} \right) \quad (6.4.14)$$

$$= \frac{1}{\sqrt{2^n}} \sum_{z_1 z_2 \dots z_n \in \{0,1\}^n} (-1)^{x_1 z_1 + x_2 z_2 + \dots + x_n z_n} |z_1\rangle |z_2\rangle \dots |z_n\rangle. \quad (6.4.15)$$

Exercise 6.5.1 Let $\mathbf{x}, \mathbf{y} \in \{0,1\}^n$ and let $\mathbf{s} = \mathbf{x} \oplus \mathbf{y}$. Show that

$$H^{\otimes n} \left(\frac{1}{\sqrt{2}}|\mathbf{x}\rangle + \frac{1}{\sqrt{2}}|\mathbf{y}\rangle \right) = \frac{1}{\sqrt{2^{n-1}}} \sum_{\mathbf{z} \in \{\mathbf{s}\}^\perp} (-1)^{\mathbf{x} \cdot \mathbf{z}} |\mathbf{z}\rangle. \quad (6.5.5)$$