

1 Title: Optimal control in the face of evolving resistance is an economic problem not a  
2 biological one.

3 Shaun R. Coutts, Helen Hicks, Alexa Varah, Kwadjo Ahodo, Rob Frekleton, Dylan Z.  
4 Childs

## Abstract

Evolved resistance to xenobiotics (e.g. antibiotics, herbicides, pesticides, fungicides) is a global threat to public health and food security. In agricultural systems non-chemical control methods can be combined with xenobiotics (Integrated Pest Management; IPM) to prolong the useful life of compounds and manage pest populations after resistance has evolved. We find IPM strategies with the highest economic returns for an arable cropping system, and perform a global sensitivity analysis to find the factors that shape those strategies. The key uncertainties are economic in nature, and farmers have an incentive to be responsive to changes in the shape of the yield loss function. Doing so effectively will require estimating, at a minimum, what yields would be in the absence of the pest, maximum weed density and how yields change with increasing pest density, with enough detail to say how much control (if any) is justified.

## Significance

Integrated pest management (IPM) applies chemical and non-chemical control methods to pest populations to manage evolved resistance. Surprisingly, despite the widespread adoption of IPM, we have a poor understanding of how to select between alternative IPM strategies. Here we consider a typical problem, a farming system in which high levels of herbicide resistance evolves repeatedly. The best IPM strategies were dependent on crop yields, yield loss caused by the weed and the maximum possible loss, land tenure, and levels of herbicide resistance. With the exception of herbicide resistance, all these factors are economic in nature, thus knowing which IPM strategy to apply where is an economic problem rather than biological one.

## Introduction

Controlling populations in the face evolving resistance to xenobiotics (i.e. antibiotics, herbicides, pesticides, fungicides) is one of the biggest challenges facing public health (Laxminarayan *et al.*, 2016; Willyard *et al.*, 2017), and food security (Denholm & Rowland, 1992; Palumbi, 2001; Hicks *et al.*, 2018). Evolved resistance also costs billions of dollars globally (Livingston *et al.*, 2016; Chesson *et al.*, 2018; Hicks *et al.*, 2018). In agricultural systems evolved resistance can be mitigated with integrated pest management (IPM), where chemical control is used in combination with non-chemical techniques such as crop rotation, cultivation and spot control (e.g. hand-weeding). IPM can be used both pro-actively to delay the evolution of resistance, and reactively to control pest populations as chemical control becomes less effective (Denholm & Rowland, 1992; Hicks *et al.*, 2018). While most current strategies to mitigate resistance focus on delaying resistance (REX Consortium, 2013), in many agricultural weeds, including *A. myosuroides*, herbicide resistance is already widespread (Hicks *et al.*, 2018). Thus, cost effective IPM strategies focusing on managing weeds in the face of existing resistance are needed.

While the concept of IPM is well established (Bottrell *et al.*, 1979), finding cost effective IPM strategies is extremely challenging (Dana *et al.*, 2014; Chalak & Pannell, 2015). Management tools need to be used in the correct combination and sequence to be effective. This results in a very large number of potential IPM strategies (i.e. different combinations and sequences), even when considering only a handful of management tools and short time horizons (Chalak & Pannell, 2015). As a result of this difficulty, there have been no attempts to rigorously search for cost effective IPM strategies in the face of rapidly evolving resistance to one of the primary management options. This is an important gap in our understanding for agricultural systems; where resistance to xenobiotics has evolved numerous times (Denholm & Rowland, 1992; Palumbi, 2001) and multiple non-chemical control options can be used in combination to deliver cost effective control (Chalak & Pannell, 2015).

Further, little is known about how robust good IPM strategies are to changes in factors such as crop yield and pest population dynamics (Epanchin-Niell & Hastings, 2010). Thus, even when effective IPM strategies are found, we do not know how generally they should apply. Robust IPM strategies would allow standard IPM strategies to be developed for entire farms or regions. If we cannot identify robust IPM strategies then farmers face the prospect of having to find workable IPM strategies for each economic and biological scenario they face. This would pose a serious barrier to IPM adoption, as IPM strategies are already seen as complex and difficult to implement (Llewellyn *et al.*, 2006). An alternative is to determine a small set of easily measurable factors that shape IPM strategies in predictable ways. So that even if cost-effective IPM strategies do change in response to economic conditions and population characteristics, it is at least straight forward to know which strategy to apply.

*Alopecurus myosuroides* was used as a test case. *A. myosuroides* is Europe’s most economically costly weed (Moss *et al.*, 2007) and one of the worlds most serious herbicide resistant grass weeds (Heap, 2014). Herbicide-resistant *A. myosuroides* costs up to £320·ha<sup>-1</sup> in yield loss and extra herbicide use (Hicks *et al.*, 2018), and imposes a total cost of £0.5bn across England (Varah *et al.*, in press). We built a population model for *A. myosuroides* where target site resistance to two herbicides is already present, although possibly at very low frequencies. To allow IPM strategies the population model has two seed bank levels so that cultivation can bury seeds, and spring crops and spot control (e.g. by hand weeding) affect survival.

We framed IPM as a combinatorial optimisation problem and used a genetic algorithm (Appendix 1) to find economically incentivised (measured by gross margin) IPM strategies (Taylor & Hastings, 2004; Carrasco *et al.*, 2010) in the face of evolution. We carried out the first global sensitivity analysis of dynamic IPM strategies (Appendix 2), testing how robust those IPM strategies were across 15,000 parameter combinations incorporate a wide range of economic, biological and psychological factors. We find that cost effective IPM strategies can change dramatically in response to changes in potential yield losses,

but those changes are predictable.

## Results and Discussion

Across a wide range of parameter combinations the most important parameters shaping IPM strategies relate to the immediate, or potential maximum, economic impact of the weed, despite much research on IPM focusing on biological factors (Colbach *et al.*, 2006). In particular parameters defining the yield loss function (Fig. 1a) have the most influence on shaping incentivized IPM strategies. The yield loss function describes the relationship between weed density and crop yield (Fig. 1b). Yield when the weed was absent ( $Y_0$ ) and the effect of weed density on yield ( $Y_D$ ) were two of the most important parameters (Fig. 1). The set of parameters that controls how large the seed bank can become ( $f_m$ ,  $f_d$  and  $\phi_b$ ; Fig. 1a), and therefore the the maximum possible loss if the weed population is uncontrolled (Fig. 1b), are also important. Although yield loss functions have been estimated for major weeds (Cousens, 1985; Doyle *et al.*, 1986; Swinton *et al.*, 1994), there is evidence that yield functions vary substantially between fields (Swinton *et al.*, 1994; Hicks *et al.*, 2018), and little attention has been paid to this variation and understanding its causes.

While the shape of the yield loss function is important because it determines how much control is justified, knowing which type of IPM strategy to employ may only require an estimate of the pest free yield ( $Y_0$ ), and one or two thresholds values of weed density where a new IPM strategy becomes advantageous. When the yield of winter wheat in the absence of *A. myosuroides* ( $Y_0$ ) is low, management intensity is lower and relies on crop rotation and tactical use of herbicide (Fig. 2, 'Y<sub>0</sub> low'). The strategy changes little when the effect of weed density on yield ( $Y_D$ ) increases. Although, more herbicide is used when the value of  $Y_D$  increases from a very low value to a slightly higher value (1% to 12% losses at high densities of *A. myosuroides*; Fig. 2g,e). When  $Y_0$  is high,  $Y_D$  shows two thresholds where IPM strategy changes. When  $Y_D$  is very low relative to the maximum *A. myosuroides*




Figure 1: a) Relative influence of each parameter on IPM strategy. Higher values indicate parameters with more influence on the structure of incentivized IPM strategies. To find which parameters were crucial in shaping IPM with high gross margin we used multi-variate boosted regression trees (Miller *et al.*, 2016) as a meta-model (Coutts & Yokomizo, 2014)(Appendix 2). We interrogated this meta-model to find important parameters accounting for high-level interactions and non-linear responses (Friedman, 2001; Miller *et al.*, 2016). Relative influence is the reduction in mean squared error attributable to each parameter. Because only relative values are only meaningful the index is re-scaled to sum to 100 across all parameters (Friedman, 2001). b) The most important parameters are related to the yield loss function. We use a linear yield function fit to empirical data (Appendix 3).




Figure 2: IPM strategies under high ( $\pounds 1668 \cdot \text{ha}^{-1}$ ) and low ( $\pounds 986 \cdot \text{ha}^{-1}$ ) values of  $Y_0$  (yield of winter wheat with no *A. myosuroides*), under increasing values (rows) of  $Y_D$  (in  $\pounds \cdot \text{plant}^{-1} \cdot \text{ha}^{-1}$ ). At the lower limit of  $Y_D$  very high *A. myosuroides* densities result in a 1% yield loss under the high  $Y_0$  scenario, and the upper limit implies a yield loss of 35%. There is initially one effective herbicide ( $R_{\text{int}} = 0.0001$ ,  $Q_{\text{int}} = 0.9$ ). We used partial dependence plots to explore how different parameter combinations change IPM strategy. Partial dependence plots show the marginal effect of a parameter of interest on IPM strategy (Friedman, 2001; Miller *et al.*, 2016). Once relevant parameter ranges were found we re-ran the genetic algorithm for those parameter combinations to generate actual IPM strategies rather than the marginal effects.

population, yield losses are never high enough to justify expenditure on control and the best strategy is to do nothing, and live with high populations of *A. myosuroides* (Fig. 2h). When  $Y_D$  increases slightly the IPM strategy shifts to a simple management regime of intermediate intensity (Fig. 2f). Once  $Y_D$  increased enough to justify intensive control, further increases did not change the IPM strategy (Fig. 2b,d).

Intensive management is costly, and so requires returns over several years to justify. As a result, valuing returns further into the future encourages more intensive management (Epanchin-Niell & Hastings, 2010). Intensive management to reduce the seed bank is only selected by the genetic algorithm when future returns are given more value (higher values of the discount rate,  $\gamma$ , Fig. 3). In agricultural systems land tenure has a crucial effect




Figure 3: The effect of discount rate ( $\gamma$ ) on the seed bank and IPM strategy (tile plots) when yields from winter wheat are high (a; £1668·ha<sup>-1</sup>) and low (b; £986·ha<sup>-1</sup>). When  $\gamma = 0.75$  rewards 5 years in the future are valued at 23% of current returns and when  $\gamma = 1$  present and future rewards are valued equally. In both cases the slope of the yield function ( $Y_D$ ) is high (£0.006·plant<sup>-1</sup>·ha<sup>-1</sup>). Initial resistance was low for both herbicides. Although we only show the first 10 years of IPM strategy, the discounted returns over 25 years are considered by the genetic algorithm.

on how investments in weed control are valued. Those who own fields can benefit from long-term investments like weed control campaigns and soil conservation, whereas those who rent fields do not (Wiese *et al.*, 1996; Fraser, 2004). The amount of rented farm land can be considerable. 54% of crop land in the USA is rented (Bigelow *et al.*, 2016), and 35% of all agricultural land in England and Wales is rented (CAAV, 2017). This has important implications for the level of control managers are incentivised to provide, and thus the spread and the evolution of resistance (Maréchal *et al.*, 2012).

Low wheat yield increases the time to pay back investments in weed control. As a result, future returns have to be valued further into the future to justify intensive control to drive down the seed bank (Fig. 3b). Thus, the effect of land tenure on weed populations at landscape scales may be more pronounced when crop yields are high, as even people with relatively low discounts rates are incentivized to control (Fig. 3a).

Resistance increased quickly with herbicide use. Even starting at low frequencies, high levels of resistance evolves after just seven herbicide applications (Fig. 4b). The best IPM



strategies are responsive to increasing resistance, drastically reducing herbicide use as higher levels of resistance evolve (Fig. 4b–d). In contrast, multiple herbicide applications per year are the norm in this cropping system, despite high levels of resistance (Hicks *et al.*, 2018). This disparity could arise from a number of contributing factors. Some managers may believe that even a little control (mortality of a few susceptible individuals) is better than no control and inaction is seen as the worst approach to weed management (Wilson *et al.*, 2008). In addition, IPM strategies are often seen as complex and more difficult to implement than the routine application of chemicals (Llewellyn *et al.*, 2006). Resistance tends to be partial and build up slowly (Moss & Hull, 2009; Hull *et al.*, 2014), so farmers may be victims of a shifting baseline (Steen & Jachowski, 2013), lowering their expectations of weed control. There may be a belief that new herbicides will become available (Hurley & Frisvold, 2016), despite no new modes of action being marketed for over 20 years (Duke, 2012). Finally, our population model was deterministic, so IPM strategies could not capture risk averse behaviour. Uncertainty in the efficacy of non-chemical control is a major impediment to adopting IPM, and farmers may prefer to stick with known chemical control, even if it is only partially effective (Hurley & Frisvold, 2016).

In our system herbicide resistance incurred a high cost, reducing gross margin (reward) by up to a quarter once high levels of resistance had evolved (Fig. 4c,d). The source of these losses depended on the initial resistance status of the population. When initial resistance was low the weed population could be initially reduced with herbicide, but then required more expensive non-chemical control to keep the weed population low so that spot control remained feasible (Fig. 4c). When resistance was initially high there were few effective control options and high weed populations reduced the yield (Fig. 4d). This result depends on the yield loss function. If high densities of the weed do not cause large yield losses then high levels of resistance are less of a concern, and high expenditure on non-chemical control is not justified.

Much work on managing resistance focuses on proactive management, delaying the initial

*de nova* evolution of resistance. In proactive management both stacking (applying different compounds at the same time) and cycling (using different compounds sequentially) can be effective (REX Consortium, 2013). In contrast we modelled reactive management, where resistant genotypes are already established in the population (Hicks *et al.*, 2018). Over a wide range of parameters, when both herbicides were effective the preference was to cycle between them (e.g. Fig. 3 and 4a). Cycling was favoured over stacking because the application of each herbicide was spread out, prolonging their useful life. We present the best case that can be hoped for in reactive management, as we assume that herbicide resistance was conferred by target site mutations. However, there is growing evidence that non-target site resistance, which frequently confers cross resistance, is widespread (Hicks *et al.*, 2018). If generalized, non-target site resistance mechanisms are present, the total amount of herbicide exposure predicts resistance level (Hicks *et al.*, 2018), and cycling is unlikely to slow the evolution of resistance.




Figure 4: The effect of initial resistance on the selected IPM strategy (tile plots) and the evolution of herbicide resistance (% survival to under both herbicides). Lighter coloured points indicate higher reward (gross margin) obtained in that time step. In this case  $Y_0 = 1668$  (high winter wheat yield) and  $Y_D = 0.0062$  (high yield penalty).

We assume that herbicide is the only action that drives the evolution of resistance. Any effective management tool will impose selection pressure, and so drive resistance to that tool (Jordan & Jannink, 1997). In reality, the spring cropping and spot control sub-actions make heavy use of glyphosate to control *A. myosuroides*. Glyphosate resistance has evolved on many separate occasions in response to prolonged, heavy use (Sammons & Gaines, 2014). As glyphosate becomes a more important part of weed management (Hicks *et al.*, 2018) resistance is likely.

## Conclusion

Combating xenobiotic resistance is ultimately a problem of behaviour change, and thus how individuals are incentivized to act (Hurley & Frisvold, 2016). Our results show that farmers have an economic incentive to be responsive to changes in the shape of the yield loss function. Doing so will require estimating, at a minimum, what yields would be in the absence of the pest, and how yields change with increasing pest density, with enough detail to say how much control (if any) is justified. The most salient piece of data on the weed population is its maximum density, as this sets the maximum possible yield loss.

[ALEXA: would be nice to conclude with a nice summary statement on what a failure to be responsive will cost farmers, the economy as whole and the environment, are at least as much as we can say at this point. Also be a good place to flag your up coming work for any things that are still unresolved. If it is all unresolved that is an important message as well I think.]

## Data Archival

Data used to fit the yield function and model code can be accessed at University of Sheffield Research Data Repository.

## Supporting Information (SI)

### Appendices

Appendix 1: Genetic Algorithm

Appendix 2: Finding Which Initial Conditions and Model Parameters Lead to Which IPM Strategies

Appendix 3: Reward Function

Appendix 4: Action Space

## Methods

We frame IPM as a combinatorial optimisation problem where the goal is to find a good combination of management tools, used in sequence. We use a genetic algorithm to solve this combinatorial problem (Taylor & Hastings, 2004; Carrasco *et al.*, 2010). Genetic algorithms cannot be checked to have found the globally optimal solution, as this would require already knowing the solution; however, genetic algorithms are efficient at selecting out comparatively poor solutions, so that over successive iterations the regions of the solution space being explored gets progressively better, resulting in a set of good (often near optimal) solutions.

Our goal is to find good IPM strategies in the face of rapidly evolving resistance, and how those strategies change in response to biological and management parameters. This problem has four parts: i) a reward function that measures how good a given IPM strategy is based on how much that strategy costs and its effectiveness, we use net present economic value; ii) a population model that translates a given IPM strategy into a population, and thus a reward; iii) an algorithm that finds IPM strategies with higher rewards, the genetic algorithm (Appendix 1); iv) finally we need to relate changes in the best IPM strategy found to changes in initial conditions and model parameters. For this we use a meta-modelling global sensitivity analysis (Coutts & Yokomizo, 2014) based on multi-variate boosted regression trees (Miller *et al.*, 2016).

### Population model

The population model links management actions to the response of the *A. myosuroides* population, and thus wheat yields. Each action  $a_j$  is a tuple of four sub-actions  $a_j =$

$\langle a_h, a_b, a_k, a_s \rangle$ . See Appendix 4 for a description of the sub-actions and all eligible combinations of these sub-actions (i.e. the full actions space,  $\mathbf{A}$ ).

The processes included in the population model limit the scope of the IPM strategies found. We use a deterministic model, and so our IPM strategies can only deal with average expected population responses, ignoring demographic uncertainty, and environmental and market variability. Also, we only model herbicide resistance that is already present in the population because *de nova* mutation is a fundamentally stochastic process.

A commonly recommended (REX Consortium, 2013) and applied (Hicks *et al.*, 2018) strategy to combat resistance is to apply xenobiotics that impair different cellular pathways (i.e. modes of action), either sequentially (cycling) or concurrently (stacking). To allow this behaviour we use a discrete time, spatially implicit model, where two independent alleles ( $R$  and  $Q$ ), each confer target site resistance to a separate herbicide. The model must also be flexible enough to accommodate non-chemical control. We include a two level seed bank (to allow plowing to take seeds out of the germinating population) and model survival as a function of resistance, herbicide choice, crop choice and spot control (where the cost increases with *A. myosuroides* density). The model tracks the number of seeds in each level of the seed bank in each of nine genotypes  $G$ , starting at the beginning of the growing season before any seeds have emerged. See Appendix 5 for a full description of the model and how each sub-action affects the population.

## Reward function

The reward function measures how good an IPM strategy is, given an initial starting condition and parameter set that the model is run under. The reward function encodes the goals of a manager. We assume farmers are primarily driven by economic returns. The economic return consists of two parts: the income made from the crop, and the costs of producing that crop. We assume that usual farm costs, such as buildings and machinery are constant from year to year, so we focus on gross margin, i.e. income - variable costs

(Redman, 2016, pp. 3–4).

To explicitly link the above ground population to the reward function we define  $N''(\mathbf{a}, n_0, t)$ , the total above ground population after all control actions, at time  $t$  given an initial population  $n_0$  and a sequence of actions

$$\mathbf{a} = \{a_j^0, a_j^1, \dots, a_j^T\} \quad (1)$$

where  $a_j^t$  is the action  $a_j \in \mathbf{A}$  taken at time  $t$  and  $T$  is the time horizon over which management is run. We assume all returns after  $T$  are ignored. The reward function is

$$R(\mathbf{a}, n_0) = \sum_{t=0}^T \gamma^t \left( Y(N''(\mathbf{a}, n_0, t)) - C(a_j^t) \right) \quad (2)$$

where  $R(\mathbf{a}, n_0)$  is the time-discounted reward for action sequence  $\mathbf{a}$  given starting population  $n_0$ ,  $\gamma \in [0, 1]$  is the discount rate. When  $\gamma = 0$  only the reward in the first time step is considered; when  $\gamma = 1$  returns in all future time steps up to  $T$  are valued equally.  $Y(N''(\mathbf{a}, n_0, t))$  is the income (in £·ha<sup>-1</sup>) from the crop chosen at time  $t$  given initial state  $n_0$  and following action sequence  $\mathbf{a}$ .  $C(a_j)$  is the cost of taking action  $a_j$ , and is composed of the cost of controlling *A. myosuroides* plus other costs that depend on the crop being grown ( $a_k$ ).

See Appendix 3 for the yield and cost models for each sub-action and parameter estimation.

## Finding good IPM strategies

Our goal is to find good strategies to manage *A. myosuroides* in the face of evolving resistance. However, it is not feasible to test every combination of management options over more than a handful of years. Genetic algorithms have been used to find good solutions to this class of problem (Taylor & Hastings, 2004; Carrasco *et al.*, 2010). The genetic algorithm starts with an randomly generated set of action sequences. These action

sequences are then iteratively improved to find a set of action sequences with a high gross margin. Genetic algorithms rely on the fact that even though the number of possible action sequences is large, many perform very poorly. The genetic algorithm explores better performing regions of the solution space more intensely. While genetic algorithms are not guaranteed to find the optimal action sequence they will find a set of actions sequences that perform well, often close to the optimal solution.

To find good action sequences we use a genetic algorithm with knock-out tournament selection, where each action sequence in a set of 1000 actions sequences is randomly paired with another, and the action sequence with the highest  $R(\mathbf{a}, n_0)$  survives to help generate new action sequences. We used pair mating between survivors and N-point cross-over to produce new action sequences. After new action sequences are created there is a process of random mutation where each  $a_j^t$  is changed to another  $a_j^t \in \mathbf{A}$  with probability  $m = 0.03$ . The algorithm used is given in Appendix 1.

## **Finding which initial conditions and model parameters lead to which IPM strategies**

It is unlikely a given IPM strategy will perform well in all scenarios. To find the parameters and initial conditions ( $n_0$ ) that shaped the IPM strategy with the highest reward, we extend the meta-modelling approach to global sensitivity analysis (outlined in (Coutts & Yokomizo, 2014)), to multivariate time series outputs (i.e. the sequences of the four sub actions). We: i) ran the genetic algorithm under 15000 different parameter sets and initial conditions, generated with Latin hyper-cube sampling (see Table S1 in Appendix 2 for upper and lower limits of each parameter); ii) used Longest Common Sub-Sequence (Toohey, 2015) as a measure of distance between these action sequences; iii) projected the resulting distance matrix into an 8D solution space using non-metric multi-dimensional scaling, implemented in the 'ecodist' R package (Goslee & Urban, 2007); and iv) predicted where each IPM solution sat in the solution space using multi-variate boosted

300 regression trees (Miller *et al.*, 2016), where the model parameters and initial conditions  
301 were predictors. See Appendix 2 for details.

302 We interrogated this multi-variate boosted regression tree to find which parameters and  
303 initial conditions were important for changing the best IPM strategy found. We used two  
304 tools, relative influence and partial dependence plots (Miller *et al.*, 2016).

## 305 References

306 Bigelow, D., Borchers, A. & Hubbs, T. (2016) *US Farmland Ownership, Tenure, and*  
307 *Transfer*. United States Department of Agriculture, Economic Research Service.

308 Bottrell, D.R. *et al.* (1979) *Integrated pest management*. United States Government Print-  
309 ing Office.

310 CAAV (2017) *Agricultural Land Occupation Survey 2016*. Central Association of Agricul-  
311 tural Valuers, UK.

312 Carrasco, L., Mumford, J., MacLeod, A., Knight, J. & Baker, R. (2010) Comprehen-  
313 sive bioeconomic modelling of multiple harmful non-indigenous species. *Ecological Eco-*  
314 *nomics*, **69**(6), 1303 – 1312. Special Section - Payments for Environmental Services:  
315 Reconciling Theory and Practice.

316 Chalak, M. & Pannell, D.J. (2015) Optimal integrated strategies to control an invasive  
317 weed. *Canadian Journal of Agricultural Economics/Revue canadienne d'agroeconomie*,  
318 **63**(3), 381–407.

319 Chesson, H.W., Kirkcaldy, R.D., Gift, T.L., Owusu-Edusei Jr, K. & Weinstock, H.S.  
320 (2018) An illustration of the potential health and economic benefits of combating  
321 antibiotic-resistant gonorrhea. *Sexually transmitted diseases*, **45**(4), 250–253.

322 Colbach, N., Busset, H., Yamada, O., Durr, C. & Caneill, J. (2006) ALOMYSYS: Mod-  
323 elling black-grass (*Alopecurus myosuroides* huds.) germination and emergence, in in-



- teraction with seed characteristics, tillage and soil climate - ii. evaluation. *European Journal of Agronomy*, **24**(2), 113–128.
- Cousens, R. (1985) A simple model relating yield loss to weed density. *Annals of applied biology*, **107**(2), 239–252.
- Coutts, S.R. & Yokomizo, H. (2014) Meta-models as a straightforward approach to the sensitivity analysis of complex models. *Population Ecology*, **56**(1), 7–19.
- Dana, E.D., Jeschke, J.M. & Garca-de Lomas, J. (2014) Decision tools for managing biological invasions: existing biases and future needs. *Oryx*, **48**(1), 5663.
- Denholm, I. & Rowland, M. (1992) Tactics for managing pesticide resistance in arthropods: theory and practice. *Annual review of entomology*, **37**(1), 91–112.
- Doyle, C.J., Cousens, R. & Moss, S.R. (1986) A model of the economics of controlling alopecurus-myosuroides huds in winter-wheat. *Crop Protection*, **5**(2), 143–150.
- Duke, S.O. (2012) Why have no new herbicide modes of action appeared in recent years? *Pest management science*, **68**(4), 505–512.
- Epanchin-Niell, R.S. & Hastings, A. (2010) Controlling established invaders: integrating economics and spread dynamics to determine optimal management. *Ecology letters*, **13**(4), 528–541.
- Fraser, E.D. (2004) Land tenure and agricultural management: soil conservation on rented and owned fields in southwest british columbia. *Agriculture and Human Values*, **21**(1), 73–79.
- Friedman, J.H. (2001) Greedy function approximation: a gradient boosting machine. *Annals of statistics*, 1189–1232.
- Goslee, S.C. & Urban, D.L. (2007) The ecodist package for dissimilarity-based analysis of ecological data. *Journal of Statistical Software*, **22**, 1–19.

- 348 Heap, I. (2014) Global perspective of herbicide-resistant weeds. *Pest Management Science*,  
349 **70**(9), 1306–1315.
- 350 Hicks, H.L., Comont, D., Coutts, S.R., Crook, L., Hull, R., Norris, K., Neve, P., Childs,  
351 D.Z. & Freckleton, R.P. (2018) The factors driving evolved herbicide resistance at a  
352 national scale. *Nature ecology & evolution*, **2**(3), 529.
- 353 Hull, R., J, C.S., L, W. & Moss, S.R. (2014) The efficacy of flufenacet based herbicides  
354 on *Alopecurus myosuroides* (blackgrass): analysis of data from 375 field trials. *Aspects*  
355 *of Applied Biology*, **127**, 49–55.
- 356 Hurley, T.M. & Frisvold, G. (2016) Economic barriers to herbicide-resistance management.  
357 *Weed Science*, **64**(sp1), 585–594.
- 358 Jordan, N. & Jannink, J. (1997) Assessing the practical importance of weed evolution: a  
359 research agenda. *Weed Research*, **37**(4), 237–246.
- 360 Laxminarayan, R., Matsoso, P., Pant, S., Brower, C., Røttingen, J.A., Klugman, K. &  
361 Davies, S. (2016) Access to effective antimicrobials: a worldwide challenge. *The Lancet*,  
362 **387**(10014), 168–175.
- 363 Livingston, M., Fernandez-Cornejo, J. & Frisvold, G.B. (2016) Economic returns to her-  
364 bicide resistance management in the short and long run: The role of neighbor effects.  
365 *Weed Science*, **64**(sp1), 595–608.
- 366 Llewellyn, R., Pannell, D., Lindner, R. & Powles, S. (2006) Targeting key perceptions  
367 when planning and evaluating extension. *Australian Journal of Experimental Agricul-*  
368 *ture*, **45**(12), 1627–1633.
- 369 Maréchal, P.Y., Henriet, F., Vancutsem, F. & Bodson, B. (2012) Ecological review of  
370 black-grass (*alopecurus myosuroides* huds.) propagation abilities in relationship with  
371 herbicide resistance. *Biotechnologie, Agronomie, Société et Environnement*, **16**(1), 103.
- 372 Miller, P.J., Lubke, G.H., McArtor, D.B. & Bergeman, C. (2016) Finding structure in  
373 data using multivariate tree boosting. *Psychological methods*, **21**(4), 583.

374 Moss, S.R. & Hull, R. (2009) The value of pre-emergence herbicides for combating  
 375 herbicide-resistant *Alopecurus myosuroides* (black-grass). *Aspects of Applied Biology*,  
 376 **91**, 79–86.

377 Moss, S.R., Perryman, S.A.M. & Tatnell, L.V. (2007) Managing herbicide-resistant black-  
 378 grass (*alopecurus myosuroides*): Theory and practice. *Weed Technology*, **21**(2), 300–  
 379 309.

380 Palumbi, S.R. (2001) Humans as the world’s greatest evolutionary force. *Science*,  
 381 **293**(5536), 1786–1790.

382 Redman, G. (2016) *John Nix Farm Managment Pocketbook*, 47th edn. Agro Business  
 383 Consultants Ltd, UK.

384 REX Consortium (2013) Heterogeneity of selection and the evolution of resistance. *Trends*  
 385 *in ecology & evolution*, **28**(2), 110–118.

386 Sammons, R.D. & Gaines, T.A. (2014) Glyphosate resistance: state of knowledge. *Pest*  
 387 *management science*, **70**(9), 1367–1377.

388 Steen, D.A. & Jachowski, D.S. (2013) Expanding shifting baseline syndrome to accom-  
 389 modate increasing abundances. *Restoration Ecology*, **21**(5), 527–529.

390 Swinton, S.M., Buhler, D.D., Forcella, F., Gunsolus, J.L. & King, R.P. (1994) Estimation  
 391 of crop yield loss due to interference by multiple weed species. *Weed Science*, **42**(1),  
 392 103–109.

393 Taylor, C.M. & Hastings, A. (2004) Finding optimal control strategies for invasive species:  
 394 a density structured model for *Spartina alterniflora*. *Journal of Applied Ecology*, **41**(6),  
 395 1049–1057.

396 Toohey, K. (2015) *SimilarityMeasures: Trajectory Similarity Measures*. URL [https://](https://CRAN.R-project.org/package=SimilarityMeasures)  
 397 [CRAN.R-project.org/package=SimilarityMeasures](https://CRAN.R-project.org/package=SimilarityMeasures). R package version 1.4.

- 398 Wiese, A.F., Salisbury, C.D., Bean, B.W., Schoenhals, M.G. & Amosson, S. (1996) Eco-  
399 nomic evaluation of field bindweed (*Convolvulus arvensis*) control in a winter wheat-  
400 fallow rotation. *Weed science*, 622–628.
- 401 Willyard, C. *et al.* (2017) Drug-resistant bacteria ranked. *Nature*, **543**(7643), 15.
- 402 Wilson, R.S., Tucker, M.A., Hooker, N.H., LeJeune, J.T. & Doohan, D. (2008) Perceptions  
403 and beliefs about weed management: perspectives of ohio grain and produce farmers.  
404 *Weed Technology*, **22**(2), 339–350.