

```
1 % Sounak Ghosh
2 % 9/28/19
3 % ECE 202 - Fall 2019 - MATLAB Exercise M5
4 % equation source: https://www.math.utah.edu/~gupta/MATH1060Fall2012/FormulaSheet.pdf
5
6 clear % clears all variables in the workplace; avoids common errors
7 clc % clears all previous outputs in the command window
8
9 %  $f(t) = 12\cos(60t + 1.8)\cos(100t + 1.2)$ 
10 %  $\cos A \cos B = 1/2 (\cos(A + B) + \cos(A - B))$ 
11
12 tms = linspace(0,200,400); % 0 to 200ms, needed for plot
13 t = tms * 10^(-3); % Converting time to milisecond
14 x = 60*t - 1.8; % first sinusoid expression in cos
15 y = 100*t + 1.2; % second sinusoid expression in cos
16 a = 12; % a is the constant
17 f1 = a*cos(x).*cos(y); % the original sinusoid function
18 f2 = (a/2)*cos(x - y); % function of the first sinusoid
19 f3 = (a/2)*cos(x + y); % function of the second sinusoid
20
21 %---- checks -----
22
23 check1 = f2 + f3 - f1; % Check function for the array with
zeroes
24 plot(tms,f1,tms,f2,tms,f3,tms,check1,'--') % plot f1, f2, f3 and check 1
25 ax = gca;
26 ax.FontSize = 18.5;
27 % legend showing the magnitude of each function
28 legend('f1 - 12*cos(x)*cos(y)', 'f2 - 6*cos(x-y)', 'f3 - 6*cos(x+y)', 'check1:f2 +
f3 - f1')
29 xlabel('time t (ms)') % x axis label
30 ylabel('f(t)') % y axis label
31 title('ECE 202, Exercise M5 - Plotting a Product of 2 Sinusoids as a Sum of 2
Sinusoids')
32 check2 = sum(abs(check1))
33
34 % The ABS(absolute function) is used to prevent the positive part cancel the
35 % the negative part. By using the ABS(absolute function) everything is
36 % becomes the same sign and reduces error points in the the plots. Instead
37 % of just adding the positive and the negative numbers which will anyway
38 % give zero, we add the positive points to just check the actual magnitude of
39 % the data. Hence, making our check more precise and solid.
40
```

```
1
2 check2 =
3
4     1.0018e-12
5
6 >>
```

### ECE 202, Exercise M5 - Plotting a Product of 2 Sinusoids as a Sum of 2 Sinusoids

