


1. $X(\omega) = \begin{cases} 1 & \omega \in A \\ -2 & \omega \in B \\ 2 & \text{otherwise, i.e. } \omega \notin A \cup B \end{cases}$ A & B are disjoint events in F .

To find cdf of X if $P(A) = \frac{1}{3}$ & $P(B) = \frac{1}{2}$



$$1 - P(A) - P(B) = 1 - \frac{1}{3} - \frac{1}{2}$$

$$= 1 - \frac{2+3}{6}$$

$$= 1 - \frac{1}{6} = \frac{5}{6} = P(X=2)$$

x	$P(X=x)$
-2	$\frac{1}{2}$
1	$\frac{1}{3}$
2	$\frac{1}{6}$ ✓

by definition of cdf $F_X(x) = P(X \leq x)$

when $x < -2$: $F_X(x) = 0$

$$-2 \leq x < 1, \quad F_X(x) = P(X = -2) = \frac{1}{2}$$

$$1 \leq x < 2, \quad F_X(x) = P(X = -2) + P(X = 1) = \frac{1}{2} + \frac{1}{3} = \frac{5}{6}$$

$$2. \quad x \geq 2 \quad F_X(x) = \frac{1}{2} + \frac{1}{3} + \frac{1}{6} = 1.$$

$$\therefore F_X(x) = \begin{cases} 0 & x < -2 \\ \frac{1}{2} & -2 \leq x < 1 \\ \frac{5}{6} & 1 \leq x < 2 \\ 1 & x \geq 2 \end{cases}$$

Ans.

2. $a < b$, $a, b \in \mathbb{R}$

(2)

Rewrite prob in form of distribution of X

(a) $P_X((a, b])$ i.e. $X > a$ and $X \leq b$

from definition of cdf $F_X(x) = P(X \leq x)$

here $P_X((a, b]) = P(a < X \leq b) = P(X \leq b) - P(X \leq a)$
 $= F_X(b) - F_X(a)$

(b) $P_X((a, b))$

So $X > a$ and $X < b$

\therefore Prob $P_X((a, b)) = P(a < X < b)$


$P(a < X < b) = P(a < X \leq b) = F_X(b) - F_X(a)$

where $F_X(b^-) = \lim_{t \rightarrow b^-} F_X(t)$ is the value of cdf

approaches as we come up to b from left

(c) $P_X([a, b]) = P(a \leq X \leq b) = F_X(b) - F_X(a^-)$

where $F_X(a^-) = \lim_{t \rightarrow a^-} F_X(t)$



Outcome $\omega = (u, v)$ is a point uniformly chosen within the square $[0,1] \times [0,1]$. The distance from (u, v) to the four edges are $\{u, 1-u, v, 1-v\}$

So $X(u, v) = \min(u, 1-u, v, 1-v)$, distance of the nearest edge.

Since by definition,

cdf $F_X(t) = P(X \leq t)$, for $t < 0$, $F_X(t) = 0$ since distances cannot be negative. For $t \geq 1/2$, $F_X(t) = 1$.

This is because max value of $X = 0.5$, at the center of side 1, the farthest any point can be from an edge

For $0 \leq t \leq 0.5$, the event $\{X > t\}$ means point is at least t away from every side

ie. $t < u < 1-t$, $t < v < 1-t$

Inner square $(t, 1-t) \times (t, 1-t)$ whose length is $1-2t$

so the area is $P(X > t) = (1-2t)^2$ for $0 \leq t \leq \frac{1}{2}$

$\therefore \forall 0 \leq t \leq 0.5$

$$F_X(t) = 1 - (1-2t)^2 = 4t - 4t^2$$

when $t < 0$, $F_X(t) = 0$

2. $t \geq 0.5$, $F_X(t) = 1$

$$\therefore \text{cdf } F_X(t) = \begin{cases} 0 & t < 0 \\ 4t - 4t^2 & 0 \leq t \leq 0.5 \\ 1 & t \geq 0.5 \end{cases}$$

Ans

④. Median of X defined as real number x_m

$$P(X \leq x_m) = 0.5$$

To find median of the r.v with pdf

$$f_X(x) = \begin{cases} be^{-x} & x > 0 \\ 0 & x < 0 \end{cases}$$

by defⁿ of prob density function, $\int_{-\infty}^{\infty} f_X(x) dx = 1$

by defⁿ here $\int_0^{\infty} be^{-x} dx = 1$

$$\text{or } b[-e^{-x}]_0^{\infty} = 1 \quad \text{or } b(0 - (-e^0)) = 1$$

$$\text{or } b = 1$$

$$\therefore \text{pdf is } F_X(x) = \int_0^x e^{-t} dt = 1 - e^{-x} \quad \forall x > 0$$

$$P(X \leq x_m) = 0.5$$

$$F_X(x_m) = 0.5$$

$$\text{or } 1 - e^{-x_m} = 0.5$$

$$e^{-x_m} = 0.5$$

$$\therefore -x_m = \ln(0.5)$$

$$\therefore x_m = -\ln(0.5) = \ln(2)$$

Answer

⑤ pdf $f_X(x) = \begin{cases} 10/x^2 & x > 10 \\ 0 & x \leq 10 \end{cases} \leftarrow \text{Given}$

Prob that a single device function for at least 15 hours

$$p = P(X > 15) = \int_{15}^{\infty} \frac{10}{x^2} dx = \left[-\frac{10}{x} \right]_{15}^{\infty} = +\frac{10}{15} = \frac{2}{3}$$

Let Y = number that last > 15

we want $P(Y > 3)$

$Y \sim \text{Binomial}(n=6, p=\frac{2}{3})$, since devices are independent (5)

$$P(Y \geq 3) = 1 - P(Y \leq 2) \\ = 1 - [P(Y=0) + P(Y=1) + P(Y=2)]$$

$$P(Y=k) = \binom{n}{k} (p)^k (1-p)^{n-k} \\ = \binom{6}{k} \left(\frac{2}{3}\right)^k \left(1 - \frac{2}{3}\right)^{6-k} = \binom{6}{k} \left(\frac{2}{3}\right)^k \left(\frac{1}{3}\right)^{6-k}$$

$$P(Y=0) = \binom{6}{0} \cdot 1 \cdot \left(\frac{1}{3}\right)^6 = 1 \cdot \frac{1}{3^6}$$

$$P(Y=1) = \binom{6}{1} \left(\frac{2}{3}\right)^1 \left(\frac{1}{3}\right)^5 = 6 \cdot \frac{2}{3} \cdot \frac{1}{3^5} = \frac{12}{3^6}$$

$$P(Y=2) = \binom{6}{2} \left(\frac{2}{3}\right)^2 \left(\frac{1}{3}\right)^4 = 15 \cdot \left(\frac{2}{3}\right)^2 \left(\frac{1}{3}\right)^4 \\ = 15 \cdot \frac{4}{3^6} = \frac{60}{3^6}$$

$$\therefore P(Y \leq 2) = \frac{1+12+60}{3^6} = \frac{73}{3^6}$$

$$\therefore P(Y \geq 3) = 1 - \frac{73}{3^6} = \frac{656}{729} \approx 0.9 \quad \underline{\text{Ans}}$$

(5) Prob density function Pdf of an exponential random variable with parameter λ

$$f_X(x) = \lambda e^{-\lambda x} \quad \forall x \geq 0$$

$$= 0, \quad \forall x < 0$$

$$F(X=x) = \int_0^x \lambda e^{-\lambda t} dt = -e^{-\lambda t} \Big|_0^x = 1 - e^{-\lambda x}$$

$$\text{cdf } F_X(x) = 1 - e^{-\lambda x}$$

(6)

we want to find two points x_1 & x_2 creating three intervals $[0, x_1)$, $[x_1, x_2)$ and $[x_2, \infty)$ so that prob of X in those interval must be $1/3$

$$P(0 \leq X \leq x_1) = \frac{1}{3}$$

$$1 - e^{-\lambda x_1} = \frac{1}{3}$$

$$\text{or } e^{-\lambda x_1} = \frac{2}{3}$$

$$\text{or } \lambda x_1 = \ln\left(\frac{3}{2}\right)$$

$$x_1 = \frac{1}{\lambda} \ln\left(\frac{3}{2}\right)$$

$$P(X \leq x_2) = \frac{2}{3}$$

$$1 - e^{-\lambda x_2} = \frac{2}{3}$$

$$\text{or } e^{-\lambda x_2} = \frac{1}{3}$$

$$\therefore x_2 = \frac{1}{\lambda} \ln(3)$$

\therefore the three intervals are

$$[0, x_1) \Rightarrow \left[0, \frac{\ln(3/2)}{\lambda}\right)$$

$$[x_1, x_2) \Rightarrow \left[\frac{\ln(3/2)}{\lambda}, \frac{\ln(3)}{\lambda}\right)$$

$$\text{and } [x_2, \infty) \Rightarrow \left[\frac{\ln(3)}{\lambda}, \infty\right)$$

each interval will prob $1/3$. Ans

(7) Discrete random variable X is geometric random variable if


$$P(X = k) = p q^{k-1} \quad \text{--- (1)} \quad k = 1, 2, 3, \dots$$

where $p, q > 0$ & $p + q = 1$.

To show, for natural numbers m & n

(7)

$$P(X > m+n \mid X > m) = P(X > n)$$



$$\begin{aligned} P(X > m+n \mid X > m) &= \frac{P(X > m+n, \overset{\text{and}}{X} > m)}{P(X > m)} \\ &= \frac{P(X > m+n)}{P(X > m)} \end{aligned}$$

$$P(X > m+n \mid X > m) = \frac{(1-p)^{m+n}}{(1-p)^m} =$$

Given ① $P(X = k) = pq^{k-1} = p(1-p)^{k-1}$

$$P(X > k) = P(X = k+1) + P(X = k+2) + \dots$$

$$= \sum_{i=k+1}^{\infty} pq^{i-1} = pq^k + pq^{k+1} + \cancel{pq^{k+2}} + \dots$$

$$= pq^k (1 + q + q^2 + \dots)$$

$$= pq^k \frac{1}{1-q} \quad \text{geometric series}$$

$$= pq^k \frac{1}{p}$$

$$= q^k$$

$$\therefore P(X > m+n \mid X > m) = \frac{q^{m+n}}{q^m} = q^n$$

$$= P(X > n)$$

$$\therefore P(X > m+n \mid X > m) = P(X > n)$$

□

7(b) Reverse statement,

8

X is +ve integer-valued random variable

$$P(X > m+n | X > m) = P(X > m) \quad \forall m, n \geq 1$$

we want to show X must be geometric.

$$\text{we know } P(X > m+n | X > m) = \frac{P(X > m+n)}{P(X > m)} = P(X > n)$$

shown earlier

$$\therefore P(X > m+n) = P(X > m) \cdot P(X > n) \quad \forall m, n \geq 1$$

— ①

let $P(X > k) = f(k)$

$$\therefore k=1, \quad f(1) = P(X > 1) \text{ say } q$$

$$k=2, \quad f(2) = P(X > 2) = f(1) \cdot f(1) = q \cdot q$$

$$k=3, \quad f(3) = P(X > 3) = f(2) \cdot f(1) = q^2 \cdot q = q^3$$

$$\therefore \text{ } f(k) = q^k$$

~~we can see~~

$$P(X > m+n) =$$

$$\begin{aligned} P(X = k) &= P(X > k-1) - P(X > k) \\ &= q^{k-1} - q^k = (1-q) q^{k-1} \\ &= p q^{k-1} \end{aligned}$$

→ geometric distribution.

eqn ① \square

(8)

X have a binomial dist

to show

$$P(X = k) = \frac{p}{1-p} \cdot \frac{n-k+1}{k} P(X = k-1)$$

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, 2, \dots, n$$

$$P(X = k-1) = \binom{n}{k-1} p^{k-1} (1-p)^{n-k+1}$$

$$\begin{aligned} \frac{P(X = k)}{P(X = k-1)} &= \frac{\binom{n}{k} \cdot p^k (1-p)^{n-k}}{\binom{n}{k-1} p^{k-1} (1-p)^{n-k+1}} \\ &= \frac{\frac{n!}{k!(n-k)!}}{\frac{n!}{(k-1)!(n-k+1)!}} \cdot p \cdot \frac{1}{(1-p)} \\ &= \frac{n! (k-1)! (n-k+1)!}{k! (n-k)! n!} \cdot p \cdot \frac{1}{(1-p)} \\ &= \frac{n-k+1}{k} \cdot p \cdot \frac{1}{(1-p)} \end{aligned}$$

$$\therefore P(X = k) = \frac{p}{(1-p)} \cdot \frac{n-k+1}{k} P(X = k-1)$$

(9)