SAMPLE GUIDELINE - SYNTHETIC TEST DATA - NOT CLINICAL ADVICE

Caring for the sick child in the community: chart booklet (sample)

Adapted for testing the guideline intake pipeline.

Contents

This page covers another part of the sick child assessment.
It is not used by the cough pathway.

Who this booklet is for

Use this booklet for a child age 2 months up to 5 years.
A younger or older child: see page 12.

Greet the caregiver

This page covers another part of the sick child assessment.
It is not used by the cough pathway.

Ask and look

ASK:
- Cough? If yes, for how long?
- Convulsions?
- Is the child not able to drink or feed anything?
- Does the child vomit everything? (vomits everything?)

LOOK:
- Chest indrawing?
- If cough, count breaths in 1 minute.
- Unusually sleepy or unconscious?

Decide: danger signs

Danger signs include:
- Cough for 14 days or more
- Diarrhoea for 14 days or more
- Convulsions, not able to drink or feed, vomits everything, chest indrawing, unusually sleepy or unconscious.

If ANY danger sign: REFER URGENTLY TO HEALTH FACILITY.

Refer urgently

If fast breathing or chest indrawing, give first dose of oral antibiotic.
Explain why child needs to go to the health facility.
Write a referral note. Help arrange transport.
FOLLOW UP child on return at least once a week until child is well.

Fast breathing

In a child age 2 months up to 12 months, 50 breaths or more per minute
In a child age 12 months up to 5 years, 40 breaths or more per minute

Cough (less than 14 days) without fast breathing: no row in this booklet.

Treat at home: Fast Breathing

Give oral amoxicillin. Give twice daily for 5 days.
Advise the caregiver to return immediately if child cannot drink or feed, becomes sicker, or has blood in stool.
Follow up child in 3 days.

Diarrhoea

This page covers another part of the sick child assessment.
It is not used by the cough pathway.

Fever

This page covers another part of the sick child assessment.
It is not used by the cough pathway.

A child outside the age range

For a child under 2 months or 5 years and older:
refer child to health facility, write a referral note, and follow up child on return.

If any OTHER PROBLEM or condition you cannot manage, refer child to health facility.