SAMPLE GUIDELINE - SYNTHETIC TEST DATA - NOT CLINICAL ADVICE

Assess and classify the sick child aged 2 months up to 5 years

Check for general danger signs:
- Not able to drink or breastfeed
- Vomits everything
- Convulsions
- Lethargic or unconscious

Then ask: Does the child have cough or difficult breathing?
If yes, the health worker should:
- Count the breaths in one minute. The child must be calm.
- Look for chest indrawing.
- Look and listen for stridor.
- Look and listen for wheezing.

Fast breathing is:
- 50 breaths per minute or more if the child is 2 months up to 12 months.
- 40 breaths per minute or more if the child is 12 months up to 5 years.

Classification table: cough or difficult breathing

Row 1. Any general danger sign or stridor in calm child.
Classify as SEVERE PNEUMONIA OR VERY SEVERE DISEASE.
Give first dose of an appropriate antibiotic. Refer URGENTLY to hospital.

Row 2. Chest indrawing or fast breathing.
Classify as PNEUMONIA.
Give oral Amoxicillin for 5 days. Advise the caregiver when to return.
Follow-up in 3 days.

Row 3. No signs of pneumonia or very severe disease.
Classify as COUGH OR COLD. Soothe the throat. Advise the caregiver when to return.

If wheezing with either fast breathing or chest indrawing: give a trial of
rapid acting inhaled bronchodilator before classifying.