

# MSBD 5007 HW2

RONG Shuo

March 30, 2025

## Question1

Determine the convexity of the following functions, where  $\mathbf{x} \in \mathbb{R}^n$  and  $\mathbf{X} \in \mathbb{S}_{++}^n$  (the set of symmetric positive definite matrices). Justify your answer.

(a)  $f(\mathbf{x}) = \log(e^{x_1} + e^{x_2} + \dots + e^{x_n})$ .

(b)  $f(\mathbf{X}) = \log \det(\mathbf{X})$ .

## Answer

(a)

$$\nabla f(\mathbf{x}) = \left( \frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$$

$$\begin{aligned} \frac{\partial f}{\partial x_i} &= \frac{1}{e^{x_1} + e^{x_2} + \dots + e^{x_n}} \times \frac{\partial}{\partial x_i} (e^{x_1} + e^{x_2} + \dots + e^{x_n}) \\ &= \frac{e^{x_i}}{e^{x_1} + e^{x_2} + \dots + e^{x_n}} \end{aligned}$$

Therefore,

$$\nabla f(\mathbf{x}) = \left( \frac{e^{x_1}}{\sum_{i=1}^n e^{x_i}}, \frac{e^{x_2}}{\sum_{i=1}^n e^{x_i}}, \dots, \frac{e^{x_n}}{\sum_{i=1}^n e^{x_i}} \right)$$

We know,

$$\begin{aligned} H_{ij} &= \frac{\partial^2 f}{\partial x_i \partial x_j} \\ H_{ii} &= \frac{\partial}{\partial x_i} \left( \frac{e^{x_i}}{S} \right) = \frac{e^{x_i}(S - e^{x_i})}{S^2} \text{ if } i = j \\ H_{ij} &= -\frac{e^{x_i} e^{x_j}}{S^2} \text{ if } i \neq j \\ \nabla^2 f(\mathbf{x}) &= \frac{1}{S} \text{diag}(\mathbf{e}) - \frac{1}{S^2} \mathbf{e} \mathbf{e}^T \end{aligned}$$

where  $\mathbf{e} = (e^{x_1}, e^{x_2}, \dots, e^{x_n})^T$ , and  $S = \sum_{i=1}^n e^{x_i}$

$$\begin{aligned} \mathbf{z} \nabla^2 f(\mathbf{x}) \mathbf{z}^T &= \frac{1}{S} \sum_{i=1}^n e^{x_i} z_i^2 - \frac{1}{S^2} \left( \sum_{i=1}^n e^{x_i} z_i \right)^2 \\ \left( \sum_{i=1}^n e^{x_i} z_i \right)^2 &\leq \left( \sum_{i=1}^n e^{x_i} z_i^2 \right) \left( \sum_{i=1}^n e^{x_i} \right) \\ \frac{1}{S} \sum_{i=1}^n e^{x_i} z_i^2 &\geq \frac{1}{S^2} \left( \sum_{i=1}^n e^{x_i} z_i \right)^2 \\ \mathbf{z} \nabla^2 f(\mathbf{x}) \mathbf{z}^T &\geq 0 \end{aligned}$$

Thus, we prove that Hessian is positive semi-definite, so we can conclude  $f(\mathbf{x})$  is convex.

(b)

Let  $\mathbf{X} \in \mathbb{S}_{++}^n$  and  $\mathbf{V} \in \mathbb{S}^n$ , we define  $g(t) = \log \det(\mathbf{X} + t\mathbf{V})$ , where  $\mathbf{X} + t\mathbf{V}$  is symmetric, positive and definite.

$$\begin{aligned} g(t) &= \log \det(\mathbf{X} + t\mathbf{V}) \\ &= \log \det(\mathbf{X}^{\frac{1}{2}}(\mathbf{I} + t\mathbf{X}^{-\frac{1}{2}}\mathbf{V}\mathbf{X}^{-\frac{1}{2}})\mathbf{X}^{\frac{1}{2}}) \\ &= \log \det \mathbf{X} + \log \det(\mathbf{I} + t\mathbf{\Lambda}) \\ &= \log \det \mathbf{X} + \sum_{i=1}^n \log(1 + t\lambda_i) \end{aligned}$$

where  $\lambda_i$  is the eigenvalues of  $\mathbf{\Lambda}$

$$g''(t) = \sum_{i=1}^n \frac{-\lambda_i^2}{(1 + t\lambda_i)^2} \leq 0$$

Thus  $g''(t) \leq 0$  for all  $t$  where  $\mathbf{X} + t\mathbf{V}$  is symmetric, positive and definite, so  $g(t)$  is concave. Since  $g(t)$  is concave for any direction,  $f(\mathbf{X}) = \log \det \mathbf{X}$  is concave on  $\mathbb{S}_{++}^n$ .

## Question2

Consider the linear system  $\mathbf{A}\mathbf{x} = \mathbf{b}$ , with  $\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 1 & 4 \end{bmatrix}$  and  $\mathbf{b} \in \begin{bmatrix} 2 \\ -4 \end{bmatrix}$ , and the initial guess  $\mathbf{x}_0 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ .

- (a) Present the first two update iterations using the **steepest descent algorithm**.
- (b) Present the first two updated iterations using the **conjugate gradient algorithm**.

## Answer

(a)

We know if we want use steepest descent algorithm to solve the linear system, we need:

$$\alpha_k = \frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k}$$

where  $\mathbf{r}_{k+1} = \mathbf{r}_k - \alpha_k \mathbf{A} \mathbf{r}_k$

$$\begin{aligned} \mathbf{r}_0 &= \mathbf{b} - \mathbf{A}\mathbf{x}_0 = \begin{bmatrix} 2 \\ -4 \end{bmatrix} \\ \alpha_0 &= \frac{5}{14} \\ \mathbf{x}_1 &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \frac{5}{14} \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \begin{bmatrix} \frac{5}{7} \\ \frac{10}{7} \end{bmatrix} \\ \mathbf{r}_1 &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} - \frac{5}{14} \begin{bmatrix} 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \end{aligned}$$

next iteration:

$$\begin{aligned} \alpha_1 &= \frac{5}{16} \\ \mathbf{x}_2 &= \begin{bmatrix} \frac{5}{7} \\ \frac{10}{7} \end{bmatrix} + \frac{5}{16} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{75}{56} \\ \frac{125}{112} \end{bmatrix} \\ \mathbf{r}_2 &= \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \frac{5}{16} \begin{bmatrix} 5 \\ 6 \end{bmatrix} = \begin{bmatrix} \frac{7}{16} \\ -\frac{14}{16} \end{bmatrix} \end{aligned}$$

In conclusion,  $\mathbf{x}_1 = \begin{bmatrix} \frac{5}{7} \\ \frac{10}{7} \end{bmatrix}$   $\mathbf{x}_2 = \begin{bmatrix} \frac{75}{56} \\ \frac{125}{112} \end{bmatrix}$

(b)

### Question3

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a twice continuously differentiable functions. Suppose that for every  $\mathbf{x} \in \mathbb{R}^n$ , the eigenvalues of the Hessian matrix  $\nabla^2 f(\mathbf{X})$  lie uniformly in the interval  $[m, M]$  with  $0 < m \leq M < \infty$ .

Prove that:

- (a) The function  $f$  has a unique global minimizer  $\mathbf{x}^*$ .
- (b) For all  $\mathbf{x} \in \mathbb{R}^n$ , the following inequality holds:

$$\frac{1}{2M} \|\nabla f(\mathbf{x})\|^2 \leq f(\mathbf{x}) - f(\mathbf{x}^*) \leq \frac{1}{2m} \|\nabla f(\mathbf{x})\|^2$$

### Question4

Consider the optimization problem  $\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x})$ , where  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is a continuously differentiable function. To develop a weighted gradient descent method, let  $\mathbf{W} \in \mathbb{R}^{n \times n}$  be a symmetric positive definite (SPD) matrix. Denote by  $\mathbf{W}^{\frac{1}{2}}$  the unique SPD square root of  $\mathbf{W}$  (i.e.,  $(\mathbf{W}^{\frac{1}{2}})^2 = \mathbf{W}$ ) and by  $\mathbf{W}^{-\frac{1}{2}}$  its inverse. Given the current iterate  $\mathbf{x}^{(k)}$ , define the next iterate  $\mathbf{x}^{(k+1)}$  as the solution of the following constrained optimization problem:

$$\begin{aligned} & \min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}^{(k)}) + \langle \nabla f(\mathbf{x}^{(k)}), \mathbf{x} - \mathbf{x}^{(k)} \rangle \\ & \text{subject to } \|\mathbf{W}^{\frac{1}{2}}(\mathbf{x} - \mathbf{x}^{(k)})\|_2 \leq \alpha_k \|\mathbf{W}^{-\frac{1}{2}} \nabla f(\mathbf{x}^{(k)})\|_2 \end{aligned}$$

where  $\alpha_k > 0$  is a step-size parameter.

Answer the following questions:

- (a) Derive an explicit formula for  $\mathbf{x}^{(k+1)}$
- (b) Prove that  $\mathbf{x}^{(k+1)}$  is equivalently the unique minimizer of the unconstrained quadratic problem:

$$\min_{\mathbf{x} \in \mathbb{R}^n} \left\{ \nabla f(\mathbf{x}^{(k)}) + \langle \nabla f(\mathbf{x}^{(k)}), \mathbf{x} - \mathbf{x}^{(k)} \rangle + \frac{1}{2\alpha_k} \|\mathbf{W}^{\frac{1}{2}}(\mathbf{x} - \mathbf{x}^{(k)})\|_2^2 \right\}$$