

# EXAMPLES OF MULTIDIMENSIONAL PROBABILITY DISTRIBUTIONS

## CHAPTER CONTENTS

Multinomial Distribution .....	61
Multivariate Normal Distribution .....	62
Dirichlet Distribution .....	63
Wishart Distribution .....	70

In this chapter, popular multidimensional probability distributions and their properties such as the expectation, the variance, and the moment-generating functions are illustrated.

## 6.1 MULTINOMIAL DISTRIBUTION

The binomial distribution explained in Section 3.2 is the probability distribution of the number  $x$  of successful trials in  $n$  Bernoulli trials with the probability of success  $p$ . The *multinomial distribution* is an extension of the binomial distribution to multidimensional cases.

Let us consider a  $d$ -sided dice with the probability of obtaining each side  $\mathbf{p} = (p_1, \dots, p_d)^\top$ , where

$$p_1, \dots, p_d \geq 0 \quad \text{and} \quad \sum_{j=1}^d p_j = 1.$$

Let  $\mathbf{x} = (x^{(1)}, \dots, x^{(d)})^\top$  be the number of times each side appears when the dice is thrown  $n$  times, where

$$\mathbf{x} \in \Delta_{d,n} = \left\{ \mathbf{x} \mid x^{(1)}, \dots, x^{(d)} \geq 0, x^{(1)} + \dots + x^{(d)} = n \right\}.$$

The probability distribution that  $\mathbf{x}$  follows is the multinomial distribution and is denoted by  $\text{Mult}(n, \mathbf{p})$ .

The probability that the  $j$ th side appears  $x^{(j)}$  times is  $(p_j)^{x^{(j)}}$ , and the number of combinations each of the  $d$  sides appears  $x^{(1)}, \dots, x^{(d)}$  times for  $n$  trials is given by

$\frac{n!}{x^{(1)}! \cdots x^{(d)}!}$ . Putting them together, the probability mass function of  $\text{Mult}(n, \mathbf{p})$  is given by

$$f(\mathbf{x}) = \frac{n!}{x^{(1)}! \cdots x^{(d)}!} (p_1)^{x^{(1)}} \cdots (p_d)^{x^{(d)}}.$$

When  $d = 2$ ,  $\text{Mult}(n, \mathbf{p})$  is reduced to  $\text{Bi}(n, p_1)$ .

The binomial theorem can also be extended to *multinomial theorem*:

$$(p_1 + \cdots + p_d)^n = \sum_{\mathbf{x} \in \Delta_{d,n}} \frac{n!}{x^{(1)}! \cdots x^{(d)}!} (p_1)^{x^{(1)}} \cdots (p_d)^{x^{(d)}},$$

with which the moment-generating function of  $\text{Mult}(n, \mathbf{p})$  can be computed as

$$\begin{aligned} M_{\mathbf{x}}(\mathbf{t}) &= E[e^{\mathbf{t}^\top \mathbf{x}}] = \sum_{\mathbf{x} \in \Delta_{d,n}} e^{t_1 x^{(1)}} \cdots e^{t_d x^{(d)}} \frac{n!}{x^{(1)}! \cdots x^{(d)}!} (p_1)^{x^{(1)}} \cdots (p_d)^{x^{(d)}} \\ &= \sum_{\mathbf{x} \in \Delta_{d,n}} \frac{n!}{x^{(1)}! \cdots x^{(d)}!} (p_1 e^{t_1})^{x^{(1)}} \cdots (p_d e^{t_d})^{x^{(d)}} \\ &= (p_1 e^{t_1} + \cdots + p_d e^{t_d})^n. \end{aligned}$$

From this, the expectation of and (co)variance of  $\text{Mult}(n, \mathbf{p})$  can be computed as

$$E[x^{(j)}] = np_j \quad \text{and} \quad \text{Cov}[x^{(j)}, x^{(j')}] = \begin{cases} np_j(1 - p_j) & (j = j'), \\ -np_j p_{j'} & (j \neq j'). \end{cases}$$

## 6.2 MULTIVARIATE NORMAL DISTRIBUTION

When random variables  $y^{(1)}, \dots, y^{(d)}$  independently follow the standard normal distribution  $N(0, 1)$ , the joint probability density function of  $\mathbf{y} = (y^{(1)}, \dots, y^{(d)})^\top$  is given by

$$\begin{aligned} g(\mathbf{y}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(y^{(j)})^2}{2}\right) \\ &= \frac{1}{(2\pi)^{d/2}} \exp\left(-\frac{1}{2} \mathbf{y}^\top \mathbf{y}\right). \end{aligned}$$

The expectation and variance-covariance matrices of  $\mathbf{y}$  are given by

$$E[\mathbf{y}] = \mathbf{0} \quad \text{and} \quad V[\mathbf{y}] = \mathbf{I},$$

where  $\mathbf{0}$  denotes the zero vector and  $\mathbf{I}$  denotes the identity matrix.

Let us transform  $\mathbf{y}$  by  $d \times d$  invertible matrix  $\mathbf{T}$  and  $d$ -dimensional vector  $\boldsymbol{\mu}$  as

$$\mathbf{x} = \mathbf{T}\mathbf{y} + \boldsymbol{\mu}.$$

Then the joint probability density function of  $\mathbf{x}$  is given by

$$\begin{aligned} f(\mathbf{x}) &= g(\mathbf{y})|\det(\mathbf{T})|^{-1} \\ &= \frac{1}{(2\pi)^{d/2}\det(\boldsymbol{\Sigma})^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right), \end{aligned}$$

where  $\det(\mathbf{T})$  is the Jacobian (Fig. 4.2), and

$$\boldsymbol{\Sigma} = \mathbf{T}\mathbf{T}^\top.$$

This is the general form of the *multivariate normal distribution* and is denoted by  $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ . The expectation and variance-covariance matrices of  $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  are given by

$$\begin{aligned} E[\mathbf{x}] &= \mathbf{T}E[\mathbf{y}] + \boldsymbol{\mu} = \boldsymbol{\mu}, \\ V[\mathbf{x}] &= V[\mathbf{T}\mathbf{y} + \boldsymbol{\mu}] = \mathbf{T}V[\mathbf{y}]\mathbf{T}^\top = \boldsymbol{\Sigma}. \end{aligned}$$

Probability density functions of two-dimensional normal distribution  $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  are illustrated in Fig. 6.1. The contour lines of two-dimensional normal distributions are *elliptic*, and their *principal axes* agree with the coordinate axes if the variance-covariance matrix is diagonal. Furthermore, the elliptic contour lines become spherical if all diagonal elements are equal (i.e., the variance-covariance matrix is proportional to the identity matrix).

Eigendecomposition of variance-covariance matrix  $\boldsymbol{\Sigma}$  (see Fig. 6.2) shows that the principal axes of the ellipse are parallel to the eigenvectors of  $\boldsymbol{\Sigma}$ , and their length is proportional to the square root of the eigenvalues (Fig. 6.3).

## 6.3 DIRICHLET DISTRIBUTION




Let  $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_d)^\top$  be a  $d$ -dimensional vector with positive entries and let  $y^{(1)}, \dots, y^{(d)}$  be random variables that independently follow the gamma distribution  $\text{Ga}(\alpha_j, \lambda)$ . For  $V = \sum_{j=1}^d y^{(j)}$ , let

$$\mathbf{x} = (x^{(1)}, \dots, x^{(d)})^\top = \left( \frac{y^{(1)}}{V}, \dots, \frac{y^{(d)}}{V} \right)^\top.$$

Then the distribution that the above  $d$ -dimensional vector  $\mathbf{x}$  follows is called the *Dirichlet distribution* and is denoted by  $\text{Dir}(\boldsymbol{\alpha})$ . The domain of  $\mathbf{x}$  is given by

$$\Delta_d = \left\{ \mathbf{x} \mid x^{(1)}, \dots, x^{(d)} \geq 0, x^{(1)} + \dots + x^{(d)} = 1 \right\}.$$

Drawing a value from a Dirichlet distribution corresponds to generating a (unfair)  $d$ -sided die.

(a) When  $\Sigma$  is generic, the contour lines are elliptic(b) When  $\Sigma$  is diagonal, the principal axes of the elliptic contour lines agree with the coordinate axes(c) When  $\Sigma$  is proportional to identity, the contour lines are spherical**FIGURE 6.1**

Probability density functions of two-dimensional normal distribution  $N(\mu, \Sigma)$  with  $\mu = (0, 0)^\top$ .

For  $d \times d$  matrix  $\mathbf{A}$ , a nonzero vector  $\boldsymbol{\phi}$  and a scalar  $\lambda$  that satisfy

$$\mathbf{A}\boldsymbol{\phi} = \lambda\boldsymbol{\phi}$$

are called an *eigenvector* and an *eigenvalue* of  $\mathbf{A}$ , respectively. Generally, there exist  $d$  eigenvalues  $\lambda_1, \dots, \lambda_d$  and they are all real when  $\mathbf{A}$  is symmetric. A matrix whose eigenvalues are all positive is called a *positive definite matrix*, and a matrix whose eigenvalues are all non-negative is called a *positive semidefinite matrix*. Eigenvectors  $\boldsymbol{\phi}_1, \dots, \boldsymbol{\phi}_d$  corresponding to eigenvalues  $\lambda_1, \dots, \lambda_d$  are orthogonal and are usually normalized, i.e., they are *orthonormal* as

$$\boldsymbol{\phi}_j^\top \boldsymbol{\phi}_{j'} = \begin{cases} 1 & (j = j'), \\ 0 & (j \neq j'). \end{cases}$$

A matrix  $\mathbf{A}$  can be expressed by using its eigenvectors and eigenvalues as

$$\mathbf{A} = \sum_{j=1}^d \lambda_j \boldsymbol{\phi}_j \boldsymbol{\phi}_j^\top = \boldsymbol{\Phi} \boldsymbol{\Lambda} \boldsymbol{\Phi}^\top,$$

where  $\boldsymbol{\Phi} = (\boldsymbol{\phi}_1, \dots, \boldsymbol{\phi}_d)$  and  $\boldsymbol{\Lambda}$  is the diagonal matrix with diagonal elements  $\lambda_1, \dots, \lambda_d$ . This is called *eigenvalue decomposition* of  $\mathbf{A}$ . In MATLAB, eigenvalue decomposition can be performed by the `eig` function. When all eigenvalues are nonzero, the inverse of  $\mathbf{A}$  can be expressed as

$$\mathbf{A}^{-1} = \sum_{j=1}^d \lambda_j^{-1} \boldsymbol{\phi}_j \boldsymbol{\phi}_j^\top = \boldsymbol{\Phi} \boldsymbol{\Lambda}^{-1} \boldsymbol{\Phi}^\top.$$


For  $d \times d$  matrix  $\mathbf{A}$  and  $d \times d$  positive symmetric matrix  $\mathbf{B}$ , a nonzero vector  $\boldsymbol{\phi}$  and a scalar  $\lambda$  that satisfy

$$\mathbf{A}\boldsymbol{\phi} = \lambda\mathbf{B}\boldsymbol{\phi}$$

are called a *generalized eigenvector* and a *generalized eigenvalue* of  $\mathbf{A}$ , respectively.

## FIGURE 6.2

Eigenvalue decomposition.

**FIGURE 6.3**

Contour lines of the normal density. The principal axes of the ellipse are parallel to the eigenvectors of variance-covariance matrix  $\Sigma$ , and their length is proportional to the square root of the eigenvalues.

The probability density function of  $\text{Dir}(\alpha)$  is given by

$$f(\mathbf{x}) = \frac{\prod_{j=1}^d (x^{(j)})^{\alpha_j - 1}}{B_d(\alpha)},$$

where

$$B_d(\alpha) = \int_{\Delta_d} \prod_{j=1}^d (x^{(j)})^{\alpha_j - 1} d\mathbf{x} \quad (6.1)$$

is the  $d$ -dimensional *beta function*.

For  $x = (\sqrt{p} \sin \theta)^2$  with  $0 \leq p \leq 1$ , Eq. (4.6) implies

$$\begin{aligned} & \int_0^p x^{\alpha-1} (p-x)^{\beta-1} dx \\ &= \int_0^{\frac{\pi}{2}} (\sqrt{p} \sin \theta)^{2(\alpha-1)} \left( p - (\sqrt{p} \sin \theta)^2 \right)^{\beta-1} \frac{dx}{d\theta} d\theta \\ &= \int_0^{\frac{\pi}{2}} p^{\alpha-1} (\sin \theta)^{2(\alpha-1)} p^{\beta-1} (\cos \theta)^{2(\beta-1)} \cdot 2p \sin \theta \cos \theta d\theta \\ &= p^{\alpha+\beta-1} \left( 2 \int_0^{\frac{\pi}{2}} (\sin \theta)^{2\alpha-1} (\cos \theta)^{2\beta-1} d\theta \right) \\ &= p^{\alpha+\beta-1} B(\alpha, \beta). \end{aligned}$$

Letting  $p = 1 - \sum_{j=1}^{d-2} x^{(j)}$ , the integration with respect to  $x^{(d-1)}$  in

$$\begin{aligned} B_d(\alpha) &= \int_0^1 (x^{(1)})^{\alpha_1-1} \times \int_0^{1-x^{(1)}} (x^{(2)})^{\alpha_2-1} \times \\ &\quad \dots \times \int_0^{1-\sum_{j=1}^{d-2} x^{(j)}} (x^{(d-1)})^{\alpha_{d-1}-1} \\ &\quad \times \left(1 - \sum_{j=1}^{d-1} x^{(j)}\right)^{\alpha_d-1} dx^{(1)} dx^{(2)} \dots dx^{(d-1)} \end{aligned}$$

can be computed as

$$\begin{aligned} &\int_0^{1-\sum_{j=1}^{d-2} x^{(j)}} (x^{(d-1)})^{\alpha_{d-1}-1} \left(1 - \sum_{j=1}^{d-1} x^{(j)}\right)^{\alpha_d-1} dx^{(d-1)} \\ &= \left(1 - \sum_{j=1}^{d-2} x^{(j)}\right)^{\sum_{j=d-1}^d \alpha_j - 1} B(\alpha_{d-1}, \alpha_d). \end{aligned}$$

Similarly, letting  $p = 1 - \sum_{j=1}^{d-3} x^{(j)}$ , the integration with respect to  $x^{(d-2)}$  in the above equation can be computed as

$$\begin{aligned} &\int_0^{1-\sum_{j=1}^{d-3} x^{(j)}} (x^{(d-2)})^{\alpha_{d-2}-1} \left(1 - \sum_{j=1}^{d-2} x^{(j)}\right)^{\sum_{j=d-1}^d \alpha_j - 1} dx^{(d-2)} \\ &= \left(1 - \sum_{j=1}^{d-3} x^{(j)}\right)^{\sum_{j=d-2}^d \alpha_j - 1} B\left(\alpha_{d-2}, \sum_{j=d-1}^d \alpha_j\right). \end{aligned}$$

Repeating this computation yields

$$B_d(\alpha) = \prod_{j=1}^{d-1} B\left(\alpha_j, \sum_{j'=j+1}^d \alpha_{j'}\right).$$

Applying

$$B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)}$$

given in Eq. (4.7) to the above equation, the  $d$ -dimensional beta function can be expressed by using the gamma function as

$$B_d(\alpha) = \prod_{j=1}^{d-1} \frac{\Gamma(\alpha_j) \Gamma\left(\sum_{j'=j+1}^d \alpha_{j'}\right)}{\Gamma\left(\sum_{j'=j}^d \alpha_{j'}\right)} = \frac{\prod_{j=1}^d \Gamma(\alpha_j)}{\Gamma(\alpha_0)},$$

where

$$\alpha_0 = \sum_{j=1}^d \alpha_j.$$

When  $d = 2$ ,  $x^{(2)} = 1 - x^{(1)}$  holds and thus the Dirichlet distribution is reduced to the beta distribution shown in Section 4.4:

$$f(x) = \frac{x^{\alpha_1-1}(1-x)^{\alpha_2-1}}{B(\alpha_1, \alpha_2)}.$$

Thus, the Dirichlet distribution can be regarded as a multidimensional extension of the beta distribution.

The expectation of  $\text{Dir}(\alpha)$  is given by

$$\begin{aligned} E[x^{(j)}] &= \frac{\int x^{(j)} \prod_{j'=1}^d (x^{(j')})^{\alpha_{j'}-1} dx^{(j)}}{B_d(\alpha)} \\ &= \frac{B_d(\alpha_1, \dots, \alpha_{j-1}, \alpha_j + 1, \alpha_{j+1}, \dots, \alpha_d)}{B_d(\alpha_1, \dots, \alpha_{j-1}, \alpha_j, \alpha_{j+1}, \dots, \alpha_d)} \\ &= \frac{\Gamma(\alpha_1) \cdots \Gamma(\alpha_{j-1}) \Gamma(\alpha_j + 1) \Gamma(\alpha_{j+1}) \cdots \Gamma(\alpha_d) \times \Gamma(\alpha_0)}{\Gamma(\alpha_0 + 1) \times \Gamma(\alpha_1) \cdots \Gamma(\alpha_{j-1}) \Gamma(\alpha_j) \Gamma(\alpha_{j+1}) \cdots \Gamma(\alpha_d)} \\ &= \frac{\Gamma(\alpha_j + 1) \Gamma(\alpha_0)}{\Gamma(\alpha_0 + 1) \Gamma(\alpha_j)} = \frac{\alpha_j \Gamma(\alpha_j) \Gamma(\alpha_0)}{\alpha_0 \Gamma(\alpha_0) \Gamma(\alpha_j)} \\ &= \frac{\alpha_j}{\alpha_0}. \end{aligned}$$

The variance and covariance of  $\text{Dir}(\alpha)$  can also be obtained similarly as


$$\text{Cov}[x^{(j)}, x^{(j')}] = \begin{cases} \frac{\alpha_j(\alpha_0 - \alpha_j)}{\alpha_0^2(\alpha_0 + 1)} & (j = j'), \\ -\frac{\alpha_j \alpha_{j'}}{\alpha_0^2(\alpha_0 + 1)} & (j \neq j'). \end{cases}$$

Probability density functions of  $\text{Dir}(\alpha)$  when  $d = 3$  are illustrated in Fig. 6.4. When  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha$ , the probability density takes larger values around the corners of the triangle if each  $\alpha < 1$ , the probability density is uniform if each  $\alpha = 1$ , and the probability density takes larger values around the center if each  $\alpha > 1$ . When  $\alpha_1, \alpha_2, \alpha_3$  are nonuniform, the probability density tends to take large values around the corners of the triangle with large  $\alpha_j$ .

When the Dirichlet parameters  $\alpha$  are all equal, i.e.,

$$\alpha_1 = \cdots = \alpha_d = \alpha,$$



**FIGURE 6.4**

Probability density functions of Dirichlet distribution  $\text{Dir}(\alpha)$ . The center of gravity of the triangle corresponds to  $x^{(1)} = x^{(2)} = x^{(3)} = 1/3$ , and each vertex represents the point that the corresponding variable takes one and the others take zeros.

the probability density function is simplified as

$$f(\mathbf{x}) = \frac{\Gamma(\alpha d)}{\Gamma(\alpha)^d} \prod_{j=1}^d (x^{(j)})^{\alpha-1}.$$

This is called *symmetric Dirichlet distribution* and is denoted by  $\text{Dir}(\alpha)$ . The common parameter  $\alpha$  is often referred to as the *concentration parameter*.

## 6.4 WISHART DISTRIBUTION

Let  $\mathbf{x}_1, \dots, \mathbf{x}_n$  be  $d$ -dimensional random variables independently following the normal distribution  $N(\mathbf{0}, \Sigma)$ , and let

$$\mathbf{S} = \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top$$

be the *scatter matrix* which is assumed to be invertible. The probability distribution that  $\mathbf{S}$  follows is called the *Wishart distribution* with  $n$  degrees of freedom and is denoted by  $W(\Sigma, d, n)$ . The probability density function of  $W(\Sigma, d, n)$  is given by

$$f(\mathbf{S}) = \frac{\det(\mathbf{S})^{\frac{n-d-1}{2}} \exp\left(-\frac{1}{2} \text{tr}(\Sigma^{-1} \mathbf{S})\right)}{\det(2\Sigma)^{\frac{n}{2}} \Gamma_d\left(\frac{n}{2}\right)},$$

where  $\det(\cdot)$  denotes the determinant of a matrix,  $\text{tr}(\cdot)$  denotes the trace of a matrix,  $\Gamma_d(\cdot)$  denotes the  $d$ -dimensional *gamma function* defined by

$$\Gamma_d(a) = \int_{\mathbf{S} \in \mathbb{S}_d^+} \det(\mathbf{S})^{a-\frac{d+1}{2}} \exp(-\text{tr}(\mathbf{S})) d\mathbf{S}, \quad (6.2)$$

and  $\mathbb{S}_d^+$  denotes the set of all  $d \times d$  positive symmetric matrices.

When  $\Sigma = \frac{1}{2} \mathbf{I}$ , the definition of  $d$ -dimensional gamma function immediately yields

$$\int_{\mathbf{S} \in \mathbb{S}_d^+} f(\mathbf{S}) d\mathbf{S} = 1. \quad (6.3)$$

Since the Jacobian is given by  $\det(2\Sigma)^{\frac{d+1}{2}}$  when  $\mathbf{S}$  is transformed to  $\frac{1}{2}\Sigma^{-1}\mathbf{S}$  (Fig. 4.2), Eq. (6.3) is also satisfied for generic  $\Sigma \in \mathbb{S}_d^+$ .

$d$ -dimensional gamma function  $\Gamma_d(\cdot)$  can be expressed by using two-dimensional gamma function  $\Gamma(\cdot)$  as

$$\begin{aligned} \Gamma_d(a) &= \pi^{\frac{d(d-1)}{4}} \prod_{j=1}^d \Gamma\left(a + \frac{1-j}{2}\right) \\ &= \pi^{\frac{d-1}{2}} \Gamma_{d-1}(a) \Gamma\left(a + \frac{1-d}{2}\right). \end{aligned}$$

An operator that transforms an  $m \times n$  matrix  $\mathbf{A} = (\mathbf{a}_1, \dots, \mathbf{a}_n)$  to the  $mn$ -dimensional vector,

$$\text{vec}(\mathbf{A}) = (\mathbf{a}_1^\top, \dots, \mathbf{a}_n^\top)^\top,$$

is called the *vectorization operator*. The operator that transforms an  $m \times n$  matrix  $\mathbf{A}$  and a  $p \times q$  matrix  $\mathbf{B}$  to an  $mp \times nq$  matrix as

$$\mathbf{A} \otimes \mathbf{B} = \begin{pmatrix} a_{1,1}\mathbf{B} & \cdots & a_{1,n}\mathbf{B} \\ \vdots & \ddots & \vdots \\ a_{m,1}\mathbf{B} & \cdots & a_{m,n}\mathbf{B} \end{pmatrix}$$

is called the *Kronecker product*. The vectorization operator and the Kronecker product satisfy the following properties:

$$\begin{aligned} \text{vec}(\mathbf{ABC}) &= (\mathbf{C}^\top \otimes \mathbf{A})\text{vec}(\mathbf{B}) \\ &= (\mathbf{I} \otimes \mathbf{AB})\text{vec}(\mathbf{C}) \\ &= (\mathbf{C}^\top \mathbf{B}^\top \otimes \mathbf{I})\text{vec}(\mathbf{A}), \\ (\mathbf{A} \otimes \mathbf{B})^{-1} &= \mathbf{A}^{-1} \otimes \mathbf{B}^{-1}, \\ (\mathbf{A} \otimes \mathbf{B})(\mathbf{C} \otimes \mathbf{D}) &= (\mathbf{AC} \otimes \mathbf{BD}), \\ \text{tr}(\mathbf{A} \otimes \mathbf{B}) &= \text{tr}(\mathbf{A})\text{tr}(\mathbf{B}), \\ \text{tr}(\mathbf{AB}) &= \text{vec}(\mathbf{A}^\top)^\top \text{vec}(\mathbf{B}), \\ \text{tr}(\mathbf{ABCD}) &= \text{vec}(\mathbf{A}^\top)^\top (\mathbf{D}^\top \otimes \mathbf{B})\text{vec}(\mathbf{C}). \end{aligned}$$

These formulas allow us to compute the product of big matrices efficiently.

### FIGURE 6.5

Vectorization operator and Kronecker product.

Thus, the Wishart distribution can be regarded as a multidimensional extension of the gamma distribution. When  $d = 1$ ,  $\mathbf{S}$  and  $\mathbf{\Sigma}$  become scalars, and letting  $S = x$  and  $\Sigma = 1$  yields

$$f(x) = \frac{x^{\frac{n}{2}-1} \exp\left(-\frac{x}{2}\right)}{2^{\frac{n}{2}} \Gamma\left(\frac{n}{2}\right)}.$$

This is equivalent to the probability density function of the chi-squared distribution with  $n$  degrees of freedom explained in Section 4.3.

The moment-generating function of  $W(\Sigma, d, n)$  can be obtained by transforming  $S$  to  $(\frac{1}{2}\Sigma^{-1} - T)S$  as

$$\begin{aligned} M_S(T) &= E[e^{\text{tr}(TS)}] \\ &= \int_{S \in \mathbb{S}_d^+} \frac{\det(S)^{\frac{n-d-1}{2}} \exp(-\text{tr}((\frac{1}{2}\Sigma^{-1} - T)S))}{2^{\frac{dn}{2}} \det(\Sigma)^{\frac{n}{2}} \Gamma_d(\frac{n}{2})} dS \\ &= \det(I - 2T\Sigma)^{-\frac{n}{2}}. \end{aligned}$$

The expectation and variance-covariance matrices of  $S$  that follows  $W(\Sigma, d, n)$  are given by

$$E[S] = n\Sigma \quad \text{and} \quad V[\text{vec}(S)] = 2n\Sigma \otimes \Sigma,$$

where  $\text{vec}(\cdot)$  is the vectorization operator and  $\otimes$  denotes the *Kronecker product* (see Fig. 6.5).