Can Consistent Forces Recover Molecular Shapes?

Shedrack Eze (seze) **Olufunke Abodunrin (oabodunr)**
Emmanuel Aaron (eaaron) **Muhammed Idris (midris)**
Carnegie Mellon University Group 9, Path 3

Abstract

A molecule's shape is the three-dimensional arrangement of its atoms, the tiny pieces of matter; forces push and pull those atoms. Learned forces can guide a search for stable shapes, but accurate predictions on stored examples do not guarantee the correct destination. An energy baseline, our comparison model, calculates forces from negative slopes of a predicted energy number, while a direct model predicts force arrows themselves. Consistency training penalizes disagreement between direct predictions and forces calculated from energy.

On aspirin, we compare force errors and measured computing cost using validation examples reserved for model selection. Direct and consistency models meet our error-matching rule at one consistency strength, but all five tested models recover zero of ten targets. This failure includes the energy baseline, even though all target shapes appear in training, so the pilot cannot rank successful recovery. Inaccurate forces at reference shapes and excessive coordinate steps leave the recovery question open.

1 Why this problem matters

Training changes a model's adjustable numbers, called parameters, to approximate expensive physics calculations from reference examples.

Energy is a number assigned to an atomic arrangement. A stretched spring gives a useful picture: stretching it further stores more energy, and its pull points toward lower energy. Energy slope means how quickly energy changes when an atom moves; the energy baseline calculates forces from the negative slopes. The direct model predicts force arrows themselves, while consistency training penalizes disagreement with forces calculated from negative energy slopes.

Our objective is correct-shape recovery at comparable force error, with computing cost measured separately; success requires both the assigned target shape and the force stopping rule.

2 What earlier studies establish

Landscape17 supplies molecular shapes, reference forces and paths between stable arrangements [1]. It motivates checking final shapes beyond prediction errors, while Bigi et al. describe problems when direct forces do not follow one energy function [2]. Together, these studies motivate separating a force-based stop from structural success.

E3x provides networks with three-dimensional symmetry rules [3], while MACE supplies an established energy-derived reference [4]. Earlier work already studies physical constraints in training losses [5]. Our contribution compares error, cost and recovery under shared conditions; consistency training itself is not new.

11-785 Introduction to Deep Learning, Fall 2026. Group 9 midterm report, 23 October 2026.

3 Data and a fair comparison

We use aspirin from Landscape17 [1], with 21 atoms in each molecule. Reference energies and forces come from hybrid density-functional theory, a detailed calculation based on electron physics, so these reference labels are calculated answers rather than exact experimental truth.

A transition state is a high-energy arrangement with two downhill branches toward stable arrangements, called minima. A connection group contains both branches' records, which stay in the same partition to avoid separating the two sides of one transition state. Training adjusts parameters, validation selects settings and saved models, and the test partition is reserved for later trained-model evaluation.

Training/validation/test contain 26/5/6 connections and 4,891/974/1,127 records. Training-only normalization puts energy and force errors on comparable scales, while shared minima mean that records are not independent examples of new destinations. Each branch targets the minimum it should reach; empirical classifications corrected 26 of 74 targets previously assigned from filename ordering. Earlier model-free software checks parsed test coordinates and labels but did not choose models or solver settings; no trained model has been scored on the test partition.

3.1 Inputs, outputs and the three model types

Our E3x graph neural networks link nearby atoms, called nodes; layers, or successive calculation stages, pass learned information along these links. Inputs are atom types and positions; outputs are molecular energy and per-atom force arrows. All types use 64 learned number/vector channels, two interactions and eight radial functions describing pair distances.

Links extend to 5 Å, where an angstrom (Å) is 10^{-10} metres. Internal features carry scalar numbers and vectors with direction. Rotating a molecule rotates its predicted forces without changing its energy, properties called equivariance for forces and invariance for energy. The energy baseline has 43,521 parameters; models with separate energy and force outputs have 43,649.

The energy baseline calculates $\widehat{\mathbf{F}}^E = -\nabla_{\mathbf{R}}\widehat{E}$. Here, $\widehat{E}$ is predicted energy, $\mathbf{R}$ contains atom positions, and the gradient gives energy slopes along position coordinates. Both the direct and consistency models also predict energy trained against reference labels, so they are not force-only models. Ordinary direct training has no agreement penalty, while consistency training adds one.

3.2 Training loss and measured success

The loss combines normalized squared energy error, normalized squared force error and, where applicable, force–energy disagreement; training adjusts parameters to reduce this number:

$$\mathcal{L} = w_E \,\mathrm{mean}(e_E^2) + w_F \,\mathrm{mean}(e_F^2) + \lambda \,\mathrm{mean}\left[\left((\widehat{\mathbf{F}}^D - \widehat{\mathbf{F}}^E)/s_F\right)^2\right]. \tag{1}$$

Here, e_E and e_F are errors divided by training-derived scales; s_F is the force scale, and superscript D means direct prediction. Weights $w_E = 0.014584365670390719$ and $w_F = 1.9854156343296092$ balance supervised errors. The energy baseline omits the final term, while direct training uses $\lambda = 0$ and consistency training uses a positive λ, the consistency strength.

Adam updates parameters at learning rate 0.003 using 64-bit arithmetic. Each of 200,000 updates samples four records that can repeat, giving 800,000 exposures. Two seeds specify random starting parameters and sampling, providing a limited repeatability check.

Validation occurs every 500 updates, including zero, and BEST is the first saved model state, or checkpoint, with minimum validation force error.

For each validation connection, force mean absolute error (MAE) averages the absolute prediction error across every record, every atom and all three coordinate directions. The reported score averages these five connection scores equally; smaller values are better.

Scores use meV/Å: a millielectronvolt is one thousandth of an electronvolt, an energy unit, and energy divided by distance measures force.

Recovery additionally requires maximum absolute predicted force component ≤ 0.01 eV/Å and aligned structural RMSD ≤ 0.05 Å. Root-mean-square deviation (RMSD) measures typical atom-position disagreement after rotation, translation and allowed atom-symmetry matching. Small predicted force is a stopping condition; it is not small prediction error.

4 What we trained and learned

4.1 A fitting plateau changed the training recipe

Early small-subset E3x training plateaued, but MACE, which also derives forces from energy, fitted the same 128 training records to 0.001515 of the zero-force squared error. Different settings prevent a controlled comparison, but the plateau did not establish an unavoidable data or energy-derived-force limit.

Later 1,000-update energy and direct fits selected update zero as BEST: training improved energy but worsened force predictions.

We then completed three 200,000-update energy baselines (Figure 1). A used width 32 and learning rate 0.001; B used width 64 and 0.003. Both used equal normalized energy/force weights, while C retained B's structure and rate but used the balanced weights in Section 3.2. Width counts internal feature channels.




Figure 1: Energy-baseline validation, seed 0. Markers show BEST; the dashed line is the declared 50 meV/Å adequacy rule. Each point uses all 974 validation records, with equal connection weighting.

C reached 34.587 meV/Å, versus 40.370 for B and 42.108 for A, and passed the adequacy rule. Predicting zero force gives 62.504 meV/Å, so C reduces that error by 44.664%. However, C's energy error is higher than B's: 126.990 versus 28.178 meV per molecule. We selected C on the energy baseline and transferred its recipe without retuning the other model types.

4.2 Agreement improves, but recovery still needs a separate test

We reused C seed zero and trained nine more fits, giving two seeds per row in Table 1. Consistency strength 0.1 reduces normalized force–energy disagreement squared error to 0.49–0.61, versus 116–379 for direct training. Its measured fit-plus-score time is 1.985 times direct prediction, excluding other diagnostics and overhead. This is not inference speed.

Table 1: Validation error at BEST. Force columns use meV/Å; energy uses meV per molecule. Seed limits show two observed values, not confidence bounds.

Model	Mean force MAE	Seed min–max	Mean energy MAE
Energy-derived	34.893	34.587–35.198	128.725
Direct	38.186	37.291–39.082	76.684
Consistency $\lambda = 0.1$	38.024	36.773–39.275	87.006
Consistency $\lambda = 1$	34.683	33.723–35.643	112.341
Consistency $\lambda = 10$	39.725	39.060–40.390	161.662

Only strength 0.1 meets our declared matching rule against direct prediction: the mean-error gap must be within 10% of the smaller mean and within both seed spans. A span is maximum minus minimum.

BEST is optimistic because it selects among 401 checks. Over updates 150,000–200,000, the last 50,000 updates, baseline errors vary by 2.2–3.1 meV/Å standard deviation across correlated checks. Nine subsequent fits finish 0.1–7.4 meV/Å above BEST, and the 0.162 meV/Å matching gap is smaller than this variation. Matching does not establish equivalence.

MACE Run A independently scores 5.349 meV/Å with equal connection weighting after 2,048 epochs, or training-data passes. Architecture, exposure and hardware differ; this is an external reference, not a controlled efficiency comparison. MACE recovery remains unmeasured.

5 The shape-search pilot fails for every evaluated checkpoint

A relaxation procedure repeatedly moves atoms along predicted forces, and we tested five saved E3x checkpoints from the same ten branch starts across five validation connections. Each branch has its own target, and all six distinct target minima appear in training. Leaving the starts unchanged recovers zero targets, while every evaluated checkpoint also recovers 0/10 (Table 2). This fails our declared pilot pass rule, so successful recovery cannot rank the methods.

The procedure proposes $\mathbf{R}_{\mathrm{trial}} = \mathbf{R} + \eta\widehat{\mathbf{F}}$, where η converts predicted forces into coordinate changes. It starts at 0.2 Å^2/eV and can grow to 1.0. Backtracking halves a failed trial's scale; each branch permits 1,500 force attempts.

The procedure checks force convergence first; otherwise, forces pointing broadly in compatible directions can permit acceptance. There is no energy-decrease test or per-atom movement cap, so a scale limit does not cap movement when forces are large.

Table 2: All checkpoints recover 0/10 original targets. "Wrong stop" meets the predicted-force rule but misses the target shape. Reference-start controls begin at the six target minima; they are separate follow-up checks, called post-hoc analyses.

Checkpoint	Wrong stop	No convergence	Reference-start recovery
Energy C, seed 0	2	8	4/6
Direct, seed 0	4	6	1/6
Direct, seed 1	4	6	0/6
Consistency 0.1, seed 0	6	4	0/6
Consistency 0.1, seed 1	9	1	0/6

Across 50 model/branch outcomes, 25 stop with wrong geometry and 25 fail to converge. Reusing the same five connections does not create 50 independent paths. The reference-start controls favour the energy checkpoint under this procedure, but do not isolate a consistency-penalty effect.

5.1 Why a small predicted force can be misleading

A 155-evaluation check reproduced all 50 stopping flags. Every model exceeds the stopping force threshold at all 11 stored aspirin minima, which include the six targets. The reference labels themselves stay below 2.745 meV/Å. This mismatch at exact reference shapes does not exclude nearby shapes meeting both tolerances.

All 23 force-converged direct/consistency endpoints have disconnected 5 Å neighbour graphs, with atoms in different fragments beyond the model's interaction cutoff. These are not intact molecular shapes, although internal fragment forces can remain. Only seven endpoints have every atom isolated, where the direct model predicts exactly zero force automatically; this does not establish a correct molecular minimum.

Four selected trace replays reproduced the saved endpoints. Two separated, with largest accepted atom movements of 436.721 Å for direct prediction and 268.047 Å for energy-derived forces. The other two remained connected, with maxima 0.086492 Å for consistency and 0.006083 Å for energy-

derived forces. The separated energy trace ended without convergence; both force-converged energy endpoints remained connected.

These checks expose model and procedure limitations, without proving that the procedure alone caused failures or that real molecular break-up occurred. Better agreement on stored examples cannot guarantee correct shapes, and zero recovery cannot establish method equivalence.

Evidence covers aspirin, five validation connections and two seeds, with shared minima. Validation selects settings and BEST; test generalization is unestablished. Failed and incomplete trials remain recorded.

6 A related representation check and the remaining work

Faculty suggested comparing models with and without rotation symmetry; our internal-feature check is related but different. E64 is the existing width-64 model with scalar numbers and directional vectors; we reused its two fits. Four new scalar-only fits match its width (S64) or approximate parameter count (SP). All models retain force rotation symmetry, so this is not a no-symmetry comparison. E64 had lower validation force error in both seeds under E64-selected settings, without separate scalar tuning. Findings remain preliminary, await independent interpretation review, and establish neither feature causality nor better recovery; records: `reports/evidence/2026-10-20/representation/`.

Changes from the proposal. We completed two seeds rather than the proposed three, and paracetamol replication is deferred until aspirin recovery is feasible. A revised adequacy protocol followed the preserved small-subset failures; responsibilities changed, and the representation check adds an extension. Our models use a 5 Å cutoff; faculty suggested studying interactions beyond 7 Å as future work. These results do not show that missing long-range interactions caused the failures.

Three next actions. First, verify the MACE loader against saved predictions and run the unchanged ten-start and six-reference controls. Second, use these controls to assess a separately declared per-atom movement cap on the same checkpoints and starts, labelling reused-validation analysis as post-hoc.

Third, measure closed-loop work and prediction time on identical hardware, declaring any matched-wall-time training budget before new runs. Work adds products of matching force and displacement components around a closed path; forces from one smooth energy have an exact zero closed-loop integral. Numerical integration can introduce residual error.

Team contributions and reproducibility. Shedrack completed data preparation, model/training integration and current MACE/E3x experiments; Olufunke authored the recovery evaluator. Emmanuel Aaron owns physical/cost analysis; Muhammed Idris owns MACE loading, prediction checks and independent reproduction.

Assigned work is not completed contribution credit. Research code/evidence: fixed snapshot 66d0821. Before submission, confirm contributions, private grader access, live course instructions and team approval.

References

[1] V. Cărare et al. Global properties of the energy landscape: a testing and training arena for machine learned potentials. *npj Computational Materials*, 12:9, 2026.

[2] F. Bigi, M. F. Langer and M. Ceriotti. The dark side of the forces: assessing non-conservative force models for atomistic machine learning. *ICML*, PMLR 267:4384–4414, 2025.

[3] O. T. Unke and H. Maennel. E3x: E(3)-equivariant deep learning made easy. arXiv:2401.07595, 2024.

[4] I. Batatia et al. MACE: Higher order equivariant message passing neural networks for fast and accurate force fields. *NeurIPS*, 35:11423–11436, 2022.

[5] M. Takamoto, V. Zaverkin and M. Niepert. Physics-informed weakly supervised learning for interatomic potentials. *ICML*, PMLR 267:58282–58310, 2025.