Do Consistent Learned Forces Find the Right Molecular Shape?

Shedrack Eze (seze) **Olufunke Abodunrin (oabodunr)**
Emmanuel Aaron (eaaron) **Muhammed Idris (midris)**
Path 3: Mentored MLIP Project

Abstract

Predicting the forces on atoms can make molecular simulations much faster than repeated quantum-chemistry calculations. Some neural networks predict those forces directly; others calculate them from a learned energy. Direct predictions can point toward the wrong stable molecular shape even when their average error is low. This project asks whether training direct forces to agree more closely with the learned energy improves recovery of the correct shape from unseen pathways. Models will use the same reference data and be compared on shape recovery, force error, and computing cost. The answer is unknown.

1 Overview and Context

A molecule can take several shapes as its atoms move. A reference quantum-chemistry calculation assigns each shape an energy and a force on every atom. Low-energy shapes are stable; higher-energy pathways connect them. Repeating the reference calculation during a simulation is expensive, so a neural network can learn an approximation. The usual score asks how close its forces are to known examples. That score does not tell whether following the predicted forces from a new pathway ends at the right stable shape.

An energy-based model calculates forces as the downhill direction of its predicted energy. A direct-force model predicts the arrows themselves, which may be faster, but its arrows can disagree with the energy it predicts. Earlier research already found that direct forces can score well on individual examples yet perform poorly in searches for transition states. The question here is more specific: if training makes the direct arrows agree better with the learned energy, do they lead to the correct stable shape more often on unseen paths, even after accounting for force error and cost?

2 Related Work and Background

Geometry-aware models such as PaiNN already learn energy-based forces [1]. Bigi et al. found direct-force problems in simulations and optimization [2]; Zhao et al. showed that lower force error need not improve reaction transition-state search [3]. Landscape17 documented missed conformational paths and extra stable shapes [5]. Force/energy disagreement and conservation-related losses have also been studied [6, 7]. The proposed contribution is to test whether reducing disagreement improves recovery from unseen conformational paths at similar error, not to claim a new penalty.

3 Methodology

Landscape17 provides public atomic positions, energies, and forces from one hybrid density-functional-theory (DFT) calculation [5]. It includes 37 aspirin and nine paracetamol paths between stable shapes. Entire paths, not nearby frames, will be assigned to separate training, validation, and test groups. Shared shapes will be flagged. Labels from rMD17 use a different DFT calculation and will not be mixed in.

Three models will share one E3x-based network [4]: energy-derived forces, direct forces, and direct forces trained to agree with the energy-derived forces. The extra training cost will be counted; agreement is not guaranteed. A MACE model on the same split provides an established energy-based comparison [8].

Validation paths will select model settings. Training time, total computing cost, and time per force prediction will be reported.

Each model will start from the same unseen positions and follow its predicted forces with the same relaxation procedure. Main measures are average force-component error and the share reaching the stable DFT reference shape. Wrong endpoints and failed runs count separately. A closed-loop test checks whether forces behave as slopes of one energy surface; speed and forces at reference stationary points are secondary. A direct model's energy output alone cannot justify reaction barriers or rates. The appendix specifies the checks.

The key comparison is correct-shape recovery for consistency-trained versus ordinary direct forces at similar average error. Energy-based forces are a reference point. Whole paths are the unit of analysis. If errors cannot be matched, only the error–cost–recovery tradeoff will be reported.

4 Baseline and Extensions

The energy-based E3x model is the baseline. Ordinary direct E3x is the control for a predeclared sweep of consistency-training strengths. Aspirin is the main test; paracetamol checks whether the pattern repeats in another molecule. MACE provides a stronger established comparison. Adding more pathway training data is a separate experiment because Landscape17 already studied its general benefit [5].

5 Results and Analysis

The hypothesis is that making direct forces more consistent with energy will improve correct-shape recovery when average force errors are similar. If any gain comes only from lower force error, or if consistency improves without better recovery, that hypothesis will not be supported. No models have been trained. Variation across training runs and paths will be reported.

6 Discussion

There are few independent paths, some share stable shapes, and models may reach different error levels. Whole-path splits, fixed starting positions, and model choices made only on validation data limit unfair comparisons. A stopped direct-force relaxation is not necessarily an energy minimum. Even a positive result would apply to these molecules and this network; it would not establish reaction rates or a complete landscape. Zhao et al.'s reaction-search findings remain relevant prior work [3].

7 Future Directions

Before the October 23 midterm: check the Landscape17 data, fix the path splits and starting positions, and test the first energy-based and direct models. Before the December 9 course report: run the consistency sweep, MACE comparison, paired relaxations, and paracetamol check.

8 Administrative

Provisional roles: data/evaluation (Eze), model implementation (Abodunrin), experiments (Aaron), and infrastructure/writing (Idris). Modeling has not begun; the team is currently reviewing the literature and refining the study design. Project repository (private): `https://github.com/Shedrackeze002/idl-molecular-force-consistency`.

Appendix

Experimental Protocol Details

The public Landscape17 archive contains 37 aspirin and nine paracetamol transition-path groups, with atomic coordinates and hybrid-DFT energy/force labels along the paths and at stationary points [5]. Path identifiers, rather than individual frames, define the partitions. Validation and test path groups are disjoint; duplicate structures and shared minima will be audited before fitting. Hyperparameters, penalty strengths, and MAE-matched checkpoint pairs are chosen using training and validation data only. Test groups are evaluated once after selection. Three seeds and a same-split MACE reference are planned core controls; a pilot will size the dedicated compute needed.

The direct consistency arm minimizes an energy/force objective plus $\lambda_C \|\hat{\mathbf{F}}^{\mathrm{D}} + \nabla \hat{E}\|_2^2$ on training geometries. The $\lambda_C = 0$ direct model and gradient-force model are controls. Head parameter counts, equal-update and equal-wall-time results, total GPU-hours, and batched force-inference latency will be reported on identical hardware.

Held-out path frames and transition-state coordinates will not tune the models; a minimum shared with training paths will be labeled as seen. Identical perturbed starts from held-out paths will be relaxed with a fixed force-driven optimizer and evaluation budget. Success requires convergence and a symmetry-aware structural match to the connected reference minimum; unmatched converged fixed points and nonconvergence are separate outcomes. Closed-loop work is measured on preselected geometry loops with numerical refinement. The primary analysis compares path-group outcomes across models within validation-defined force-MAE bands, with uncertainty reported over groups and seeds. Full transition-state networks, barrier heights, and reaction rates are outside this proposal's direct-force claims.

References

[1] K. T. Schütt, O. T. Unke, and M. Gastegger. Equivariant message passing for the prediction of tensorial properties and molecular spectra. *ICML*, 2021.

[2] F. Bigi, M. F. Langer, and M. Ceriotti. The dark side of the forces: assessing non-conservative force models for atomistic machine learning. *ICML*, 2025.

[3] Q. Zhao et al. Harnessing machine learning to enhance transition state search with interatomic potentials and generative models. *Advanced Science*, 12:e06240, 2025.

[4] O. T. Unke and H. Maennel. E3x: E(3)-equivariant deep learning made easy. arXiv:2401.07595, 2024.

[5] V. Cărare et al. Global properties of the energy landscape: a testing and training arena for machine learned potentials. *npj Computational Materials*, 12:9, 2026 (online 2025).

[6] Anonymous. When forces disagree: a data-free fast diagnostic from internal consistency of direct-force neural network potentials. OpenReview manuscript under review, 2026.

[7] M. Takamoto, V. Zaverkin, and M. Niepert. Physics-informed weakly supervised learning for interatomic potentials. *ICML*, 2025.

[8] I. Batatia et al. MACE: Higher order equivariant message passing neural networks for fast and accurate force fields. *NeurIPS*, 2022.