

BMF 1-1

Prove the correctness of the Horner's rule:

- Show that

$$(a \oplus b) \otimes c = (a \otimes c) \oplus (b \otimes c) \iff (\otimes c) \cdot \oplus / = \oplus / \cdot (\otimes c) * \cdot$$

- Show that

$$f = \oplus / \cdot \otimes / * \cdot \text{tails}$$

satisfy the equations $f [] = e$ and $f(x \# [a]) = f x \odot a$.

Proof: (a) \Leftarrow is trivial, by applying both sides to $[a, b]$ we obtain

$$(a \oplus b) \otimes c = (\otimes c)(a \oplus b) = (\otimes c) \cdot \oplus / [a, b] = \oplus / \cdot (\otimes c) * [a, b] = \oplus / [a \otimes c, b \otimes c] = (a \otimes c) \oplus (b \otimes c).$$

\Rightarrow can be proved by induction: The base $n = 1$ is trivial. If it holds for arrays of length n , then for any array $[a_1, \dots, a_{n+1}]$,

$$\begin{aligned} (\otimes c) \cdot \oplus / [a_1, \dots, a_{n+1}] &= (a_1 \oplus \dots \oplus a_{n+1}) \otimes c = ((a_1 \oplus \dots \oplus a_n) \otimes c) \oplus (a_{n+1} \otimes c) \\ &= ((\otimes c) \cdot \oplus / [a_1, \dots, a_n]) \oplus (a_{n+1} \otimes c) = (\oplus / \cdot (\otimes c) * [a_1, \dots, a_n]) \oplus ((\otimes c) * [a_{n+1}]) \\ &= \oplus / \cdot (\otimes c) * [a_1, \dots, a_{n+1}]. \end{aligned}$$

Hence it holds for any array.

(b) $f [] = \oplus / \cdot \otimes / * \cdot \text{tails} [] = \oplus / \cdot \otimes / * [] = \oplus / [e] = e$, and

$$\begin{aligned} f(x \# [a]) &= \oplus / \cdot \otimes / * \cdot \text{tails}(x \# [a]) = \oplus / \cdot \otimes / * \cdot ((\# [a]) * \cdot \text{tails} x \# []) \\ &= \oplus / \cdot ((\otimes / * \cdot (+ + [a]) * \cdot \text{tails} x) \# (\otimes / * [])) \\ &= \oplus / \cdot ((\otimes a) * \cdot \otimes / * \cdot \text{tails} x \# []) \\ &= ((\otimes a) \cdot (\oplus / \cdot \otimes / * \cdot \text{tails} x)) \oplus e \\ &= (f x \otimes a) \oplus e = f x \odot a \end{aligned}$$

BMF 1-2

Code the derived linear algorithm for mss in Haskell:

```
mss :: [Integer] -> Integer
mss xs = snd $ foldl f (0,0) xs
    where f = \ (s,ms) x -> let s' = if s+x > 0 then s+x else 0
                             ms' = max s' ms in (s',ms')
```

BMF 1-3

Prove the Segment Decomposition Theorem:

Suppose S, T are defined by $S = \oplus / \cdot f * \cdot \text{segs}$, $T = \oplus / \cdot f * \cdot \text{tails}$. If T can be expressed in the form $T = h \cdot \odot \nearrow_e$, then we have $S = \oplus / \cdot h * \cdot \odot \nearrow_e$.

Proof:

$$\begin{aligned}
S &= \otimes / \cdot f * \cdot \text{segs} = \otimes / \cdot f * \cdot \# / \cdot \text{tails} * \cdot \text{inits} \\
(\text{map promotion}) &= \otimes / \cdot (\# / \cdot f * *) \cdot \text{tails} * \cdot \text{inits} \\
(\text{reduce promotion}) &= (\otimes / \cdot \otimes / *) \cdot f * * \cdot \text{tails} * \cdot \text{inits} \\
&= \otimes / \cdot \otimes / * \cdot f * * \cdot \text{tails} * \cdot \text{inits} \\
(\text{map distribution}) &= \otimes / \cdot (\otimes / \cdot f * \cdot \text{tails}) * \cdot \text{inits} \\
(T = h \cdot \odot \not\rightarrow_e) &= \otimes / \cdot (h \cdot \odot \not\rightarrow_e) * \cdot \text{inits} \\
(\text{map distribution}) &= \otimes / \cdot h * \cdot (\odot \not\rightarrow_e) * \cdot \text{inits} \\
&= \otimes / \cdot h * \cdot \odot \not\rightarrow_e
\end{aligned}$$