

## Contents

<b>1 Homework 1: Schröder-Bernstein Theorem</b>	<b>3</b>
1.1 PSA	3
1.2 PSB: Countable and Uncountable Sets	4
1.3 PSC: Schröder-Bernstein Theorem	6
1.4 PSD: Details of Dedekind Cut	8
<b>2 Homework 2: Cesàro sum</b>	<b>9</b>
2.1 PSA	9
2.2 PSB: Calculation of Limits	11
2.3 PSC: Riemann Rearrangement Theorem	13
2.4 PSD: Cesàro Sum	14
2.5 PSE: Definition of $\sqrt[n]{x}$ and $b^x$	15
2.6 PSF	17
2.7 PSG: Banach-Mazur Game	18
2.8 Problem H	18
2.9 Problem I: Binary Expansion	19
2.10 Problem J	20
<b>3 Homework 3: Basel Problem</b>	<b>20</b>
3.1 PSA	20
3.2 PSB	22
3.3 PSC	25
3.4 PSD	26
3.5 PSE: Reciprocal Sum of Primes	27
3.6 PSF: Euler's "Proof" of the Basel Problem	27
<b>4 Homework 4: Topology</b>	<b>29</b>
4.1 PSA: Topology on Metric Spaces	29
4.2 PSB	31
4.3 PSC: Root of Function:	32
4.4 PSD: Calculation of Limits	33
4.5 Problem E	36
4.6 Problem F	37
4.7 Problem G	37
<b>5 Homework 5: Infinity of Prime</b>	<b>37</b>
5.1 PSA	37
5.2 PSB: Uniform Continuity	39
5.3 PSC: Existence of Limits	40
5.4 PSD: Problems on Uniform Continuity	41
5.5 PSE:	42
5.6 PSF: Calculate Limits	42
5.7 Problem G	43
5.8 Problem H	43
5.9 Problem I	44
5.10 Ex: Proof of the infinity of primes using topology	44
<b>6 Homework 6: Takagi Function</b>	<b>44</b>
6.1 PSA: Calculating Derivatives	44
6.2 PSB	49

6.3	PSC	50
6.4	PST: Takagi Function	51
<b>7</b>	<b>Homework 7: Émile Borel Lemma</b>	<b>55</b>
7.1	PSA	55
7.2	PSB	56
7.3	PSC	59
7.4	PSD	59
7.5	PSE: Roots of Polynomials	59
7.6	PSF: Émile Borel's Lemma	60
7.7	PSG: Midterm Test Part B	64
<b>8</b>	<b>Homework 8; Strum-Liouville Theory</b>	<b>68</b>
8.1	PSA: Convex functions	68
8.2	PSB	70
8.3	PSC	71
8.4	PSD	73
8.5	PSE	74
8.6	PSF	75
8.7	PSG	75
8.8	Problem S: Strum-Liouville Theory	76
<b>9</b>	<b>Homework 9: Stone-Weierstrass Theorem</b>	<b>79</b>
9.1	PSA	79
9.2	PSB: Calculating Integrals	81
9.3	Problem W: Stone-WeierstraSS Theorem	88
<b>10</b>	<b>Homework 10: Irrationality of <math>\pi</math></b>	<b>91</b>
10.1	PSA	91
10.2	PSB: On $\zeta(2)$	95
10.3	PSC: Calculation of Integrals	98
<b>11</b>	<b>Homework 11: Density of Sum of Squares</b>	<b>102</b>
11.1	PSA: Riemann Integral	102
11.2	PSB: Convex Functions	104
11.3	PSC: Integrals and Derivatives	105
11.4	PSD: Calculation of improper integrals	109
11.5	PSE: Density of sum of squares	113
<b>12</b>	<b>Homework 12: Oscillatory Integral</b>	<b>122</b>
12.1	PSA: Stieltjes Integral	122
12.2	PSB: Convergence of Improper Integrals	125
12.3	PSC: Oscillatory Integral	128

# 1 Homework 1: Schröder-Bernstein Theorem

## 1.1 PSA

**A1) Suppose a non-empty set  $X \subset \mathbb{R}$  has an upper bound, and  $M$  is an upper bound of  $X$ . The following two propositions are equivalent:**

- $M = \sup X$ .
- For any  $\varepsilon > 0$ , there exists an  $x \in X$  such that  $x > M - \varepsilon$ .

Proof:

$$M = \sup X \iff \forall M' < M, \exists x \in X, x > M' \iff \forall \varepsilon = M - M' > 0, \exists x \in X, x > M - \varepsilon.$$

**A2) Prove that every non-empty open interval contains infinitely many rational numbers.**

Proof: We only need to find one rational number  $q$  in the interval  $(a, b)$ , then we can apply the process to  $(a, q)$  and so on.

By the Archimedean rule, there is a positive integer  $N$  such that  $N(b - a) > 2$ , hence there exists an integer  $p$  such that  $p = \lfloor bN \rfloor \in (aN, bN)$ , and  $q = \frac{p}{N} \in (a, b) \cap \mathbb{Q}$ .

**A3) Let  $(X, d)$  be a metric space,  $Y \subset X$ . We define the distance function on  $Y$ :**

$$d_Y : Y \times Y \rightarrow \mathbb{R}, (y_1, y_2) \mapsto d_Y(y_1, y_2) = d(y_1, y_2).$$

Prove that  $d_Y$  is a distance function, and  $(Y, d_Y)$  is a metric space. We call  $d_Y$  the induced metric on  $Y$ , and  $(Y, d_Y)$  is called a subspace.

Proof: Trivial, since  $d_Y(y_1, y_2) = d(y_1, y_2)$ .

**A4) Let  $\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R} = \{(x_1, \dots, x_n) \mid x_i \in \mathbb{R}, \dots, x_n \in \mathbb{R}\}$ , for any  $x, y \in \mathbb{R}^n$ , we define**

$$d(x, y) = \sqrt{(x_1 - y_1)^2 + \cdots + (x_n - y_n)^2}, \quad x = (x_1, \dots, x_n), y = (y_1, \dots, y_n).$$

Prove that  $(\mathbb{R}^n, d)$  is a metric space.

Proof:

1.  $d(x, y) = 0 \iff x_i = y_i, \forall 1 \leq i \leq n \iff x = y$ .
2.  $d(x, y) = d(y, x)$  is trivial.
3.  $d(x, y) + d(y, z) \geq d(x, z)$  is the Minkowski inequality:

$$\left( \sqrt{\sum_{i=1}^n a_i^2} + \sqrt{\sum_{i=1}^n b_i^2} \right)^2 = \sum_{i=1}^n a_i^2 + b_i^2 + 2 \sqrt{\sum_{i=1}^n a_i^2 \sum_{i=1}^n b_i^2} \geq \sum_{i=1}^n a_i^2 + b_i^2 + 2a_i b_i = \sum_{i=1}^n (a_i + b_i)^2.$$

**A5) Given a metric space  $(X, d)$ , and  $Y \subset X$ . If for any  $x \in X$  and  $\varepsilon > 0$ , there exists  $y \in Y$  such that  $d(y, x) < \varepsilon$ , then we say  $Y$  is dense in  $X$ . Prove that the set of rational numbers is dense in  $\mathbb{R}$ .**

Proof: For any  $x \in \mathbb{R}$ , let  $N = \lfloor x \rfloor$ , then for any  $\varepsilon > 0$ , let  $q > 1/\varepsilon$ . Then for  $p \in [Nq, (N+1)q] \cap \mathbb{Z}$ , choose  $p$  such that  $|x - p/q|$  is minimal. Suppose  $p/q < x$ , then

$$2 \left| x - \frac{p}{q} \right| < \left| x - \frac{p}{q} \right| + \left| x - \frac{p+1}{q} \right| = \frac{1}{q} < \varepsilon.$$

Hence  $d(x, p/q) < \varepsilon$ .

**A6) For  $(x, y) \in \mathbb{R}^2$ , if its coordinates  $x$  and  $y$  are rational numbers, then we call this point a rational point. Prove that  $(\mathbb{R}^2, d)$  (refer to question A4) the set of rational points in  $\mathbb{R}^2$  is dense.**

Proof: By A5),  $\overline{\mathbb{Q}} = \mathbb{R}$ . Hence for any  $(x, y) \in \mathbb{R}^2$  and  $\varepsilon > 0$ , there exists  $(a, b) \in \mathbb{Q}^2$  such that  $|a - x|, |b - y| < \varepsilon/2$ . Then

$$d((x, y), (a, b)) = \sqrt{(a - x)^2 + (b - y)^2} < \varepsilon.$$

Hence  $\mathbb{Q}^2$  is dense in  $\mathbb{R}^2$

**A7) Prove that the axiom (F) and (O), and the boundedness principle imply the Archimedean axiom (A).**

Proof: Otherwise assume that  $\mathbb{N}$  has an upper bound. Then  $M = \sup \mathbb{N}$  exists. Let  $\varepsilon = 1/2$  then there is an  $n \in \mathbb{N}$  such that  $n > M - \varepsilon$ . Hence  $n + 1 > M$ , leading to contradiction.

**A8) (Existence of irrational numbers) Let  $X = \{x \in \mathbb{Q} \mid x^2 < 2\}$  be a bounded set, and  $\sqrt{2} = \sup X$ . Prove that  $\sqrt{2}$  is an irrational number.**

Proof: If  $\sqrt{2} = s = p/q$  is rational, then  $p^2 \geq 2q^2$ , otherwise let  $x = s(2 - s^2)/4 + s$ , then  $s < x$  and  $x^2 < 2$ , a contradiction. If  $s^2 > 2$ , then  $x = s(2 - s^2)/4 < s$  and  $x^2 > 2$ , hence  $x$  is an upper bound of  $X$ , leading to contradiction. Therefore  $s^2 = 2$  which is impossible.

**A9) Prove that every open interval contains infinitely many irrational numbers.**

Proof: Otherwise the open interval will be a countable set.

## 1.2 PSB: Countable and Uncountable Sets

Let  $\mathbb{N}$  denote the set of natural numbers (including 0).  $X$  is a set, if there is an injective map  $f : X \rightarrow \mathbb{N}$ , then we say  $X$  is countable; if  $X$  is not countable, then we say  $X$  is uncountable.

**B1) Prove that finite sets are countable.**

Proof: For any finite set  $X = \{a_1, \dots, a_n\}$ , the map  $f : a_k \mapsto k$  is an injective, hence  $X$  is countable.

**B2) Prove that subsets of countable sets are countable.**

Proof: If  $X$  is countable and  $Y \subset X$ , then there is an injective map  $f : X \rightarrow \mathbb{N}$ , so  $f|_Y : Y \rightarrow \mathbb{N}$  is an injective map, hence  $Y$  is countable.

**B3) Prove that if  $X$  is a countable set, then we can always write  $X = \{x_1, x_2, x_3, \dots\}$  (that is, the elements of  $X$  can be indexed by natural numbers).**

Proof: Let  $I = \{n \in \mathbb{N} : f^{-1}(n) \neq \emptyset\}$ ,  $x_k = f^{-1}(\min I \setminus \{f(x_1), \dots, (x_{k-1})\})$ . Then  $x_x \in X$ , and for any  $x \in X$ ,  $f(x) \in I$  hence  $x \in \{x_1, \dots, x_{f(x)}\}$ . Therefore  $X = \{x_1, \dots, x_n, \dots\}$ .

**B4) Prove that the set of rational numbers  $\mathbb{Q}$  is countable.**

Proof: List every positive rational number as below:

$$\frac{1}{1}, \frac{1}{2}, \frac{2}{1}, \frac{1}{3}, \frac{2}{2}, \frac{3}{1}, \frac{1}{4}, \frac{2}{3}, \dots$$

such that  $p/q$  is before  $m/n$  if  $p+q < m+n$  or  $p+q = m+n$  and  $p < m$ , then every number in  $\mathbb{Q}_{>0}$  is listed at least once. Hence  $\mathbb{Q}_{>0}$  is countable and so is  $\mathbb{Q}$ .

**B5) Prove that the countable union of countable sets is countable, that is, if  $X_1, X_2, \dots, X_n, \dots$  are all countable sets, then their union  $\bigcup_{n=1}^{\infty} X_n$  is also a countable set.**

Proof: Assume  $X_n$  are disjoint. Since  $X_n$  are countable, we can write

$$X_n = \{a_1^{(n)}, a_2^{(n)}, \dots, a_m^{(n)}, \dots\}.$$

Then

$$\bigcup_{n=1}^{\infty} X_n = \{a_1^{(1)}, a_2^{(1)}, a_1^{(2)}, a_3^{(1)}, \dots\}$$

where the order is the same as in B4). Hence  $\bigcup_{n \geq 1} X_n$  is countable.

**B6) If  $X$  is countable, and the map  $f : X \rightarrow Y$  is surjective, then  $Y$  is countable.**

Proof: Since  $X$  is countable, there is an injective map  $g : X \rightarrow \mathbb{N}$ . Let

$$h : Y \rightarrow \mathbb{N}, y \mapsto \min g(f^{-1}(\{y\})).$$

then  $g$  is injective, hence  $Y$  is countable.

**B7) Prove the following using proof by contradiction:  $\mathbb{R}$  is uncountable.**

B7-1) Suppose  $J \subset \mathbb{R}$  is a closed interval and its length  $|J| > 0$ . For any  $x \in \mathbb{R}$ , there always exists an interval  $I \subset J$  such that  $|I| > 0$  and  $x \notin I$ .

Proof: Any closed interval  $J = [a, b]$  can be written in the form  $J = A \cup B \cup C$ , where  $A = [a, \frac{2a+b}{3}]$ ,  $B = [\frac{2a+b}{3}, \frac{a+2b}{3}]$ ,  $C = [\frac{a+2b}{3}, b]$ , and  $x$  can only be in at most 2 of these sets. Hence we can choose a set  $I$  in  $A, B, C$ .

B7-2) Prove that if  $\{x_1, x_2, \dots\}$  is a countable subset of  $\mathbb{R}$ , then there exists a nested interval sequence  $I_1 \supset I_2 \supset \dots$  such that for any  $n$ ,  $x_n \notin I_n$ .

Proof: Simple application of B7-1)

B7-3) Prove that  $\mathbb{R}$  is uncountable.

Proof: If  $\mathbb{R}$  is countable, write  $\mathbb{R} = \{r_1, r_2, \dots\}$ , then set  $I_0 = [0, 1]$ . By B7-2) we can obtain a sequence  $I_0 \supset I_1 \supset \dots$  such that  $x_n \notin I_n$  for any  $n$ . Hence

$$\bigcap_{n=0}^{\infty} I_n = \emptyset,$$

leading to contradiction.

**B8) Prove that if  $X$  is an uncountable set, and  $A$  is a countable subset of  $X$ , then  $X - A$  is uncountable.**

Proof: Otherwise suppose that both  $A$  and  $X - A$  is countable, then there exist injective mappings  $f : A \rightarrow \mathbb{N}$  and  $g : X - A \rightarrow \mathbb{N}$ . Define

$$h : X \rightarrow \mathbb{N}, x \mapsto \begin{cases} 2f(x), & x \in A, \\ 2g(x) + 1, & x \notin A. \end{cases}$$

Then  $h$  is injective, hence  $X$  is countable.

**B9) Prove that any interval of non-zero length (open or closed) is uncountable.**

Proof: Same as B7).

Or use the fact that  $\mathbb{R}$  is the countable union of intervals of the same length, and the countable union of countable sets is still countable.

**B10) Prove that the set of complex numbers  $\mathbb{C}$  is uncountable.**

Proof:  $\mathbb{C}$  has an uncountable subset  $\mathbb{R}$ .

**B11) Suppose  $\mathcal{I}$  is a collection of non-overlapping closed intervals, satisfying the following property: for any  $I, J \in \mathcal{I}$ , if  $I \neq J$ , then their intersection is empty, i.e.,  $I \cap J = \emptyset$ . Prove that  $\mathcal{I}$  is countable.**

Proof: For any  $I \in \mathcal{I}$ , there exists a rational number  $r_I \in I$ . Consider  $f : \mathcal{I} \rightarrow \mathbb{Q}$ ,  $I \mapsto r_I$ , then  $f$  is injective. Since  $\mathbb{Q}$  is countable, so is  $\mathcal{I}$ .

### 1.3 PSC: Schröder-Bernstein Theorem

Suppose  $X$  and  $Y$  are two sets, and mappings  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$  are both injective. Let  $X' = X - g(Y)$ .

**C1) If  $X$  is a finite set, prove that there exists a bijection  $\varphi : X \rightarrow Y$ .**

Proof:  $g : Y \rightarrow X$  is injective and  $X$  is finite,  $\implies Y$  is finite. Hence  $|X| \leq |Y|$ , and  $|X| \geq |Y|$ , so  $|X| = |Y|$ . Therefore we can write  $X = \{x_1, x_2, \dots, x_n\}$  and  $Y = \{y_1, y_2, \dots, y_n\}$ , and obtain

$$\varphi : X \rightarrow Y, x_k \mapsto y_k.$$

**C2) If  $X$  is countable, prove that there exists a bijection  $\varphi : X \rightarrow Y$ .**

Proof: Assume  $X$  is infinite, then  $Y$  is countable (by  $g$ ) and infinite (by  $f$ ). Hence we can list  $X = \{x_1, x_2, \dots\}$  and  $Y = \{y_1, y_2, \dots\}$  and define

$$\varphi : X \rightarrow Y, x_k \mapsto y_k.$$

---

From now on, we impose no restrictions on  $X$ . Let  $h : X \rightarrow X$  be the composite map  $h = g \circ f$ .

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow h & & \downarrow g \\ X & \leftarrow & \end{array}$$

**C3) Consider the set family  $\mathcal{F} = \{A \subset X \mid X' \cup h(A) \subset A\}$ . Prove that  $\mathcal{F}$  is non-empty.**

Proof:  $X \in \mathcal{F}$ .

**C4) Prove that if  $A \in \mathcal{F}$ , then  $X' \cup h(A) \in \mathcal{F}$ .**

Proof: If  $A \in \mathcal{F}$  then  $X' \cup h(A) \subset A$ , hence (let  $B$  denote  $X' \cup h(A)$ )

$$X' \cup h(B) \subset X' \cup h(A) = B.$$

**C5) We define**

$$A_0 = \bigcap_{A \in \mathcal{F}} A = \{x \in X \mid \text{for any } A \in \mathcal{F}, \text{ we have } x \in A\}.$$

Prove that  $A_0 \in \mathcal{F}$ .

Proof:

$$X' \cup h(A_0) \subset X' \cup \left( \bigcap_{A \in \mathcal{F}} h(A) \right) = \bigcap_{A \in \mathcal{F}} X' \cup h(A) \subset \bigcap_{A \in \mathcal{F}} A = A_0.$$

Hence  $A_0 \in \mathcal{F}$ .

**C6) Prove that  $X' \cup h(A_0) = A_0$ .**

Proof:

$$A_0 \in \mathcal{F} \implies X' \cup h(A_0) \in \mathcal{F} \implies A_0 \subset X' \cup h(A_0).$$

The other side is proved in C5).

**C7) Let  $B_0 = X - A_0$ . Prove that  $f(A_0) \cap g^{-1}(B_0) = \emptyset$  and  $f(A_0) \cup g^{-1}(B_0) = Y$ .**

Proof: If  $f(A_0) \cap g^{-1}(B_0) \neq \emptyset$ , then there exist  $a \in A_0, b \in B_0$  such that  $f(a) = g^{-1}(b)$ , i.e.  $b = h(a)$ . Since  $a \in A_0$ , for any  $A \in \mathcal{F}$ ,  $a \in A$ , hence  $b = h(a) \in X' \cup h(A) \subset A$ . Therefore  $b \in A_0$ , a contradiction.

Otherwise if there exists  $y \in Y$  such that  $y \notin f(A_0) \cup g^{-1}(B_0)$ , then  $g(y) \notin B_0 \implies g(y) \in A_0$ . Let  $z = g(y) \in A_0 \cap g(Y)$ , then  $z \notin X'$  so  $z \in h(A_0)$  by C6). Let  $z = h(t)$  then  $y = f(t) \in f(A_0)$  since  $g$  is injective, leading to contradiction.

**C8) We define the map  $\varphi : X \rightarrow Y$ . For  $x \in X$ , we require**

$$\varphi(x) = \begin{cases} f(x), & \text{if } x \in A_0; \\ g^{-1}(x), & \text{if } x \in B_0. \end{cases}$$

Prove that this is a bijection.

Proof:

1.  $\varphi$  is injective: for any  $x, y \in A_0, x \neq y$ ,  $\varphi(x) \neq \varphi(y)$  since  $f$  is injective. For any  $x, y \in B_0, x \neq y$ ,  $\varphi(x) \neq \varphi(y)$  since  $g$  is a mapping. For any  $x \in A_0, y \in B_0$ ,  $\varphi(x) \neq \varphi(y)$  since  $f(A_0) \cap g^{-1}(B_0) = \emptyset$ .
2.  $\varphi$  is surjective:  $\varphi(X) = \varphi(A_0 \cup B_0) = f(A_0) \cup g^{-1}(B_0) = Y$ .

Based on the above, we have proved:

**Theorem (Schröder-Bernstein).** If there exist injective maps  $f : X \rightarrow Y$  and  $g : Y \rightarrow X$ , then there exists a bijection  $\varphi : X \rightarrow Y$  between the two sets.

### 1.4 PSD: Details of Dedekind Cut

The goal of this part of the exercise is to complete the part of the Dedekind cut construction method taught in class, thereby providing a complete proof for the construction of real numbers.

**D1) Prove that if  $X$  and  $Y$  are both Dedekind cuts, then the product  $X \cdot Y$  as defined in the lecture is also a Dedekind cut, i.e.,**

$\times : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}, (X, Y) \mapsto X \cdot Y,$

is well-defined. (Hint: You only need to prove the case where  $X > 0, Y > 0$ .)

Proof: The set  $X \cdot Y$  is define as  $Z = \bar{0} \cup \{x \cdot y : x, y \geq 0, x \in X, y \in Y\}$ . Let  $Z' = \mathbb{Q} - Z$ , then

1.  $Z \neq \emptyset, Z' \neq \emptyset$ , since for any  $x \in X', y \in Y', x \cdot y \notin Z$ .
2. For any  $z \in Z, z' \in Z'$ , if  $z' < z$  then  $z > 0$ . So assume  $z = x \cdot y, x \in X, y \in Y, x, y \geq 0$ , then  $z' = x \cdot (yz'/z) \in Z$ , a contradiction.
3. If  $Z$  has a maximal element  $z = x \cdot y, x, y \geq 0, x \in X, y \in Y$ , then since  $x, y$  are both not maximal, there exists  $x' \in X, y' \in Y$ , such that  $x < x', y < y'$  so  $z < z' = x' \cdot y' \in Z$ , a contradiction.

**D2) Prove that  $(X \cdot Y) \cdot Z = X \cdot (Y \cdot Z)$ . ( $\implies$  (F5))**

Proof: We only need to verify the case where  $X, Y, Z > 0$ . Then both  $(X \cdot Y) \cdot Z$  and  $X \cdot (Y \cdot Z)$  are the set

$$\bar{0} \cup \{x \cdot y \cdot z : x, y, z \geq 0, x \in X, y \in Y, z \in Z\}.$$

**D3) Prove that  $X \cdot Y = Y \cdot X$ . ( $\implies$  (F6))**

Proof: Same as D2).

**D4) Prove that  $X \cdot (Y + Z) = X \cdot Y + X \cdot Z$ . ( $\implies$  (F9))**

Proof: We can assume that  $X, Y, Z > 0$ , then

$$X \cdot (Y + Z) = \{xy + xz : x \in X, y \in Y, z \in Z\}$$

while

$$X \cdot Y + X \cdot Z = \{xy + x'z : x, x' \in X, y \in Y, z \in Z\}.$$

Hence  $X \cdot (Y + Z) \subset X \cdot Y + X \cdot Z$ .

For any  $xy + x'z \in X \cdot Y + X \cdot Z$ , suppose  $x \geq x'$ , then

$xy + xz \in X \cdot (Y + Z)$  and  $xy + x'z \leq xy + xz$ , so  $xy + x'z \in X \cdot Y + X \cdot Z$ , therefore  $X \cdot Y + X \cdot Z = X \cdot (Y + Z)$ .

**D5) Prove that  $\bar{1} \cdot X = X$  and  $\bar{1} \neq \bar{0}$ . ( $\implies$  (F7))**

Proof: Assume that  $X > 0$ , then  $\bar{1} \cdot X = \{u \cdot v : u < 1, v \in X\}$ . Foy any  $u < 1, v \in X$ ,  $u \cdot v < v$  hence  $u \cdot v \in X$ . For any  $x \in X$ , there exists  $x' \in X, x' > x$ , then  $x = x' \cdot (x/x') \in \bar{1} \cdot X$ . Therefore  $\bar{1} \cdot X = X$  and  $1/2 \in \bar{1} \setminus \bar{0}$ , so  $\bar{1} \neq \bar{0}$ .



**D6) Prove that if  $X \cdot Y = \bar{0}$ , then  $X = \bar{0}$  or  $Y = \bar{0}$ ; conversely, if  $X \geq \bar{0}, Y \geq \bar{0}$ , then  $X \cdot Y \geq \bar{0}$ . ( $\implies$  (O5))**

Proof: Otherwise there exists  $x, x' \in X, y, y' \in Y$ , such that  $x, y > 0, x', y' < 0$ . Hence  $xy, x'y \in X \cdot Y$ , where  $xy > 0 > x'y$ , so  $X \cdot Y \neq \bar{0}$ .

Suppose  $X, Y > 0$ , then there exists  $x \in X, y \in Y$  such that  $x, y > 0$ , hence  $0 < xy \in X \cdot Y$ , so  $X \cdot Y > \bar{0}$ .

**D7)  $X$  is a positive Dedekind cut. Prove that for any integer  $n$ , there exist  $x \in X, x' \in X'$  such that**

$$1 < \frac{x'}{x} < 1 + \frac{1}{n}.$$

Proof: Let  $l_0 = x \in X, r_0 = x' \in X'$ . Define  $l_n, r_n$  as follows: If  $(l_{n-1} + r_{n-1})/2 \in X$ , then  $l_n = (l_{n-1} + r_{n-1})/2, r_n = r_{n-1}$ , otherwise  $l_n = l_{n-1}, r_n = (l_{n-1} + r_{n-1})/2$ . Then

$$0 < \frac{r_n - l_n}{l_n} \leq \frac{1}{2} \frac{r_{n-1} - l_{n-1}}{l_{n-1}}.$$

Hence there exist such  $x, x'$ .

**D8) Prove that for any Dedekind cuts  $X$  and  $Y$ , if  $Y \neq \bar{0}$ , there exists a unique Dedekind cut  $Z$  such that**

$$Y \cdot Z = X.$$

We denote  $Z$  as  $\frac{X}{Y}$ . When  $X = \bar{1}$ , we also denote it as  $Y^{-1}$ . ( $\implies$  (F8))

Proof: By D6),  $Z$  is unique. By D2) we can assume that  $X = \bar{1}$ , and  $Y > 0$ . Let

$$Z = \left\{ \frac{1}{y} : y \in Y' \right\} \cup \bar{0} \cup \{0\}.$$

Then by D7),  $Y \cdot Z = \bar{1}$ .

## 2 Homework 2: Cesàro sum

### 2.1 PSA

**A1)  $\{x_n\}_{n \geq 1}$  is a bounded real sequence. Prove that there is a subsequence  $\{x_{n_i}\}_{i \geq 1}$  such that  $\lim_{i \rightarrow \infty} x_{n_i}$  exists and**

$$\lim_{i \rightarrow \infty} x_{n_i} = \limsup_{n \rightarrow \infty} x_n.$$

Proof: Let  $M = \limsup_{n \rightarrow \infty} x_n < \infty$ , then for any  $\varepsilon = 1/i > 0$  there exists  $N \geq n_{i-1}$  such that  $M \leq \sup_{k \geq N} x_k < M + \varepsilon$ . Hence there exists  $n_i \geq N$  such that  $x_{n_i} \in (M - \varepsilon, M + \varepsilon)$ . Take the sequence  $\{x_{n_i}\}_{i \geq 1}$  then  $\lim_{i \rightarrow \infty} x_{n_i} = \limsup_{n \rightarrow \infty} x_n$ .

**A2)  $\{x_n\}_{n \geq 1}$  is a real sequence. Prove that  $\{x_n\}_{n \geq 1}$  converges iff  $\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n$ .**

Proof: Since a sub-sequence of a Cauchy sequence converge to the same value as the original sequence,  $\implies$  is trivial by A1).

$\Leftarrow \lim_{n \rightarrow \infty} \sup_{k \geq n} x_k - \inf_{k \geq n} x_k = 0$  implies  $x_n$  is Cauchy, hence convergent.

**A3)**  $\{x^{(k)}\}_{k \geq 1} \subset \mathbb{R}^n$ , where  $x^{(k)} = (x_1^{(k)}, x_2^{(k)}, \dots, x_n^{(k)})$ . Then  $\{x^{(k)}\}_{k \geq 1}$  converges in  $\mathbb{R}^n$  iff for any  $i = 1, 2, \dots, n$ ,  $\{x_i^{(k)}\}_{k \geq 1}$  converges.

Proof: Use Cauchy sequences and the fact that for  $x = (x_1, x_2, \dots, x_n)$ ,

$$\max\{|x_k| : 1 \leq k \leq n\} \leq \|x\| \leq \sum_{k=1}^n |x_k|.$$

**A4)** Suppose  $\{z_n\}_{n \geq 1}, \{w_n\}_{n \geq 1}$  are two convergent complex sequences. Prove that if  $\lim_{n \rightarrow \infty} w_n \neq 0$ , then the sequence  $\{z_n/w_n\}_{n \geq 1}$  converges.

Proof: Suppose  $z = \lim_{n \rightarrow \infty} z_n$  and  $w = \lim_{n \rightarrow \infty} w_n$ , then

$$\left| \frac{z_n}{w_n} - \frac{z}{w} \right| \leq \frac{|w| \cdot |z_n - z|}{|w \cdot w_n|} + \frac{|z| \cdot |w_n - w|}{|w \cdot w_n|}.$$

Hence  $\left| \frac{z_n}{w_n} - \frac{z}{w} \right| \rightarrow 0$ , so  $\lim_{n \rightarrow \infty} z_n/w_n = z/w$ .

**A5)** Suppose  $\{a_n\}_{n \geq 1}$  is a monotonically decreasing sequence of positive reals, and  $\lim_{n \rightarrow \infty} a_n = 0$ . Prove that the series

$$a_1 - a_2 + a_3 - a_4 + \dots + (-1)^{n-1} a_n + \dots$$

converges.

Proof: Suppose  $a_n = a_1 - \sum_{k=1}^n b_k$ , then  $b_k \geq 0$  and  $\sum_{k=1}^{\infty} b_k = a_1$ . The series

$$\sum_{n=1}^{\infty} (-1)^{n-1} a_n = \sum_{n=1}^{\infty} b_{2n} < a_1$$

clearly converges.

**A6)**  $\{a_n\}_{n \geq 1} \subset \mathbb{C}$ . Prove that if  $\sum_{k=1}^{\infty} |a_k|$  converges, then  $\sum_{k=1}^{\infty} a_k$  converges.

Proof:  $\sum_{k=1}^{\infty} |a_k|$  converges implies for any  $\varepsilon > 0$ , there exists  $N$  such that for any  $n \geq N$ ,  $p \geq 0$ ,  $\sum_{k=n}^{n+p} |a_k| < \varepsilon$ . Note that  $\left| \sum_{k=n}^{n+p} a_k \right| \leq \sum_{k=n}^{n+p} |a_k|$ , so  $\sum_{k=1}^{\infty} a_k$  converges.

**A7)** Prove that we can define the exponential function on  $\mathbb{C}$ :

$$\exp : \mathbb{C} \rightarrow \mathbb{C}, z \mapsto \exp(z) = e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Proof: Use A6).

**A8)**  $\{a_n\} \subset \mathbb{C}$ . Suppose for any  $n \in \mathbb{N}$ ,  $a_n \neq 0$ . Let  $P_n = a_1 \cdot a_2 \cdots a_n$ . If  $\lim_{n \rightarrow \infty} P_n$  exists and is not 0, we call  $\prod_{n=1}^{\infty} a_n$  convergent and let  $\prod_{n=1}^{\infty} a_n = \lim_{n \rightarrow \infty} P_n$ . Prove that  $\prod_{n=1}^{\infty} a_n$  converges iff for any  $\varepsilon > 0$ , there exists  $N$  such that for any  $n \geq N$ ,  $p \geq 0$ ,

$$|a_n \cdot a_{n+1} \cdots a_{n+p} - 1| < \varepsilon.$$

Proof: If  $\lim_{n \rightarrow \infty} P_n = P$  exists and is non-zero, then for any  $\varepsilon > 0$ , there exists  $N$  such that for any  $n \geq N$ ,  $|P_n - P| < \varepsilon P/4$  and  $|P_n| > P/2$ . Then for any  $n \geq N$ ,  $p \geq 0$ ,  $|P_{n+p}/P_n - 1| < \varepsilon$ .

If for any  $\varepsilon > 0$ , there exists  $N$  such that for any  $n \geq N$ ,  $p \geq 0$ ,  $|P_{n+p} - P_n| < \varepsilon |P_n|$ , then let  $\varepsilon = 1$  we infer that  $P_n$  is bounded by some constant  $M$ . Hence the sequence  $\{P_n\}$  is Cauchy, and  $P = \lim_{n \rightarrow \infty} P_n$  cannot be zero, otherwise there is no such  $N$  for  $\varepsilon = 1/2$ .

**A9) Prove that  $\exp(x)$  is monotonically increasing on  $\mathbb{R}$ .**

Proof: For  $x, y \in \mathbb{R}$ ,

$$\exp(x) \cdot \exp(y) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \cdot \sum_{m=0}^{\infty} \frac{y^m}{m!} = \sum_{k=0}^{\infty} \sum_{n+m=k} \frac{x^n y^m}{k!} = \sum_{k=0}^{\infty} \frac{(x+y)^k}{k!} = \exp(x+y).$$

$\exp(x) \cdot \exp(-x) = \exp(0) = 1$  implies  $\exp(x) > 0$  for all  $x \in \mathbb{R}$ , so if  $x > y$ ,  $\exp(x)/\exp(y) = \exp(x-y) > 1 \implies \exp(x) > \exp(y)$ .

**A10) Suppose  $P(x)$  and  $Q(x)$  are polynomials of degree  $n, m$ , where  $m > n$ . Prove that**

$$\lim_{n \rightarrow \infty} \frac{Q(n)}{P(n)} = 0, \quad \lim_{n \rightarrow \infty} \frac{Q(n)}{e^n} = 0.$$

Proof: Suppose  $P(x) = \sum_{k=0}^n a_k x^k$  and  $Q(x) = \sum_{k=0}^m b_k x^k$ , then there exists  $N$  such that for any  $x \geq N$ ,  $|P(x)| > |a_n| x^n / 2$ ,  $|Q(x)| \leq \sum_{k=0}^m |b_k| \cdot x^m$ , and  $e^x \geq x^{m+1} / (m+1)!$ , hence

$$\left| \frac{Q(x)}{P(x)} \right| \leq \frac{2 \sum_{k=0}^m |b_k|}{|a_n|} \cdot x^{m-n} \rightarrow 0, \quad \left| \frac{Q(x)}{e^x} \right| \leq (m+1)! \sum_{k=0}^m |b_k| \cdot x^{-1} \rightarrow 0.$$

## 2.2 PSB: Calculation of Limits

**B1)**

$$\lim_{n \rightarrow \infty} \frac{n+10}{2n-1} = \frac{1}{2}.$$

**B2)**

$$\lim_{n \rightarrow \infty} \frac{\sqrt{n}+10}{2\sqrt{n}-1} = \frac{1}{2}.$$

**B3)**

$$\lim_{n \rightarrow \infty} \underbrace{0.99 \cdots 9}_{n \text{ times}} = 1.$$

**B4)**

$$\lim_{n \rightarrow \infty} \frac{1}{n(n+3)} = 0.$$

**B5)**

$$\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$$

B6)

$$\lim_{n \rightarrow \infty} \frac{2^n}{n!} = 0.$$

B7)

$$\lim_{n \rightarrow \infty} \frac{n!}{n^n} = 0.$$

B8)

$$\lim_{n \rightarrow \infty} \sqrt{n+10} - \sqrt{n+1} = 0.$$

B9)

$$\lim_{n \rightarrow \infty} \frac{1+2+\cdots+n}{n^2} = \frac{1}{2}.$$

B10)

$$\lim_{n \rightarrow \infty} \frac{1^2+2^2+\cdots+n^2}{n^3} = \frac{1}{3}.$$

B11)  $a > 0$

$$\lim_{n \rightarrow \infty} a^{1/n} = 1.$$

B12)  $a > 1$

$$\lim_{n \rightarrow \infty} \frac{n^{10000}}{a^n} = 0.$$

B13)

$$\lim_{n \rightarrow \infty} \frac{2^n + n}{3^n + n^2} = 0.$$

B14)

$$\lim_{n \rightarrow \infty} \frac{3^n + 2^n}{3^n + n^2} = 1.$$

B15)

$$\lim_{n \rightarrow \infty} \sqrt{n}(\sqrt{n+1} - \sqrt{n}) = \frac{1}{2}.$$

B16) same as B12)

B17)

$$\lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^n = e^{-1}.$$

B18)

$$\lim_{n \rightarrow \infty} \left(1 - \frac{1}{5n}\right)^{n+2019} = e^{-1/5}.$$

B19)

$$\lim_{n \rightarrow \infty} (n^3 + n^2 + 9n + 1)^{1/n} = 1.$$

B20)

$$\lim_{n \rightarrow \infty} (2018^n + 2019^n)^{1/n} = 2019.$$

## 2.3 PSC: Riemann Rearrangement Theorem

Suppose  $\sum_{n=1}^{\infty} a_n$  is conditionally convergent, we will prove that for and  $\alpha \in \mathbb{R} \cup \{-\infty, \infty\}$ , we can rearrange the sequence such that the new series sums to  $\alpha$ . Suppose  $\varphi : \mathbb{Z}_{\geq 1} \rightarrow \mathbb{Z}_{\geq 1}$  is a bijection, let  $b_k = a_{\varphi(k)}$ , then the sequence  $\{b_k\}_{k \geq 1}$  is called a rearrangement of  $\{a_n\}_{n \geq 1}$ . Let all non-negative terms of  $\{a_n\}_{n \geq 1}$ , listed in the same order as in  $\{a_n\}$  be  $c_1, c_2, \dots$ , and the negative terms be  $d_1, d_2, \dots$ .

**C1) Prove that**  $\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} d_n = 0$ .

Proof: Since  $\sum_{n=1}^{\infty} a_n$  is conditionally convergent,  $c_n, d_n$  both have infinite terms and  $\lim_{n \rightarrow \infty} a_n = 0$ . Therefore  $\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} d_n = 0$ .

**C2) Prove that**  $\sum_{n=1}^{\infty} c_n = \sum_{n=1}^{\infty} b_n = \infty$ .

Proof: Since  $\sum_{n=1}^{\infty} a_n$  is not absolutely convergent, the two series can not be both convergent. If one converges and the other doesn't, then  $\sum_{n=1}^{\infty} a_n$  will diverge. Hence they both diverge.

**C3) Prove that for any  $\alpha \in \mathbb{R}$ , there exists a rearrangement  $\{b_n\}$  of  $\{a_n\}$  such that  $\sum_{k=1}^{\infty} b_k = \alpha$ .**

Proof: Suppose  $\alpha \geq 0$ . Inductively define the indices  $u_i$  and  $v_i$  as follows ( $u_0 = v_0 = 0$ ): For  $i \geq 1$ , let  $u_i$  be the least index such that  $u_i > u_{i-1}$  and

$$\sum_{j=1}^{u_i} c_j - \sum_{j=1}^{v_{i-1}} d_j \geq \alpha,$$

and  $v_i$  be the least index such that  $v_i > v_{i-1}$  and

$$\sum_{j=1}^{u_i} c_j - \sum_{j=1}^{v_i} d_j \leq \alpha.$$

Let  $\varphi$  be the permutation such that

$$b_1 = c_1, b_2 = c_2, \dots, b_{u_1} = c_{u_1}, b_{u_1+1} = -d_1, \dots, b_{u_1+v_1} = -d_{v_1}, \dots$$

Since  $\sum_{n=1}^{\infty} c_n = \sum_{n=1}^{\infty} d_n = \infty$ ,  $u_i$  and  $v_i$  all exists, so  $\varphi$  is indeed a bijection. By definition we know that

$$\left| \sum_{j=1}^{u_i} c_j - \sum_{j=1}^{v_{i-1}} d_j - \alpha \right| \leq c_{u_i-1},$$

and

$$\left| \sum_{j=1}^{u_i} c_j - \sum_{j=1}^{v_i} d_j - \alpha \right| \leq d_{v_i-1}.$$

Since  $\lim_{n \rightarrow \infty} c_n = \lim_{n \rightarrow \infty} d_n = 0$ , the two values above both tend to 0. Note that the series  $\sum_{n=1}^{\infty} b_n$  is monotonic between these indices, hence  $\sum_{n=1}^{\infty} b_n = \alpha$ .

**C4) Prove that there exists a rearrangement  $\{x_k\}$  of  $\{a_n\}$  such that  $\sum_{k=1}^{\infty} x_k = \infty$ .**

Proof: Define  $u_i$  and  $v_i$  as in C3), such that

$$\sum_{j=1}^{u_i} c_j - \sum_{j=1}^{v_{i-1}} d_j \geq i \geq \sum_{j=1}^{u_i} c_j - \sum_{j=1}^{v_i} d_j.$$

Same as C3) define the sequence  $x_k$  and clearly  $\sum_{n=1}^{\infty} x_k = \infty$ .

## 2.4 PSD: Cesàro Sum

For a real sequence  $\{a_n\}_{n \geq 1}$ , let  $\sigma_n = \frac{a_1 + a_2 + \dots + a_n}{n}$ .

**D1) Suppose  $\lim_{n \rightarrow \infty} a_n = a$ , prove that  $\lim_{n \rightarrow \infty} \sigma_n = a$ .**

Proof: For any  $n > 0$ ,

$$|\sigma_n - a| \leq \sum_{i=1}^N \frac{|a_i - a|}{n} + \sum_{i=N+1}^n \frac{|a_i - a|}{n} \leq \frac{MN}{n} + \varepsilon(N),$$

where  $M = |a| + \sup_{i \leq N} |a_i|$ , and  $\varepsilon(N) = \sup_{i > N} |a_i - a|$ . By  $\lim_{n \rightarrow \infty} a_n = a$  we know  $\varepsilon(N) \rightarrow 0$ , hence  $\lim_{n \rightarrow \infty} \sigma_n = a$ .

**D2) Construct a divergent sequence  $\{a_n\}$  such that  $\lim_{n \rightarrow \infty} \sigma_n = 0$ .**

Solution:  $a_n = (-1)^{n-1}$ ,  $\sigma_n \in [0, 1/n]$ .

**D3) Determine whether there exists  $\{a_n\}_{n \geq 1}$  such that for any  $n \geq 1$ ,  $a_n > 0$  and  $\limsup_{n \rightarrow \infty} a_n = \infty$  but  $\lim_{n \rightarrow \infty} \sigma_n = 0$ .**

Solution: Let

$$a_n = \begin{cases} 2^{-n}, & n \neq 2^k, \\ k, & n = 2^k. \end{cases}$$

Then  $\limsup_{n \rightarrow \infty} a_n = \infty$  and  $a_n > 0$ , but for any  $n$ , suppose  $n \in [2^{k-1}, 2^k]$ , then

$$\sigma_n \leq \frac{1}{n} \cdot \left(1 + \frac{k(k+1)}{2}\right) \leq \frac{k(k+1)}{2^{k-1}}.$$

Hence  $\lim_{n \rightarrow \infty} \sigma_n = 0$ .

**D4) For  $k \geq 1$ , denote  $b_k = a_{k+1} - a_k$ . Prove that for any  $n \geq 2$ ,  $a_n - \sigma_n = \sum_{k=1}^{n-1} kb_k/n$ .**

Proof:

$$\sum_{k=1}^{n-1} kb_k = \sum_{k=1}^{n-1} k(a_{k+1} - a_k) = (n-1)a_n - \sum_{k=1}^{n-1} a_k = n(a_n - \sigma_n).$$

**D5) Suppose  $\lim_{k \rightarrow \infty} kb_k = 0$  and  $\{\sigma_n\}_{n \geq 1}$  converges. Prove that  $\{a_n\}_{n \geq 1}$  also converges.**

Proof: By D1),  $\lim_{k \rightarrow \infty} kb_k = 0$  implies

$$\lim_{n \rightarrow \infty} a_n - \sigma_n = \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^{n-1} kb_k}{n} = \lim_{k \rightarrow \infty} kb_k = 0.$$

Therefore  $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \sigma_n$  exists.

**D6) Suppose  $\{kb_k\}_{k \geq 1}$  is bounded, i.e.  $b_k = O(k^{-1})$ , and  $\lim_{n \rightarrow \infty} \sigma_n = \sigma$ . Prove that  $\lim_{n \rightarrow \infty} a_n = \sigma$ .**

Proof: Note that for  $m < n$ ,

$$a_n - \sigma_n = \frac{m}{n-m}(\sigma_n - \sigma_m) + \frac{1}{n-m} \sum_{k=m+1}^n a_n - a_k.$$

Therefore since  $\sigma_n$  is a Cauchy sequence, and  $|a_n - a_k| \leq M(n-k)/k$ , we can choose  $n, m$  to show that  $\lim_{n \rightarrow \infty} a_n - \sigma_n = 0$ .

## 2.5 PSE: Definition of $\sqrt[n]{x}$ and $b^x$

**E1) Given  $n \in \mathbb{N}$  and  $x > 0$ , prove that if  $y_1, y_2 > 0$  satisfy  $y_1^n = x = y_2^n$ , then  $y_1 = y_2$ .**

Proof: Note that  $y_1^{n-1} + y_1^{n-2}y_2 + \cdots + y_2^{n-1} > 0$ , and

$$0 = y_1^n - y_2^n = (y_1 - y_2) \cdot (y_1^{n-1} + y_1^{n-2}y_2 + \cdots + y_2^{n-1}).$$

Hence  $y_1 = y_2$ .

**E2) Prove that if  $x > 0$ , then the set  $E(x) = \{t \in \mathbb{R} : t^n < x\}$  is non-empty and has an upper-bound.**

Proof: Note that  $0 \in E(x)$  and  $E(x)$  has the upper-bound  $\max\{1, x\}$ .

**E3) Prove that  $y = \sup E(x)$  satisfy  $y^n = x$  and  $y > 0$ .**

Proof:  $y = \sup E(x) \implies y^n = x$  since  $t^n$  is continuous on  $\mathbb{R}$ , and  $y^n = x$  and  $0 \in E(x)$  implies  $y > 0$ .

**E4) Prove that the mapping  $\sqrt[n]{\cdot} : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}, x \mapsto \sqrt[n]{x} = y$  is well-defined. Denote  $\sqrt[n]{x}$  as  $x^{1/n}$ .**

Proof: Use E3).

**E5) Prove the  $\sqrt[n]{\cdot} : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$  is a bijection.**

Proof: By E1) it is injective, and  $\sqrt[n]{y^n} = y$  implies it is surjective. Hence it is a bijection.

**E6)  $a, b > 0$ ,  $n \in \mathbb{N}$ , prove that  $(ab)^{1/n} = a^{1/n}b^{1/n}$ .**

Proof: Use E5) and  $(xy)^n = x^n y^n$ .

**E7) Suppose  $b > 1$ ,  $m, n, p, q \in \mathbb{Z}$  where  $n, q > 0$ . Let  $r = \frac{m}{n} = \frac{p}{q}$ , prove that**

$$(b^m)^{1/n} = (b^p)^{1/q}.$$

Proof: Use  $(b^m)^q = (b^p)^n$  and E5).

**E8) Prove that for any  $r \in \mathbb{Q}$ ,  $r \mapsto b^r$  is well-defined.**

Proof: For  $r = p/q$ , where  $q > 0$ ,  $\gcd(p, q) = 1$ , let  $b^r = (b^p)^{1/q}$ , then for any  $r = m/n$ ,  $b^r = (b^m)^{1/n}$ .

**E9) Prove that for  $r, s \in \mathbb{Q}$ ,  $b^{r+s} = b^r b^s$ .**

Proof: Suppose  $r = p/q, s = m/n$ , where  $n, q > 0$ , then

$$b^{r+s} = b^{(mq+np)/nq} = (b^{mq} \cdot b^{np})^{1/nq} = (b^m)^{1/n} \cdot (b^p)^{1/q} = b^r b^s.$$

**E10) For  $x \in \mathbb{R}$ , let  $B(x) = \{b^t : t \in \mathbb{Q}, t \leq x\}$ . Prove that  $B(x)$  is non-empty and has an upper-bound. Define  $b^x = \sup B(x)$ .**

Proof:  $B(x)$  is clearly non-empty and bounded by  $b^{\lfloor x \rfloor + 1}$ .

**E11) Prove that if  $r \in \mathbb{Q}$ , then**

$$b^r = \sup B(r), \forall r \in \mathbb{Q}.$$

Proof:  $b^r \in B(r)$  and since  $b^t$  is monotonically increasing,  $b^r \geq \sup B(r)$ , hence  $b^r = \sup B(r)$ .



**E12) Prove that for any  $x, y \in \mathbb{R}$ ,  $b^{x+y} = b^x b^y$ .**

Proof: For any  $b^t \in B(x)$ ,  $b^s \in B(y)$ ,  $t \leq x$  and  $s \leq y$ , so  $t + s \leq x + y$  and  $b^{t+s} \in B(x + y)$ , hence  $b^{x+y} \geq b^x b^y$ . For any  $b^t \in B(x + y)$ ,  $t$  can be written in the form  $t = u + v$  where  $b^u \in B(x)$ ,  $b^v \in B(y)$ , so  $b^{x+y} \leq b^x b^y$ .

**E13\*) Prove that when  $b = e$ , the function derived from E10) (denoted as  $e^x$ ) is the same as  $\exp(x) = \sum_{n=0}^{\infty} x^n/n!$ .**

Proof: From  $\exp(1) = e$ ,  $\exp(0) = 1$  and  $\exp(x + y) = \exp(x) \cdot \exp(y)$  we know that for  $n \in \mathbb{Z}$ ,  $\exp(n) = e^n$ . For  $r = p/q \in \mathbb{Q}$ ,

$$(e^r)^q = e^p = \exp(p) = \exp(r)^q,$$

so by E5)  $e^r = \exp(r)$ . Since  $\exp$  is continuous, for any  $x \in \mathbb{R}$ ,  $e^x = \exp(x)$ .

## 2.6 PSF

Given  $\alpha > 0$  and  $x_1 > \sqrt{\alpha}$ , we define inductively  $\{x_n\}_{n \geq 1}$ :

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{\alpha}{x_n} \right), n \geq 1.$$

**F1) Prove that  $\{x_n\}$  is monotonically decreasing and  $\lim_{n \rightarrow \infty} x_n = \sqrt{\alpha}$  (which is defined in E).**

Proof: Note that

$$x_{n+1} - x_n = \frac{\alpha - x_n^2}{2x_n}.$$

Hence we can prove by induction that  $x_n > \sqrt{\alpha}$  and  $x_n > x_{n+1}$ .  $x_n$  is decreasing and bounded, so  $\lim_{n \rightarrow \infty} x_n = A$  exists, and  $A = (A + \alpha/A)/2$ . Therefore  $\lim_{n \rightarrow \infty} x_n = A = \sqrt{\alpha}$ .

**F2) Let  $\varepsilon_n = x_n - \sqrt{\alpha}$ . Prove that  $\varepsilon_{n+1} = \frac{\varepsilon_n^2}{2x_n} < \frac{\varepsilon_n^2}{2\sqrt{\alpha}}$ .**

Proof:

$$\frac{\varepsilon_n^2}{2x_n} = \frac{x_n^2 + \alpha - 2x_n\sqrt{\alpha}}{2x_n} = \frac{1}{2} \left( x_n + \frac{\alpha}{x_n} \right) - \sqrt{\alpha} = x_{n+1} - \sqrt{\alpha} = \varepsilon_{n+1}.$$

**F3) Prove that if  $\beta = 2\sqrt{\alpha}$ , then  $\varepsilon_{n+1} < \beta(\varepsilon_1/\beta)^{2^n}$ .**

Proof:  $\varepsilon_{n+1}/\beta < (\varepsilon_n/\beta)^2$ , hence  $\varepsilon_{n+1} < \beta(\varepsilon_1/\beta)^{2^n}$ .

**F4) Let  $\alpha = 3, x_1 = 2$ . Verify that  $\varepsilon_1/\beta < 0.1$ ,  $\varepsilon_5 < 4 \cdot 10^{-16}$ ,  $\varepsilon_6 < 4 \cdot 10^{-32}$ .**

Now we consider  $\alpha > 1$  and  $y_1 > \sqrt{\alpha}$ , and define

$$y_{n+1} = \frac{\alpha + y_n}{1 + y_n} = y_n + \frac{\alpha - y_n^2}{1 + y_n}, n \geq 1$$

**F6) Prove that  $\{y_{2k-1}\}$  is monotonically decreasing.**

Proof: Note that

$$y_{n+2} = \frac{\alpha + y_{n+1}}{1 + y_{n+1}} = \frac{\alpha + \frac{\alpha + y_n}{1 + y_n}}{1 + \frac{\alpha + y_n}{1 + y_n}} = \frac{2\alpha + (\alpha + 1)y_n}{(\alpha + 1) + 2y_n}$$

hence

$$y_{n+2} - y_n = \frac{2(\alpha - y_n^2)}{(\alpha + 1) + 2y_n}, \quad y_{n+2} - \sqrt{\alpha} = \frac{(\sqrt{\alpha} - 1)^2}{(\alpha + 1) + 2y_n} (y_n - \sqrt{\alpha}).$$

Therefore  $y_1 > \sqrt{\alpha}$  implies  $\sqrt{\alpha} < y_{2n+1} < y_{2n-1}$ .

**F7) Prove that  $\{y_{2k}\}$  is monotonically increasing.**

Proof:  $y_2 = (\alpha + y_1)/(1 + y_1) < \sqrt{\alpha}$ , so same as F6),  $y_{2k} > y_{2k-2}$  and  $y_{2k} < \sqrt{\alpha}$ .

**F8) Prove that  $\lim_{n \rightarrow \infty} y_n = \sqrt{\alpha}$ .**

Proof:  $\{y_{2n-1}\}$  is decreasing and bounded by  $\sqrt{\alpha}$ , so  $\lim_{n \rightarrow \infty} y_{2n-1} = A$  exists and  $A = (2\alpha + (\alpha + 1)A)/((\alpha + 1) + 2A)$ , so  $A = \sqrt{\alpha}$ . Likewise  $\lim_{n \rightarrow \infty} y_{2n} = \sqrt{\alpha}$ , hence  $\lim_{n \rightarrow \infty} y_n = \sqrt{\alpha}$ .

**F9) Compare the rates of convergence between  $x_n$  and  $y_n$ .**

Solution: Let  $\delta_n = |y_n - \sqrt{\alpha}|$ , then  $\delta_n \sim c^n \delta_1$ , hence  $x_n$  converges faster than  $y_n$ .

## 2.7 PSG: Banach-Mazur Game

Alice and Bob are playing a game: Alice selects a closed interval  $W_1$  first, then Bob choose a subinterval  $L_1$  of  $W_1$ , such that the length of  $L_1$  is less than half of the length of  $W_1$ ; they take turns choosing intervals  $W_n$  and  $L_n$ , such that  $L_n \subset W_n \subset L_{n-1}$  and  $|L_n| < |W_n|/2 < |L_{n-1}|/4$ , obtaining

$$W_1 \supset L_1 \supset W_2 \supset L_2 \supset \cdots \supset W_n \supset L_n \supset \cdots$$

Alice and Bob find that

$$\bigcap_{n \geq 1} W_n = \bigcap_{n \geq 1} L_n = \{x\}$$

is a real number. If  $x \in \mathbb{Q}$  then Alice wins, otherwise Bob wins. Who has a winning strategy?

**Solution:** Bob will win. We show that if  $\mathbb{Q}$  is replaced with any set  $M$  that is of first category, Bob can still win.

$M$  can be written as the union of a countable number of nowhere dense sets. Then in every move of Bob, he can choose  $L_n$  such that it does not intersect the  $n$ th such nowhere dense set. Hence the final number  $x$  is not in  $M$ .

## 2.8 Problem H

Consider the set  $\mathcal{P} = \{\{p_n\}_{n \geq 1} : p_n \in \mathbb{Z}, p_1 \geq 2, p_{n+1} \geq p_n^2\}$ .

**H1) For any  $p = \{p_n\}_{n \geq 1} \in \mathcal{P}$ , define the sequence**

$$a_n = \prod_{k=1}^n \left(1 + \frac{1}{p_k}\right).$$

Prove that  $f(p) = \lim_{n \rightarrow \infty} a_n$  exists and  $f(p) \in (1, 2]$ .

Proof: Note that  $p_n \geq p_1^{2^{n-1}}$ , then

$$a_n \leq \prod_{k=1}^n \left(1 + \frac{1}{p_1^{2^{k-1}}}\right) = \frac{1 - p_1^{-2^n}}{1 - p_1^{-1}} < \frac{1}{1 - p_1^{-1}}.$$

So the sequence  $\{a_n\}$  is monotonic and bounded, hence  $f(p) = \lim_{n \rightarrow \infty} a_n$  exists. Since  $a_n \in (1 + 1/p_1, \frac{1}{1-p_1^{-1}})$ , we obtain  $f(p) \in [1 + 1/p_1, \frac{1}{1-p_1^{-1}}] \subset (1, 2]$ .

**H2) Prove that  $f : \mathcal{P} \rightarrow (1, 2]$  is a bijection.**

Proof: For any  $p = \{p_n\}, q = \{q_n\} \in \mathcal{P}$ , if  $p \neq q$ , take the least  $k$  such that  $p_k \neq q_k$  and suppose  $q_k \geq p_k + 1$ , then for any  $n > k$ ,

$$\begin{aligned} a_n &= \prod_{t=1}^n \left(1 + \frac{1}{p_t}\right) \geq \prod_{t=1}^k \left(1 + \frac{1}{p_t}\right) \cdot \left(1 + \frac{1}{p_{k+1}}\right) \\ b_n &= \prod_{t=1}^n \left(1 + \frac{1}{q_t}\right) \leq \prod_{i=1}^{k-1} \left(1 + \frac{1}{p_i}\right) \cdot \frac{1 - q_k^{-2^{n-k}}}{1 - q_k^{-1}} \end{aligned}$$

Therefore

$$b_n \leq \prod_{t=1}^k \left(1 + \frac{1}{p_t}\right) \leq (1 + C)a_n$$

for all  $n > k$  where  $C = p_{k+1}^{-1} > 0$ , hence  $f(q) \leq (1 + C)f(p) < f(p)$ , hence  $f$  is injective.

For any  $x \in (1, 2]$ , inductively define  $p = \{p_n\} \in \mathcal{P}$  as follows: For any  $n \geq 1$ , Let  $t$  be the least integer such that  $a_n \leq x$  and  $t \geq p_{n-1}^2$  (clearly such  $t$  exists). If  $a_n = x$ , then let  $p_n = t - 1$ ,  $p_m = p_n^{2^{m-n}}$  for all  $m > n$ , then  $f(p) = x$ . Otherwise let  $p_n = t$ . Note that for any  $n$  such that  $p_n > p_{n-1}^2$ ,

$$|x - a_n| \leq 2^{-2^n},$$

therefore  $f(p) = x$ , and  $f$  is surjective.

**H3) Prove that  $\mathcal{P}$  is uncountable.**

Proof: By H2) and the fact that  $(1, 2]$  is uncountable.

## 2.9 Problem I: Binary Expansion

Consider the set  $\mathcal{S} = \{\{s_n\}_{n \geq 0} : s_n \in \{-1, 1\}\}$ .

**I1) For any  $s = \{s_n\}_{n \geq 0} \in \mathcal{S}$ , define the sequence**

$$c_n = \sum_{k=0}^n \frac{s_0 s_1 \cdots s_k}{2^k}.$$

Prove that  $h(s) = \lim_{n \rightarrow \infty} c_n$  exists and  $h(s) \in [-2, 2]$ .

Proof:  $h(s)$  exists since  $c_n$  is clearly a Cauchy sequence, and  $c_n \in [-2, 2]$  hence  $h(s) \in [-2, 2]$ .

**I2) Prove that  $h : \mathcal{S} \rightarrow [-2, 2]$  is surjective. Determine whether  $h$  is injective.**

Proof: Consider any  $x \in [-2, 2]$ , we can choose  $s_n$  such that  $|c_n - x| \leq 2^{-n}$ . Hence there exists  $s = \{s_n\} \in \mathcal{S}$  such that  $h(s) = \lim_{n \rightarrow \infty} c_n = x$ , so  $h$  is surjective.

Consider  $s = \{1, -1, 1, 1, 1, \dots\} \in \mathcal{S}$  and  $s' = \{-1, -1, 1, 1, \dots\}$ , then  $h(s) = h(s') = 0$ , hence  $h$  is not injective.

**I3) For  $s = \{s_n\}_{n \geq 0} \in \mathcal{S}$ , prove that**

$$2 \sin\left(\frac{\pi}{4} c_n\right) = s_0 \sqrt{2 + s_1 \sqrt{2 + \dots + s_n \sqrt{2}}}.$$

Proof: We prove by induction on  $n$ . The base  $n = 0$  is trivial. If the statement holds for  $n - 1$ , then let  $s' = \{s_{n+1}\}_{n \geq 0} \in \mathcal{S}$ , we have

$$2 \sin\left(\frac{\pi}{4} c_n\right) = 2 \sin s_0 \left(\frac{\pi}{4} + \frac{1}{2} \cdot \frac{\pi}{4} c'_{n-1}\right) = s_0 \sqrt{2 + \sin\left(\frac{\pi}{4} c'_{n-1}\right)}.$$

By the induction hypothesis, the statement also holds for  $n$ .

**I4) Calculate the limit**

$$\lim_{n \rightarrow \infty} \sqrt{2 + \sqrt{2 + \dots + \sqrt{2}}}$$

Solution: Consider  $s = \{s_n = 1\}_{n \geq 0} \in \mathcal{S}$ , then  $c_n = 2 - 2^n$  hence  $\lim_{n \rightarrow \infty} 2 \sin(\pi c_n / 4) = 2$ .

## 2.10 Problem J

**Problem:  $k \geq 2$  is a given integer. Define the sequence  $\{a_n\}$  as follows:**

$$a_0 > 0 \text{ already given, } a_{n+1} = a_n + a_n^{-1/k}, n \geq 0.$$

Calculate  $\lim_{n \rightarrow \infty} a_n^{k+1} / n^k$ .

Solution: It is easy to see that  $a_n \rightarrow \infty$ , hence

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n^{\frac{k+1}{k}}}{n} &= \lim_{n \rightarrow \infty} a_n^{\frac{k+1}{k}} - a_n^{\frac{k+1}{k}} = \lim_{n \rightarrow \infty} a_n^{\frac{k+1}{k}} \left( \left( 1 + \frac{a_{n+1} - a_n}{a_n} \right)^{\frac{k+1}{k}} - 1 \right) \\ &= \lim_{n \rightarrow \infty} a_n^{\frac{k+1}{k}} \left( \left( 1 + a_n^{-\frac{k+1}{k}} \right)^{\frac{k+1}{k}} - 1 \right) = \frac{k+1}{k}. \end{aligned}$$

Therefore

$$\lim_{n \rightarrow \infty} \frac{a_n^{k+1}}{n^k} = \left( 1 + \frac{1}{k} \right)^k.$$

## 3 Homework 3: Basel Problem

### 3.1 PSA

**A1) Given  $f : (a, x_0) \cup (x_0, b) \rightarrow \mathbb{R}$ , then  $\lim_{x \rightarrow x_0} f(x)$  exists iff for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that for any  $x_1, x_2 \in (x_0 - \delta, x_0 + \delta)$ ,  $|f(x_1) - f(x_2)| < \varepsilon$ .**

Proof:  $\Leftarrow$  Let  $x_n = x_0 + 1/n$ , then  $\{f(x_n)\}$  form a Cauchy sequence, hence  $f(x_0) = \lim_{n \rightarrow \infty} f(x_n)$  exists. For any  $\varepsilon > 0$ , there exists  $N, \delta > 0$  such that for any  $x, y \in (x_0 - \delta, x_0 + \delta)$ ,  $|f(x) - f(y)| < \varepsilon$

and for any  $n > N$ ,  $|f(x_n) - f(x_0)| < \varepsilon$ , hence let  $\delta' = \min\{\delta, 1/N\}$ , then for any  $x \in (x_0 - \delta', x_0 + \delta')$ ,  $|f(x) - f(x_0)| \leq |f(x) - f(x_N)| + |f(x_N) - f(x_0)| < 2\varepsilon$ .

Hence  $\lim_{x \rightarrow x_0} f(x) = f(x_0)$  exists.

$\implies$  For any  $\varepsilon > 0$  there exists  $\delta > 0$  such that for any  $x \in (x_0 - \delta, x_0 + \delta)$ ,  $|f(x) - f(x_0)| < \varepsilon$ , hence for any  $x, y \in (x_0 - \delta, x_0 + \delta)$ ,  $|f(x) - f(y)| < 2\varepsilon$ .

**A2) Suppose  $I$  is an interval (not a point), prove that the linear space  $C(I)$  on  $\mathbb{R}$  is of infinite dimension.**

Proof:  $C(I)$  contains the subspace of all polynomials, hence is of infinite dimension.

**A3) Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both continuous, prove that  $g \circ f : X \rightarrow Z$  is also continuous.**

Proof: For any open set  $U \in Z$ ,  $g^{-1}(U) \subset Y$  is an open set, and  $f^{-1}(g^{-1}(U)) \subset X$  is an open set, hence  $(g \circ f)^{-1}(U)$  is an open set in  $X$  and therefore  $g \circ f$  is continuous on  $X$ .

**A4) Suppose  $(X, d_X)$  and  $(Y, d_Y)$  are metric spaces,  $f : X \rightarrow Y$  is continuous. If  $d'_X$  and  $d_X$  are equivalent metrics, and so are  $d'_Y$  and  $d_Y$ , then in the spaces  $(X, d'_X)$  and  $(Y, d'_Y)$ ,  $f$  is also continuous.**

Proof: The topology generated by equivalent metrics are the same.

**A5) The mapping  $f : X \rightarrow \mathbb{R}^n$  can be written in the form**

$$f : X \rightarrow \mathbb{R}^n, x \mapsto f(x) = (f_1(x), f_2(x), \dots, f_n(x)).$$

Prove that  $f$  is continuous iff  $f_i$  is continuous for every  $i = 1, 2, \dots, n$ .

Proof: Since  $f$  is continuous iff  $\forall x_n \rightarrow x, f(x_n) \rightarrow f(x)$ , and  $\{x_k = (x_k^{(1)}, \dots, x_k^{(n)})\}_{k \geq 1}$  converges iff every  $\{x_k^{(i)}\}_{k \geq 1}$  converges,  $f$  is continuous iff every  $f_i$  is continuous.

**A6) Suppose  $(X, d_X)$  is a metric space,  $(V, \|\cdot\|)$  is a normed linear space.  $f : X \rightarrow V$  and  $g : X \rightarrow V$  are continuous mappings. Prove that  $f \pm g : X \rightarrow V$  is continuous. If  $V = \mathbb{C}$  then  $f \cdot g : X \rightarrow \mathbb{C}$  is continuous. If  $V = \mathbb{C}$  and for any  $x \in X$ ,  $g(x) \neq 0$ , then  $f/g : X \rightarrow \mathbb{C}$  is continuous.**

(Choose one statement to prove.)

Proof: Since for  $\{x_n\}, \{y_n\} \subset \mathbb{C}$ ,  $\lim_{n \rightarrow \infty} x_n y_n = \lim_{n \rightarrow \infty} x_n \cdot \lim_{n \rightarrow \infty} y_n$  and if  $y_n \neq 0$ , then

$$\lim_{n \rightarrow \infty} x_n / y_n = \lim_{n \rightarrow \infty} x_n / \lim_{n \rightarrow \infty} y_n.$$

Hence  $f \cdot g, f/g$  are both continuous.

For  $\{x_n\}, \{y_n\} \subset V$ , if  $A = \lim_{n \rightarrow \infty} x_n$  and  $B = \lim_{n \rightarrow \infty} y_n$  then

$$\|x_n + y_n - A - B\| \leq \|x_n - A\| + \|y_n - B\| \rightarrow 0.$$

Hence  $f \pm g$  is continuous.

**A7) Find all discontinuities of the function**

$$f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto \begin{cases} 1/q, & \text{if } x = p/q \in \mathbb{Q}, \text{ where } q \geq 1, (p, q) = 1. \\ 0, & \text{if } x \notin \mathbb{Q}. \end{cases}$$

### 3 Homework 3: Basel Problem

---

Solution: For any  $x \in \mathbb{Q}$ ,  $f(x) \neq 0$  but for any  $\delta > 0$  there exists  $y \in (x - \delta, x + \delta)$  such that  $y \notin \mathbb{Q}$ . Hence  $|f(x) - f(y)| = f(x)$ , so  $f$  is not continuous at  $x$ .

For any  $x \notin \mathbb{Q}$ , and any  $\varepsilon > 0$ , let  $N = \lfloor 1/\varepsilon \rfloor + 1$  and  $\delta = \inf_{n \leq N} \|xn\|/n$ , then for any  $y \in (x - \delta, y + \delta)$ , if  $y \notin \mathbb{Q}$  then  $f(x) = f(y) = 0$ , if  $y = p/q \in \mathbb{Q}$  then  $q > N > 1/\varepsilon$ , hence  $|f(x) - f(y)| = f(y) = 1/q < \varepsilon$ . Therefore  $f$  is continuous at  $x$  iff  $x \notin \mathbb{Q}$ .

#### A8) Calculate

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = \lim_{x \rightarrow 0} \sum_{n=1}^{\infty} \frac{x^{n-1}}{n!} = 1.$$

#### A9) Calculate

$$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e.$$

Since  $\lim_{n \rightarrow \infty} (1 + 1/n)^n = e$  and  $(1 + 1/x)^x$  is monotonic on  $[100, \infty)$ .

#### A10) Calculate

$$\lim_{x \rightarrow -\infty} \left(1 + \frac{1}{x}\right)^x = e.$$

Since  $\lim_{x \rightarrow \infty} (1 - 1/x)^x = \lim_{x \rightarrow \infty} (1 - 1/x)^{x-1} = e$ .

### 3.2 PSB

#### B1) Calculate the following series:

1.

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \sum_{n=1}^{\infty} \frac{1}{n} - \frac{1}{n+1} = 1.$$

2.

$$\sum_{n=1}^{\infty} \frac{1}{4n^2 - 1} = \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{2n-1} - \frac{1}{2n+1} = \frac{1}{2}.$$

3.

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)(n+2)} = \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n(n+1)} - \frac{1}{(n+1)(n+2)} = \frac{1}{4}.$$

4.

$$\sum_{n=1}^{\infty} \arctan \frac{1}{n^2 + n + 1} = \sum_{n=1}^{\infty} \arctan \frac{1}{n} - \arctan \frac{1}{n+1} = \frac{\pi}{4}.$$

5.

$$\sum_{n=0}^{\infty} \frac{(-1)^n + 2}{3^n} = \frac{1}{1 + 1/3} + \frac{2}{1 - 1/3} = \frac{3}{4} + 3 = \frac{15}{4}.$$

6.

$$\sum_{n=1}^{\infty} \frac{n}{3^n} = \frac{3}{4}.$$

7.

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{2^{n-1}} = \frac{1}{1+1/2} = \frac{2}{3}.$$

8.

$$\sum_{n=1}^{\infty} \frac{2n-1}{2^n} = 3.$$

9.

$$\sum_{n=1}^{\infty} \frac{2n+1}{n^2(n+1)^2} = \sum_{n=1}^{\infty} \frac{1}{n^2} - \frac{1}{(n+1)^2} = 1.$$

10.

$$\sum_{n=1}^{\infty} \sqrt{n+2} - 2\sqrt{n+1} + \sqrt{n} = 1 - \sqrt{2}.$$

11.

$$\sum_{n=1}^{\infty} \log \left( \frac{n(2n+1)}{(n+1)(2n-1)} \right) = \lim_{n \rightarrow \infty} \log \left( \frac{2n+1}{n+1} \right) = \log 2.$$

12.

$$\sum_{n=1}^{\infty} \frac{1}{n(n+m)} = \frac{1}{m} \sum_{n=1}^{\infty} \frac{1}{n} - \frac{1}{n+m} = \frac{1}{m} \sum_{n=1}^m \frac{1}{n}.$$

**B2) Determine whether the following series converge:**

1.

$$\sum_{n=1}^{\infty} \sqrt{n+1} - \sqrt{n} = \lim_{n \rightarrow \infty} \sqrt{n+1} - 1 = \infty.$$

2.

$$\sum_{n=1}^{\infty} \frac{\sqrt{n+1} - \sqrt{n}}{n} \leq \sum_{n=1}^{\infty} \frac{1}{2n\sqrt{n}}$$

converges.

3.

$$\sum_{n=2}^{\infty} (\sqrt[n]{n} - 1)^n$$

converges, since  $\limsup_{n \rightarrow \infty} \sqrt[n]{(\sqrt[n]{n} - 1)^n} = 0 < 1$ .

4.

$$\sum_{n=1}^{\infty} \frac{1}{1+x^n}$$

converges if  $|x| > 1$  and diverges if  $|x| \leq 1$ .

5.

$$\sum_{n=1}^{\infty} \frac{1}{n2^n} \leq \sum_{n=1}^{\infty} \frac{1}{2^n} = 1.$$

converges

6.

$$\sum_{n=1}^{\infty} \left( \frac{n^2}{3n^2+1} \right)^n \leq \sum_{n=1}^{\infty} \frac{1}{3^n} < 1.$$

converges.

7.

$$\sum_{n=1}^{\infty} \frac{1}{n^{1+1/n}} \geq \sum_{n=1}^{\infty} \frac{1}{2n}$$

diverges.

8.

$$\sum_{n=2}^{\infty} \frac{1}{(\log n)^{\log n}} = \sum_{n=2}^{\infty} \frac{1}{n^{\log \log n}} \leq C + \sum_{n=100}^{\infty} \frac{1}{n^2}$$

converges.

9.

$$\sum_{n=1}^{\infty} \frac{n^{n+1/n}}{(n + \frac{1}{n})^n}$$

diverges, since

$$\lim_{n \rightarrow \infty} \frac{n^{n+1/n}}{(n + \frac{1}{n})^n} = \exp \lim_{n \rightarrow \infty} \left( \frac{\log n}{n} - n \log \left( 1 + \frac{1}{n^2} \right) \right) = 1.$$

10.

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1} \sqrt{n}}{n+1}$$

converges (conditionally), since the partial sum of  $(-1)^{n-1}$  is bounded and  $\frac{\sqrt{n}}{n+1}$  monotonically tends to 0.

11.

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt[n]{n}}$$

diverges since  $(-1)^{n-1} n^{-1/n}$  does not tend to 0.

12.

Let  $H_n = 1 + 1/2 + \dots + 1/n$ .

$$\sum_{n=1}^{\infty} \frac{H_n \sin nx}{n}$$

converges since the partial sum of  $\sin nx$  is bounded and  $\frac{H_n}{n}$  monotonically tends to 0.

**B3) Determine whether the following series converge (absolutely):**

1.

$$\sum_{n=2}^{\infty} \frac{(-1)^n}{n \log n}$$

converges since the partial sum of  $(-1)^n$  is bounded and  $\frac{1}{n \log n}$  monotonically tends to 0, but only conditionally by C3).

2.

$$\sum_{n=2}^{\infty} \frac{\sin(n\pi/4)}{\log n}$$

converges since the partial sum of  $\sin(n\pi/4)$  is bounded and  $\frac{1}{\log n}$  monotonically tends to 0, but only conditionally since  $\sum_{n=2}^{\infty} \frac{1}{\log(4n+2)}$  tends to infinity.

3.

$$\sum_{n=1}^{\infty} (-1)^n \frac{n-1}{n+1} \frac{1}{\sqrt[3]{n}}$$



converges since  $\frac{n-1}{(n+1)^{\frac{1}{3}}n}$  monotonically tends to 0, but only conditionally since  $\sum_{n=1}^{\infty} n^{-1/3}$  diverges.

4.  
 $a > 1$ .

$$\sum_{n=1}^{\infty} (-1)^{n(n-1)/2} \frac{n^{10}}{a^n}$$

converges absolutely since there exists  $C > 0$  such that for  $n > C$ ,  $n^{10}a^{-n} \leq a^{-n/2}$ , and  $\sum_{n=1}^{\infty} a^{-n/2}$  converges.

### 3.3 PSC

Suppose the integer  $b \geq 2$ ,  $f : [1, \infty) \rightarrow \mathbb{R}_{>0}$  is monotonically decreasing.

**C1) Prove that**

$$(b-1)b^{k-1}f(b^k) \leq \sum_{j=b^{k-1}}^{b^k-1} f(j) \leq (b-1)b^{k-1}f(b^{k-1}).$$

Proof: There are  $(b-1)b^{k-1}$  integers in  $[b^{k-1}, b^k - 1]$ , and since  $f$  is monotonically decreasing, for any  $j \in [b^{k-1}, b^k - 1]$ ,  $f(j) \in [f(b^k), f(b^{k-1})]$ .

**C2) Prove that the series**

$$\sum_{n=1}^{\infty} f(n) \text{ and } \sum_{n=1}^{\infty} b^n f(b^n)$$

converge or diverge simultaneously.

Proof: From C1),

$$\sum_{k=1}^{\infty} (b-1)b^{k-1}f(b^k) \leq \sum_{n=1}^{\infty} f(n) = \sum_{k=1}^{\infty} \sum_{j=b^{k-1}}^{b^k-1} f(j) \leq \sum_{k=1}^{\infty} (b-1)b^{k-1}f(b^{k-1}).$$

Therefore the two series converge or diverge simultaneously.

**C3) Prove that  $\sum_{n=2}^{\infty} \frac{1}{n \log n}$  diverges.**

Proof: Consider  $f(x) = \frac{1}{x \log x}$  which is monotonically decreasing. Note that

$$\sum_{n=2}^{\infty} 2^n f(2^n) = \sum_{n=2}^{\infty} \frac{1}{n \log 2} = \infty.$$

From C2) we know that  $\sum_{n=2}^{\infty} f(n)$  diverges.

**C4) Prove that  $\sum_{n=100}^{\infty} \frac{1}{n \log n \log \log n}$  diverges.**

Proof: Consider  $f(x) = \frac{1}{x \log x \log \log x}$  which is monotonically decreasing. From C3),

$$\sum_{n=100}^{\infty} 2^n f(2^n) = \sum_{n=100}^{\infty} \frac{1}{n \log 2 \cdot \log(n \log 2)}$$

diverges. Hence from C2) we know that  $\sum_{n=100}^{\infty} f(n)$  diverges.

**C5) Prove that  $\sum_{n=1}^{\infty} n^{-s}$  converges iff  $s > 1$ .**

Proof: Consider  $f(x) = x^{-s}$  which is monotonically decreasing. Note that

$$\sum_{n=1}^{\infty} 2^n f(2^n) = \sum_{n=1}^{\infty} 2^{-n(s-1)} = \frac{2^{1-s}}{1 - 2^{1-s}}.$$

**C6) Suppose  $s > 1$ , prove that  $\sum_{n=2}^{\infty} \frac{1}{n(\log n)^s}$  and  $\sum_{n=10}^{\infty} \frac{1}{n \log n (\log \log n)^s}$  converges.**

Proof: Same as C3) and C4).

### 3.4 PSD

For  $\{a_n\}_{n \geq 1} \subset \mathbb{R}$ ,

- $\alpha \in \mathbb{R}$ , if for any  $\varepsilon > 0$ , there are infinitely many  $n$  such that  $a_n \in (\alpha - \varepsilon, \alpha + \varepsilon)$ , then we call  $\alpha$  a limit point of  $\{a_n\}_{n \geq 1}$ .
- Likewise define limit points for  $\alpha = \pm\infty$ .

**D1) Prove that  $\alpha \in \mathbb{R}$  is a limit point of  $\{a_n\}_{n \geq 1}$  iff there is a sub-sequence  $\{a_{n_k}\}_{k \geq 1}$  which converges to  $\alpha$ .**

Proof:  $\Leftarrow$  is trivial.  $\Rightarrow$  Let  $\varepsilon = 1/k$  then there exists  $a_{n_k}$  such that  $|a_{n_k} - \alpha| < \varepsilon$ . Hence  $\lim_{k \rightarrow \infty} a_{n_k} = \alpha$ .

**D2) Prove that  $+\infty$  is a limit point of  $\{a_n\}_{n \geq 1}$  iff there is a sub-sequence  $\{a_{n_k}\}_{k \geq 1}$  such that  $\lim_{k \rightarrow \infty} a_{n_k} = \infty$ .**

Proof: Same as D1).

**D3) Let  $E = \{\alpha \in \mathbb{R} \cup \{\pm\infty\} : \alpha \text{ is a limit point of } \{a_n\}\}$ . Prove that  $E \neq \emptyset$ .**

Proof: If  $\{a_n\}$  is unbounded, then by D2)  $E \cap \{\pm\infty\} \neq \emptyset$ . If  $\{a_n\}$  is bounded, then by Bolzano-Weierstrass theorem,  $E \neq \emptyset$ .

**D4) Prove that  $E \subset \mathbb{R}$  iff  $\{a_n\}$  is bounded.**

Proof: Use D2)

**D5) Suppose  $\{a_n\}_{n \geq 1}$  is bounded. Prove that  $\sup E = \limsup_{n \rightarrow \infty} a_n$ ,  $\inf E = \liminf_{n \rightarrow \infty} a_n$ .**

Proof: Let  $M = \limsup_{n \rightarrow \infty} a_n$ , then for any  $\varepsilon > 0$ , there exists  $n$  such that  $M - \varepsilon < \sup_{k \geq n} a_k < M + \varepsilon$ , hence there exists  $k \geq n$  such that  $|a_k - M| < \varepsilon$ , so  $M \in E$ .

For any  $\alpha \in E$ , there is a sub-sequence  $\{a_{n_k}\} \rightarrow \alpha$ , hence

$$\alpha = \lim_{k \rightarrow \infty} a_{n_k} \leq \lim_{k \rightarrow \infty} \sup_{m \geq n_k} a_{n_k} = \limsup_{n \rightarrow \infty} a_n = M.$$

Therefore  $M = \sup E$ . Substitute  $a_n \rightarrow -a_n$  and we obtain  $\inf E = \liminf_{n \rightarrow \infty} a_n$ .

**D6) Suppose  $\{a_n\}_{n \geq 1}$  is bounded. Let  $a^* = \limsup_{n \rightarrow \infty} a_n$ . Prove that**

i)  $a^* \in E$ , i.e.  $\sup E \in E$ .

Proof: See the proof of D5).

ii) For any  $x > a^*$ , there exists  $N \in \mathbb{Z}_{\geq 1}$  such that for any  $n > N$ ,  $a_n < x$ .

Proof: If there is an infinite sub-sequence  $\{a_{n_k}\}_{k \geq 1}$  such that  $a_{n_k} \geq x$ , then  $\{a_{n_k}\}$  has a limit point  $a' > x > a^*$ , contradicting  $a^* = \sup E$ .

**D7) Construct an example of  $\{a_n\}_{n \geq 1}$  such that  $E \cap \mathbb{R} \neq \emptyset$  and  $E \not\subset \mathbb{R}$ .**

Solution: Since  $\mathbb{Q}$  is countable, let  $\{a_n\}_{n \geq 1}$  iterate every element of  $\mathbb{Q}$ , then  $E = \mathbb{R} \cup \{\pm\infty\}$  is an infinite set.

**D8) Construct  $\{a_n\}_{n \geq 1}$  such that  $E$  is an infinite set.**

Solution: Same as D7).

### 3.5 PSE: Reciprocal Sum of Primes

Define the  $\zeta$ -function:

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s}.$$

We have proved the formula:

$$\zeta(s) = \prod_{p \in \mathcal{P}} \frac{1}{1 - p^{-s}}.$$

Prove that the series

$$\sum_{p \in \mathcal{P}} p^{-s}$$

converges when  $s > 1$ , and diverges when  $0 < s \leq 1$ .

Proof: We know that for  $|a_n| < 1$ ,  $\prod_{n=1}^{\infty} (1 - a_n)$  converges iff  $\sum_{n=1}^{\infty} a_n$  converges. Hence by  $\zeta(s)^{-1} = \prod_{p \in \mathcal{P}} (1 - p^{-s})$ , we obtain  $\sum_{p \in \mathcal{P}} p^{-s}$  converges iff  $s > 1$ .

### 3.6 PSF: Euler's "Proof" of the Basel Problem

For any  $\theta \in \mathbb{R}, n \in \mathbb{Z}$ , prove the identity

$$\frac{\sin((2n+1)\theta)}{(2n+1)\sin\theta} = \prod_{k=1}^n \left(1 - \frac{\sin^2\theta}{\sin^2(k\pi/(2n+1))}\right).$$

Further prove that for any  $x \in \mathbb{R}$ ,

$$\frac{\sin(\pi x)}{\pi x} = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2}\right).$$

Proof: (1) By induction there is a polynomial  $P_n(x)$  such that  $P_n(\sin\theta) = \sin(2n+1)\theta$  for any  $\theta \in \mathbb{R}$  and  $\deg P_n = 2n+1$ . For any  $k = 1, 2, \dots, n$ , and  $\theta = \pm k\pi/(2n+1)$ ,  $\sin((2n+1)\theta) = 0$ , hence  $P_n$  has roots 0 and  $\pm \sin(k\pi/(2n+1))$  for  $k = 1, 2, \dots, n$ . Since  $\deg P_n = 2n+1$ ,

$$P_n(x) = Cx \prod_{k=1}^n \left(1 - \frac{x^2}{\sin^2(k\pi/(2n+1))}\right)$$

for some  $C \in \mathbb{R}$ . Let  $x = \sin \theta$  and consider the derivatives on both sides when  $\theta = 0$ , then we obtain  $C = 2n + 1$ , therefore

$$\frac{\sin((2n+1)\theta)}{(2n+1)\sin\theta} = \prod_{k=1}^n \left(1 - \frac{\sin^2\theta}{\sin^2(k\pi/(2n+1))}\right).$$

(2) Let  $m = 2n + 1$ . From (1) we know that for any  $x \in \mathbb{C}$  and  $k < n$ ,  $\sin x = U_k^{(n)} \cdot V_k^{(n)}$ , where

$$U_k^{(n)} = m \sin \frac{x}{m} \prod_{j=1}^k \left(1 - \frac{\sin^2(x/m)}{\sin^2(j\pi/m)}\right),$$

$$V_k^{(n)} = \prod_{j=k+1}^n \left(1 - \frac{\sin^2(x/m)}{\sin^2(j\pi/m)}\right).$$

Clearly, for any  $k \in \mathbb{N}$ ,

$$\lim_{n \rightarrow \infty} U_k^{(n)} = U_k = x \prod_{j=1}^k \left(1 - \frac{x^2}{j^2\pi^2}\right).$$

and for any  $x \in \mathbb{C}$  and  $j \in \mathbb{N}$ ,

$$\left| \frac{\sin^2(x/m)}{\sin^2(j\pi/m)} \right| \leq \frac{x^2}{4j^2} \cdot K(|x|/m)^2,$$

where  $K(x) = \sum_{n=0}^{\infty} |x|^n / (2n+1)!$  is monotonic on  $[0, \infty)$  and  $K(0) = 1$ . Note that for  $\alpha_i \in \mathbb{C}$ ,

$$\left| 1 - \prod_{j=1}^n (1 - \alpha_n) \right| \leq \sum_{j=1}^n \left( \sum_{k=1}^n |\alpha_k| \right)^j.$$

Hence for any  $x \in \mathbb{C}$  and  $\varepsilon > 0$ , there exists  $N$  such that for any  $k \geq N$ , and any  $n > k$ ,  $|V_k^{(n)} - 1| < \varepsilon$ , since

$$|V_k^{(n)} - 1| \leq \sum_{j=1}^{\infty} \left( \sum_{l=k+1}^{\infty} \frac{x^2}{4l^2} K(|x|/m)^2 \right)^j \leq \sum_{j=1}^{\infty} \left( K(|x|/(2k+1))^2 \cdot \frac{x^2}{k} \right)^j \rightarrow 0.$$

i.e. for any  $x \in \mathbb{C}$

$$\lim_{k \rightarrow \infty} \sup_{n > k} |V_k^{(n)} - 1| = 0.$$

And likewise we know that there is a constant  $M$  such that for any  $n > k$ ,  $|x| < k$ ,  $|U_k^{(n)}| \leq M$ . Therefore for any  $x \in \mathbb{C}$ ,

$$\sin x = x \lim_{n \rightarrow \infty} \prod_{k=1}^n \left(1 - \frac{x^2}{k^2\pi^2}\right) = x \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2\pi^2}\right).$$

Note:

From the formula above, we can formally deduce that

$$\sin(\pi x) = \pi x (1 - \zeta(2)x^2 + \zeta(4)x^4 + \cdots).$$

Compare it to  $\sin z = x - x^3/6 + \cdots$ , and we get  $\zeta(2) = \pi^2/6$ .

## 4 Homework 4: Topology

### 4.1 PSA: Topology on Metric Spaces

**A1) Suppose  $(X, d_X)$  and  $(Y, d_Y)$  are metric spaces,  $f : X \rightarrow Y$  is a mapping. Prove that the two following definitions of continuity is equivalent:**

- Suppose  $x_0 \in X$ , if for any  $\{x_n\}_{n \geq 1} \subset X$  such that  $\lim_{n \rightarrow \infty} x_n = x_0$ , we have  $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$ , then we say  $f$  is continuous at  $x_0$ . If  $f$  is continuous at every point  $x \in X$ , then  $f$  is a continuous mapping.
- Suppose  $x_0 \in X$ ,  $y_0 = f(x_0) \in Y$ . If for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that for any  $d_X(x, x_0) < \delta$ ,  $x \in X$ , we have  $d_Y(f(x), f(x_0)) < \varepsilon$ , we call  $f$  continuous at  $x_0$ . If  $f$  is continuous at every point  $x \in X$ , then  $f$  is a continuous mapping.

Proof: 1 $\Rightarrow$ 2: If there exists  $\varepsilon > 0$  such that for any  $n \geq 1$ , there exists  $x_n$  such that  $d_X(x_0, x_n) < 1/n$  but  $d_Y(f(x_n), f(x_0)) > \varepsilon$ , then  $\lim_{n \rightarrow \infty} x_n = x_0$  but  $\lim_{n \rightarrow \infty} f(x_n) \neq f(x_0)$ , a contradiction.

2 $\Rightarrow$ 1: For any  $\{x_n\}_{n \geq 1} \subset X$  such that  $\lim_{n \rightarrow \infty} x_n = x_0$ , and any  $\varepsilon > 0$ , take the corresponding  $\delta$  and  $N$  such that  $n > N \implies d(x_n, x_0) < \delta$ . Then for any  $n > N$ ,  $d(x_n, x_0) < \delta$  so  $d(f(x_n), f(x_0)) < \varepsilon$ , hence  $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$ .

**A2)  $(X, d)$  is a metric space. For any  $x \in X$ ,  $r > 0$ , let  $B(x, r) = \{y \in X : d(x, y) < r\}$ . Prove that for any  $x \in X$ ,  $r > 0$ , if  $x' \in B(x, r)$ , then there exists  $r' > 0$  such that  $B(x', r') \subset B(x, r)$ .**

If  $U = \bigcup_{\alpha \in \mathcal{A}} B(x_\alpha, r_\alpha)$ , then we call  $U$  an open set. Prove that  $U \subset X$  is open iff for any  $x \in U$ , there exists  $\delta_x > 0$  such that  $B(x, \delta_x) \subset U$ .

Proof: If  $x' \in B(x, r)$ , let  $r' = r - d(x, x')$ , then for any  $y \in B(x', r')$ ,  $d(x, y) \leq d(x, x') + d(x', y) < d(x, x') + r' = r$ , hence  $y \in B(x, r)$  so  $B(x', r') \subset B(x, r)$ .

If for any  $x \in U$ , there exists  $\delta_x > 0$  such that  $B(x, \delta_x) \subset U$ , then  $U = \bigcup_{x \in U} B(x, \delta_x)$  is open.

If  $U$  is open then for any  $x \in U$ , suppose  $x \in B(x_\alpha, r_\alpha)$  for some  $\alpha \in \mathcal{A}$ , then there exists  $\delta_x > 0$  such that  $B(x, \delta_x) \subset B(x_\alpha, r_\alpha) \subset U$ .

**A3) Let  $\mathcal{T}$  denote all open sets on  $(X, d)$ . Prove that  $\mathcal{T}$  is a topology.**

Proof: 1.  $\emptyset \in \mathcal{T}$ ,  $X = \bigcup_{x \in X} B(x, 1) \in \mathcal{T}$ . 2. If  $\{U_\alpha : \alpha \in J\} \subset \mathcal{T}$ , where  $U_\alpha = \bigcup_{x \in \mathcal{A}_\alpha} B(x, r_{\alpha, x})$  then let  $\mathcal{A} = \bigcup_{\alpha \in J} \mathcal{A}_\alpha$ ,

$$\bigcup_{\alpha \in J} U_\alpha = \bigcup_{x \in \mathcal{A}} B(x, \sup_{\alpha, x \in \mathcal{A}_\alpha} r_{\alpha, x}) \in \mathcal{T}.$$

1. If  $U_1, \dots, U_n \in \mathcal{T}$ , where  $U_k = \bigcup_{x \in \mathcal{A}_k} B(x, r_{k, x})$ , then let  $\mathcal{A} = \bigcup_{k=1}^n \mathcal{A}_k$

$$\bigcap_{k=1}^n U_k = \bigcup_{x \in \mathcal{A}} B(x, \min_{k \in \{1, \dots, n\}} r_{k, x}) \in \mathcal{T}.$$

Therefore  $\mathcal{T}$  is a topology on  $X$ .

**A4)  $(X, d)$  is a metric space. If  $F \subset X$  and  $F^C$  is open, then we call  $F$  a closed set. Prove that  $F$  is closed iff for any sequence  $\{x_n\}_{n \geq 1} \in F$ , if  $\lim_{n \rightarrow \infty} x_n = x$  then  $x \in F$ .**

Proof: Suppose  $F$  is closed, if a sequence  $\{x_n\}_{n \geq 1}$  satisfy  $\lim_{n \rightarrow \infty} x_n = x$  and  $x \in F^C$ , then there exists  $\varepsilon > 0$  such that  $B(x, \varepsilon) \subset F^C$ . However  $B(x, \varepsilon) \cap \{x_n\} \neq \emptyset$ , leading to contradiction.

If for any sequence  $\{x_n\}_{n \geq 1}$  such that  $\lim_{n \rightarrow \infty} x_n = x$ , there is  $x \in F$ , then for any  $x \in F^C$ , if for any  $\varepsilon > 0$   $B(x, \varepsilon) \not\subset F^C$ , then for any  $n \geq 1$ , take  $x_n \in B(x, \varepsilon) \cap F$ . The sequence  $\{x_n\}$  has the limit  $\lim_{n \rightarrow \infty} x_n = x$  but  $x \in F^C$ , a contradiction. Hence  $F$  is closed.

**A5) Prove that**

1.  $\emptyset$  and  $X$  are closed sets.
  2. Any intersection of closed sets are still closed.
  3. Finite unions of closed sets are still closed.
- Proof: Use A3) and de Morgan's theorem.

**A6) Suppose  $(X, d_X)$  and  $(Y, d_Y)$  are metric spaces and  $f : X \rightarrow Y$ , then the following statements are equivalent:**

1.  $f$  is continuous.
2. For any  $U \subset Y$  open,  $f^{-1}(U)$  is an open set in  $X$ .
3. For any  $F \subset Y$  closed,  $f^{-1}(F)$  is a closed set in  $X$ .

Proof:  $1 \Rightarrow 2$ : If  $f$  is continuous, then for any  $U \subset Y$  open, consider any point  $x \in f^{-1}(U)$ . Let  $y = f(x) \in U$ , then there exists  $\varepsilon > 0$  such that  $B(y, \varepsilon) \subset U$ . Since  $f$  is continuous, there exists  $\delta > 0$  such that for any  $x' \in B(x, \delta)$ ,  $f(x') \in B(y, \varepsilon) \subset U$ , hence  $B(x, \delta) \subset f^{-1}(U)$ . Therefore  $f^{-1}(U)$  is an open set in  $X$ .

$2 \Rightarrow 1$ : For any  $x \in X$  and  $\varepsilon > 0$ , consider the open set  $U = B(y, \varepsilon)$ , where  $y = f(x)$ . Since  $x \in f^{-1}(U)$  and  $f^{-1}(U)$  is an open set, there exists  $\delta > 0$  such that  $B(x, \delta) \subset f^{-1}(U)$ , therefore  $f$  is continuous.

$2 \Leftrightarrow 3$ : Note that  $f^{-1}(Y \setminus A) = X \setminus f^{-1}(A)$ .

**A7) Let  $A'$  be the set of limit points of  $A$ . Prove that  $\bar{A} = A' \cup A$ .**

Proof: For any closed set  $F \supset A$ , by A4) we know  $A' \subset F$ , hence  $A' \cup A \subset \bar{A}$ . Consider a sequence  $\{x_n\}_{n \geq 1} \subset A' \cup A$  such that  $\lim_{n \rightarrow \infty} x_n = x$  exists, for any  $n \geq 1$  we can find a  $y_n \in A$  such that  $d(x_n, y_n) \leq 2^{-n}$ , hence  $\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} x_n = x$  so  $x \in A' \cup A$ . Therefore  $A' \cup A$  is closed, and hence  $\bar{A} = A' \cup A$ .

**A8) Suppose  $(Y, d_Y)$  and  $(Z, d_Z)$  are metric spaces, define the metric on  $Y \times Z$ :**

$$d_{Y \times Z} : (Y \times Z)^2 \rightarrow \mathbb{R}_{\geq 0}, ((y_1, z_1), (y_2, z_2)) \rightarrow \sqrt{d_Y(y_1, y_2)^2 + d_Z(z_1, z_2)^2}.$$

Prove that this defines a metric and the projection mappings are continuous:

$$\pi_Y : Y \times Z \rightarrow Y, (y, z) \mapsto y; \pi_Z : Y \times Z \rightarrow Z, (y, z) \mapsto z.$$

Given a mapping  $F : X \rightarrow Y \times Z$ , then  $F$  is continuous iff  $\pi_Y \circ F$  and  $\pi_Z \circ F$  are both continuous.

Proof:  $d((y_1, z_1), (y_2, z_2)) = 0 \iff (y_1, z_1) = (y_2, z_2)$ ,  $d((y_1, z_1), (y_2, z_2)) = d((y_2, z_2), (y_1, z_1))$ , and  $d((y_1, z_1), (y_2, z_2)) \leq d((y_1, z_1), (y_3, z_3)) + d((y_3, z_3), (y_2, z_2))$  (since  $\sqrt{(x+y)^2 + (u+v)^2} \leq \sqrt{x^2 + u^2} + \sqrt{y^2 + v^2}$ ), hence  $d_{Y \times Z}$  is a metric.

Note that  $d((y_1, z_1), (y_2, z_2)) \geq d(y_1, y_2)$ , hence  $\pi_Y$  and  $\pi_Z$  are continuous.

$d((y_1, z_1), (y_2, z_2)) \leq d(y_1, y_2) + d(z_1, z_2)$ , hence  $F$  is continuous iff  $\pi_Y \circ F$  and  $\pi_Z \circ F$  are both continuous.

**A9) Prove that the operators  $+$  and  $\cdot$  on real numbers are continuous.**

Proof: For any  $(x, y), (u, v) \in \mathbb{R}^2$ ,

$$|(x + y) - (u + v)| \leq |x - u| + |y - v| \leq 2|(x, y) - (u, v)|.$$

Hence  $+$  is uniformly continuous.

$$|x \cdot y - u \cdot v| \leq |x| \cdot |y - v| + |v| \cdot |x - u|.$$

Therefore  $\cdot$  is continuous.

**A10) Prove that the operators  $+$  and  $\cdot$  on  $\mathbf{M}_n(\mathbb{R})$  are continuous.**

Proof: The proof of A9) only uses the properties of norms, and the fact that  $\|A \cdot B\| \leq \|A\| \cdot \|B\|$ . This also holds for the norm  $\|A\| = \sup_{|\mathbf{x}|=1} |A\mathbf{x}|$  on  $\mathbf{M}_n(\mathbb{R})$ , therefore  $+$  and  $\cdot$  are continuous on  $\mathbf{M}_n(\mathbb{R})$ .

**A11) Prove that  $\mathbf{GL}_n(\mathbb{R})$  is an open set on  $\mathbf{M}_n(\mathbb{R})$ .**

Proof: The mapping  $\det : \mathbf{M}_n(\mathbb{R}) \rightarrow \mathbb{R}$  is continuous, since view as  $\det : \mathbb{R}^{n^2} \rightarrow \mathbb{R}$  it is a multi-linear mapping. The set  $\mathbf{GL}_n(\mathbb{R}) = \det^{-1}(\{x \in \mathbb{R} : x \neq 0\})$ , where  $\{x \in \mathbb{R} : x \neq 0\}$  is an open set on  $\mathbb{R}$ , therefore  $\mathbf{GL}_n(\mathbb{R})$  is an open set on  $\mathbf{M}_n(\mathbb{R})$ .

**A12) Prove that  $\text{Inv} : \mathbf{GL}_n(\mathbb{R}) \rightarrow \mathbf{GL}_n(\mathbb{R}), A \mapsto A^{-1}$  is continuous.**

Proof: Note that for any  $A, B \in \mathbf{GL}_n(\mathbb{R})$ ,

$$\|A^{-1} - B^{-1}\| \leq \frac{\|A - B\|}{\|A\| \cdot \|B\|}.$$

Hence  $\text{Inv}$  is continuous.

## 4.2 PSB

Prove the following equalities:

**B1)  $\lambda > 0, \lim_{x \rightarrow \infty} \frac{x^n}{e^{\lambda x}} = 0$ .**

Proof: By definition, for  $x > 0, e^{\lambda x} \geq (\lambda x)^{n+1}/(n+1)!$ . Hence for any  $\varepsilon > 0$ , let  $M = \frac{(n+1)!}{\lambda^{n+1}\varepsilon}$ , then for any  $x > M$ ,

$$\left| \frac{x^n}{e^{\lambda x}} \right| \leq \frac{(n+1)!}{\lambda^{n+1}x} < \varepsilon.$$

Therefore

$$\lim_{x \rightarrow \infty} \frac{x^n}{e^{\lambda x}} = 0.$$

**B2)  $\alpha > 0$ , then**

$$\lim_{x \rightarrow \infty} x^\alpha \log \left( 1 + \frac{1}{x} \right) = \begin{cases} \infty, & \alpha > 1; \\ 1, & \alpha = 1; \\ 0, & 0 < \alpha < 1. \end{cases}$$

Proof: If  $0 < \alpha < 1$ , then for any  $\varepsilon > 0$ , there exists  $\delta = \varepsilon^{1/(\alpha-1)}$  such that for any  $x > \delta$ ,

$$\left| x^\alpha \log \left( 1 + \frac{1}{x} \right) \right| \leq x^{\alpha-1} < \varepsilon.$$

If  $\alpha > 1$ , then for any  $\varepsilon > 0$ , there exists  $\delta = (2\varepsilon)^{1/\alpha-1}$  such that for any  $x > \delta$ ,

$$\left| x^\alpha \log \left( 1 + \frac{1}{x} \right) \right| \geq \frac{x^\alpha}{x+1} \geq \frac{1}{2} x^{\alpha-1} > \varepsilon.$$

If  $\alpha = 1$ , then for any  $\varepsilon > 0$ , there exists  $\delta = 1/\varepsilon$  such that for any  $x > \delta$ ,

$$1 - \varepsilon \leq \frac{x}{x+1} \leq x \log \left( 1 + \frac{1}{x} \right) \leq 1.$$

Therefore

$$\lim_{x \rightarrow \infty} x^\alpha \log \left( 1 + \frac{1}{x} \right) = \begin{cases} \infty, & \alpha > 1; \\ 1, & \alpha = 1; \\ 0, & 0 < \alpha < 1. \end{cases}$$

**B3)**  $\lim_{x \rightarrow 0^+} x^{-n} e^{-1/x^2} = 0.$

Proof: If  $x < 1$ , then  $e^{-1/x^2} \leq e^{-1/x} \leq (n+1)!x^{n+1}$ , hence for any  $\varepsilon > 0$ , let  $\delta = \varepsilon/(n+1)!$ , then for any  $x \in (0, \delta)$ ,  $x^{-n} e^{-1/x^2} \leq (n+1)!x \leq \varepsilon$ . Therefore

$$\lim_{x \rightarrow 0^+} x^{-n} e^{-1/x^2} = 0.$$

**B4) We know that  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ . Calculate**

$$\lim_{x \rightarrow 0} \frac{\cos x - 1}{x}, \text{ and } \lim_{x \rightarrow 0} \frac{\cos x - 1}{x^2/2}.$$

Solution: For any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that for any  $|x| < \delta$ ,  $\sin x \in ((1-\varepsilon)x, (1+\varepsilon)x)$ . Hence

$$\left| \frac{\cos x - 1}{x} \right| \leq \left| \frac{\sqrt{1 - \sin^2 x} - 1}{x} \right| \leq \left| \frac{\sin^2 x}{x(\sqrt{1 - \sin^2 x} + 1)} \right| \leq (1+\varepsilon)^2 x \leq \delta(1+\varepsilon)^2.$$

Therefore

$$\lim_{x \rightarrow 0} \frac{\cos x - 1}{x} = 0.$$

Likewise

$$\left| \frac{\cos x - 1}{x^2/2} + 1 \right| \leq \left| \frac{\sin^2 x - x^2(1 + \sqrt{1 - \sin^2 x})/2}{x^2/2 \cdot (\sqrt{1 - \sin^2 x} + 1)} \right| \leq (2\varepsilon + \sqrt{1 - \sin^2 x} - 1).$$

Therefore

$$\lim_{x \rightarrow 0} \frac{\cos x - 1}{x^2/2} = -1.$$

### 4.3 PSC: Root of Function:

**C1) Prove that  $x^3 + 2x - 1 = 0$  has exactly one root which lies in  $(0, 1)$ .**

Proof: Let  $f(x) = x^3 + 2x - 1$ , then  $f(0) = -1$  and  $f(1) = 2$ , so  $f(0) < 0 < f(1)$ . Since  $f$  is continuous and monotonically increasing on  $(0, 1)$ , there is exactly one root in  $(0, 1)$ .



**C2) Suppose  $0 \leq \lambda < 1$ ,  $b > 0$ , determine whether the equation  $x - \lambda \sin x = b$  has a solution.**

Solution:

**C3) Prove that  $\sin x = 1/x$  has infinitely many roots.**

Proof: For any  $n \in \mathbb{N}$ , let  $x_n = (2n + 1/2)\pi$ ,  $y_n = (2n + 3/2)\pi$ , and  $f(x) = \sin x - 1/x$ , then  $f(x_n) = 1 - 1/x_n > 0$ ,  $f(y_n) = -1 - 1/y_n < 0$ , therefore  $f$  has a root in  $(x_n, y_n)$ , and hence  $f$  has infinitely many roots.

**C4) Assume  $f \in C([0, 2])$  and  $f(0) = f(2)$ . Prove that  $f(x) - f(x + 1) = 0$  has a root in  $[0, 1]$ .**

Proof: Let  $g(x) = f(x) - f(x + 1)$ , then  $g(0) = f(0) - f(1) = -g(1)$  and  $g \in C([0, 1])$ . Therefore  $g$  has a root in  $[0, 1]$ .

**C5) Prove that  $x^3 + 3 = e^x$  has a solution in  $\mathbb{R}$ .**

Proof: Let  $f(x) = e^x - x^3 - 3$ , then  $\lim_{x \rightarrow \infty} f(x) = \infty$  and  $\lim_{x \rightarrow -\infty} f(x) = -\infty$ , therefore  $f$  has a root in  $\mathbb{R}$ .

**C6) Suppose  $f : [0, 2] \rightarrow \mathbb{R}$  is continuous and  $f(0) = f(2)$  then there exists  $x \in [1, 2]$  such that  $f(x) = f(x - 1)$ .**

Exactly the same as C4)?

**C7)  $f : \mathbb{R} \rightarrow \mathbb{R}$ , Prove that if for any  $c \in \mathbb{R}$ ,  $|f^{-1}(c)| = 2$ , then  $f$  is not continuous.**

Proof: If  $f$  is continuous on  $\mathbb{R}$ , suppose  $f^{-1}(0) = \{a < b\}$ , then  $f|_{[a, b]}$  is bounded. Suppose  $f(\frac{a+b}{2}) > 0$ , then for any  $t \in (a, b)$ ,  $f(t) > 0$  (otherwise  $|f^{-1}(0)| > 2$ ). Consider an arbitrary  $M > y = \sup_{x \in [a, b]} f(x)$ , and take  $t \in f^{-1}(M)$ . Assume  $t < a$ , then  $f(t) = M > y/2 > f(a) = 0$ , hence there exists  $s \in (t, a)$  such that  $f(s) = y/2$ . However there are at least two elements of  $f^{-1}(y/2)$  in  $(a, b)$ , leading to contradiction.

**C8) Suppose the continuous function  $f : [a, b] \rightarrow \mathbb{R}$  is injective. If  $f(a) < f(b)$ , prove that  $f$  is monotonically increasing.**

Proof: Otherwise suppose  $f(u) > f(v)$  for some  $u < v$ . Note that for any  $c \in (a, b)$ ,  $f(a) < f(c) < f(b)$ , otherwise  $f(c) < f(a) \implies \exists d \in (c, b), f(d) = f(a)$ , or  $f(c) > f(b) \implies \exists d \in (a, c), f(d) = f(b)$ . Hence  $a < u < v < b$ . Likewise consider  $u < v < b$  we get  $f(u) > f(v) > f(b)$ , and by  $a < u < v$  we get  $f(a) > f(u) > f(v)$ , therefore  $f(a) > f(b)$ , a contradiction.

## 4.4 PSD: Calculation of Limits

$n, m$  are positive integers.

(1)

$$\lim_{x \rightarrow \infty} \frac{a_0 x^n + a_1 x^{n-1} + \cdots + a_n}{b_0 x^m + b_1 x^{m-1} + \cdots + b_m} = \begin{cases} 0, & m > n, \\ \infty, & m < n, a_0 > 0, \\ -\infty, & m < n, a_0 < 0, \\ \frac{a_0}{b_0}, & m = n. \end{cases}$$

(2)  $a > 1, b > 0$

$$\lim_{x \rightarrow \infty} \frac{x^b}{a^x} = 0.$$

(3)  $a > 0$

$$\lim_{x \rightarrow \infty} \frac{\log x}{x^a} = 0.$$

(4)  $a > 0$

$$\lim_{x \rightarrow 0^+} x^a \log x = 0.$$

(5)

$$\lim_{x \rightarrow \infty} \left( \frac{x^2 + 1}{x^2 - 2} \right)^{x^2} = \lim_{x \rightarrow \infty} \left( \frac{x + 1}{x - 2} \right)^x = e^3.$$

(6)

$$\lim_{x \rightarrow \infty} (x - \sqrt{x^2 - a}) = \lim_{x \rightarrow \infty} \frac{a}{x + \sqrt{x^2 - a}} = 0.$$

(7)

$$\lim_{x \rightarrow \infty} \sqrt{x+1} - \sqrt{x-1} = \lim_{x \rightarrow \infty} \frac{2}{\sqrt{x+1} + \sqrt{x-1}} = 0.$$

(8)

$$\lim_{x \rightarrow 0} \frac{(1+x)(1+2x)(1+3x) - 1}{x} = 1 + 2 + 3 = 6.$$

(9)

$$\lim_{x \rightarrow 1} \frac{x + x^2 + \cdots + x^n - n}{x - 1} = \frac{n(n+1)}{2}.$$

(10)

$$\lim_{x \rightarrow 1} \frac{x^{100} - 2x + 1}{x^{50} - 2x + 1} = \frac{49}{24}.$$

(11)

$$\lim_{x \rightarrow 1} \left( \frac{m}{1-x^m} - \frac{n}{1-x^n} \right) = \frac{m-n}{2}.$$

Proof: Note that

$$\begin{aligned} \lim_{x \rightarrow 1} \left( \frac{m}{1-x^m} - \frac{n}{1-x^n} \right) &= \lim_{x \rightarrow 1} \frac{m(1+x+\cdots+x^{n-1}) - n(1+x+\cdots+x^{m-1})}{(1+x+\cdots+x^{m-1})(1+x+\cdots+x^{n-1})(1-x)} \\ &= \frac{1}{mn} \cdot \lim_{x \rightarrow 1} \frac{m(x-1+\cdots+x^{n-1}-1) - n(x-1+\cdots+x^{m-1}-1)}{1-x} \\ &= \frac{1}{mn} \cdot (-m(1+2+\cdots+(n-1)) + n(1+2+\cdots+(m-1))) \\ &= \frac{m-n}{2}. \end{aligned}$$

(12)

$$\lim_{x \rightarrow 0} \frac{(1+x)^a - 1}{x} = a.$$

(diverges if  $a = 0$ ).

(13)

$$\lim_{x \rightarrow 1} \frac{x^a - 1}{x^b - 1} = \frac{a}{b}.$$

(14)

$$\lim_{x \rightarrow \infty} (\log x)^{1/x} = \lim_{x \rightarrow \infty} e^{(\log \log x)/x} = 1.$$

(15)  $a, b > 0$

$$\lim_{x \rightarrow 0} \left( \frac{a^x + b^x}{2} \right)^{1/x} = \sqrt{ab}.$$

(16)

$$\lim_{x \rightarrow \infty} \sqrt[k]{(x+a_1)(x+a_2)\cdots(x+a_k)} - x$$

Proof: Let  $y = (x+a_1)(x+a_2)\cdots(x+a_k)$  and  $s = a_1 + \cdots + a_k$ , then

$$\frac{sx^{k-1}}{ky^{(k-1)/k}} \leq \sqrt[k]{y} - x = \frac{y - x^k}{y^{(k-1)/k} + \cdots + x^{k-1}} \leq \frac{sx^{k-1} + \prod_{i=1}^k (1+a_i)x^{k-2}}{kx^{k-1}}.$$

Therefore

$$\lim_{x \rightarrow \infty} \sqrt[k]{y} - x = s = \sum_{i=1}^k a_i.$$

(17)

$$\lim_{x \rightarrow 0} \frac{(\sqrt{1+x^2} + x)^n - (\sqrt{1+x^2} - x)^n}{x} = 2n.$$

(18)

$$\lim_{x \rightarrow \frac{\pi}{2}} (\sin x)^{\tan x} = 1.$$

(19)

$$\lim_{x \rightarrow \infty} \left( \sin \frac{1}{x} + \cos \frac{1}{x} \right)^x = e.$$

(20)  $\alpha > 0$ ,

$$\lim_{x \rightarrow \infty} \frac{\sqrt{x + \sqrt{x + \sqrt{x}}}}{x^\alpha} = \begin{cases} 0, & \alpha > \frac{1}{2}, \\ 1, & \alpha = \frac{1}{2}, \\ \infty, & \alpha < \frac{1}{2}. \end{cases}$$

(21)  $\alpha > 0$ ,

$$\lim_{x \rightarrow 0^+} \frac{\sqrt{x + \sqrt{x + \sqrt{x}}}}{x^\alpha} = \begin{cases} 0, & \alpha < \frac{1}{8}, \\ 1, & \alpha = \frac{1}{8}, \\ \infty, & \alpha > \frac{1}{8}. \end{cases}$$

Proof: Note that for  $x \in (0, 1)$ ,

$$x^{1/8-\alpha} \leq \frac{\sqrt{x + \sqrt{x + \sqrt{x}}}}{x^\alpha} \leq 2x^{1/8-\alpha}.$$

And for any  $\varepsilon > 0$  there exists  $\delta = (1 + \varepsilon)\varepsilon$  such that for any  $x < \delta$ ,  $\sqrt{x + \sqrt{x + \sqrt{x}}} < \varepsilon x^{1/8}$ .  
Therefore

$$\lim_{x \rightarrow 0^+} \frac{\sqrt{x + \sqrt{x + \sqrt{x}}}}{x^\alpha} = \begin{cases} 0, & \alpha < \frac{1}{8}, \\ 1, & \alpha = \frac{1}{8}, \\ \infty, & \alpha > \frac{1}{8}. \end{cases}$$

## 4.5 Problem E

Prove that for any  $A \subset \mathbb{R}$  that is countable, there exists a monotonic function  $f : \mathbb{R} \rightarrow \mathbb{R}$ , such that the set of discontinuities of  $f$  is exactly  $A$ .

Proof: Let  $A = \{x_1, x_2, \dots\}$  and  $f(x) = \sup\{1 - 2^n : x_n < x\}$ , (define  $\sup \emptyset = 0$ ) then  $f$  is monotonically increasing and the set of discontinuities is exactly  $A$ .

## 4.6 Problem F

$f : [0, 1] \rightarrow [0, 1]$  is monotonic. Prove that  $f$  has a fixed point.

Proof: Otherwise suppose that  $f$  has no fixed points. Let  $S = \{t \in [0, 1] : f(t) > t\}$  and  $x = \sup S$ . Note that  $0 \in S$  so  $S$  is non-empty. If  $x \in S$ , then  $f(x) > x$  so  $f(f(x)) > f(x)$  ( $f$  is monotonic) then  $x < f(x) \in S$  which leads to contradiction. If  $x \notin S$ , then  $f(x) < x$ . Take  $y \in (f(x), x) \cap S$ , ( $y$  exists since  $x = \sup S$ ) then  $f(x) > f(y) > y > f(x)$ , a contradiction.

## 4.7 Problem G

Consider all self-homeomorphisms of  $[0, 1]$ , i.e.

$$\text{Homeo}([0, 1]) = \{f : [0, 1] \rightarrow [0, 1] : f \text{ is a continuous bijective}\}$$

We know that for any  $f \in \text{Homeo}([0, 1])$ ,  $f^{-1} \in \text{Homeo}([0, 1])$ . Suppose  $f, g \in \text{Homeo}([0, 1])$  and the only fixed points of  $f, g$  are 0, 1. Prove that there exists  $h \in \text{Homeo}([0, 1])$ , such that

$$h \circ f \circ h^{-1} = g.$$

Proof: Take  $x_0 = 1/2$ , and let  $I_n = [f^n(x_0), f^{n+1}(x_0)]$ ,  $J_n = [g^n(x_0), g^{n+1}(x_0)]$ . Note that  $(0, 1) = \bigcup_{n \in \mathbb{Z}} I_n = \bigcup_{n \in \mathbb{Z}} J_n$ . Define  $h_0 : I_0 \rightarrow J_0$ ,  $x \mapsto kx + b$  such that the line  $h_0$  passes  $(x_0, x_0)$  and  $(f(x_0), g(x_0))$ , i.e.  $x \mapsto \frac{g(x_0) - x_0}{f(x_0) - x_0}(x - x_0) + x_0$ . Define  $h_n : I_n \rightarrow J_n$ ,  $x \mapsto g^n \circ f^{-n}(x)$ , and  $h : [0, 1] \rightarrow [0, 1]$  such that

$$h(x) = \begin{cases} x, & x \in \{0, 1\}, \\ h_n(x), & x \in I_n. \end{cases}$$

Then for any  $x \in I_n$ ,  $f(x) \in I_{n+1}$  hence  $h(f(x)) = g^{n+1} \circ f^{-n}(x) = g(h(x))$ . Since  $h$  maps  $I_n$  to  $J_n$  bijectively,  $h$  is a bijection on  $[0, 1]$ . For any  $x \in I_n \cap I_{n+1}$  the value of  $h$  does not depend on which interval we choose, and  $h$  is continuous on any interval  $I_n$ , therefore  $h$  is a continuous bijection.

## 5 Homework 5: Infinity of Prime

### 5.1 PSA

**A1) Prove that  $e^x$  is uniformly continuous on  $(-\infty, 0]$  but not on  $\mathbb{R}$ .**

Proof: For  $y < x \leq 0$  and  $|x - y| < \varepsilon$ ,

$$e^x - e^y = e^y(e^{y-x} - 1) \leq e^\varepsilon - 1.$$

Hence  $e^x$  is uniformly continuous on  $(-\infty, 0]$ . But for any  $\delta > 0$ , there exists  $y$  and  $x = y + \delta$  such that

$$e^x - e^y = e^y \cdot (e^\delta - 1) > 1.$$

Therefore  $e^x$  is not uniformly continuous on  $\mathbb{R}$ .

**A2) Prove that the function  $f : \mathbb{R}_{>0} \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $(x, \alpha) \mapsto x^\alpha$  is continuous on  $\mathbb{R}_{>0} \times \mathbb{R}$ .**

Proof: For  $(x, \alpha), (y, \beta)$ ,

$$|x^\alpha - y^\beta| \leq |x^\alpha - y^\alpha| + |y^\alpha - y^\beta|.$$

Since  $x^\alpha$  and  $a^x$  are both continuous (as functions of  $x$ ), so is  $x^\alpha : \mathbb{R}_{>0} \times \mathbb{R} \rightarrow \mathbb{R}$ .

**A3) Prove that for any  $x, y > 0$  and  $\alpha, \beta$ ,  $(xy)^\alpha = x^\alpha y^\alpha$ ,  $(x^\alpha)^\beta = x^{\alpha\beta}$ ,  $a^{\log_a x} = x$ . If  $x > 0, y > 0$ , then  $a^{x+y} = a^x a^y$ ,  $\log_a(x \cdot y) = \log_a x + \log_a y$ .**

Proof: See PSE of HW2.

**A4) Consider the sequence of functions  $\{f_n(x)\}_{n \geq 1}$  defined on  $[0, 1]$ , where  $f_n(x) = x^n$ . Prove that for any  $a < 1$ ,  $\{f_n(x)\}_{n \geq 1}$  converges uniformly to 0 on  $[0, a]$ , but  $\{f_n(x)\}_{n \geq 1}$  does not converge uniformly on  $[0, 1)$ .**

Proof: For any  $a < 1$ , and any  $\varepsilon > 0$ , let  $N = \log_a x$ , then for any  $n > N$ ,  $f_n(x) < \varepsilon$ , hence  $\{f_n(x)\}_{n \geq 1}$  converges uniformly to 0 on  $[0, a]$ . Let  $\varepsilon = 1/2$ , then for any  $N \in \mathbb{N}$ , there exists  $1 > x > 2^{-1/N}$  such that  $f_N(x) > \varepsilon$ . Hence  $\{f_n(x)\}_{n \geq 1}$  is not uniformly convergent on  $[0, 1)$ .

**A5) Consider the sequence of functions  $\{f_n(x)\}_{n \geq 1}$ , where  $f_n(x) = \frac{nx}{1+n^2x^2}$ . Prove that  $\{f_n(x)\}_{n \geq 1}$  converges point-wise to 0 on  $\mathbb{R}$ , but does not converge uniformly.**

Proof: For any  $x \in \mathbb{R}$ , and any  $\varepsilon > 0$ , there exists  $N = 1/(x\varepsilon)$  such that for any  $n \geq N$ ,

$$\left| \frac{nx}{1+n^2x^2} \right| \leq \frac{1}{|nx|} < \varepsilon.$$

Hence  $f_n(x)$  converges to 0 for any  $x \in \mathbb{R}$ .

Let  $\varepsilon = 1/2$ , then for any  $n \in \mathbb{N}$ , there exists  $x = 1/n$  such that  $f_n(x) = \varepsilon$ , so  $f$  is not uniformly continuous on  $\mathbb{R}$ .

**A6) Consider the sequence of functions  $\{f_n(x)\}_{n \geq 1}$ , where**

$$f_n(x) = \begin{cases} \frac{nx^2}{1+n^2x^2}, & x > 0; \\ \frac{nx}{1+n^2x^2}, & x \leq 0. \end{cases}$$

Determine the convergence of  $\{f_n(x)\}_{n \geq 1}$  on  $\mathbb{R}$  (both point-wise and uniformly).

Proof: For any  $\varepsilon > 0$ , let  $N = \max\{1/\varepsilon, 1/4\varepsilon^2\}$ , then for any  $x > 0$  and  $n > N$ ,

$$|f_n(x) - x| = \left| \frac{x}{1+nx} \right| < \frac{1}{n} < \varepsilon.$$

For any  $x < 0$ ,

$$|f_n(x) - x| = \left| \frac{x}{1+n^2x^2} \right| \leq \frac{1}{2\sqrt{n}} < \varepsilon.$$

Hence  $\{f_n\}_{n \geq 1}$  converges uniformly to  $x$ .

**A7) Given  $\varphi : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$  such that  $\varphi(0) = 0$ ,  $\lim_{x \rightarrow \infty} \varphi(x) = 0$ ,  $\varphi$  is continuous and not identically zero. Prove that the sequences  $\{f_n(x)\}_{n \geq 1}$  and  $\{g_n(x)\}_{n \geq 1}$  converge point-wise to 0, but uniformly, where  $f_n(x) = \varphi(nx)$ ,  $g_n(x) = \varphi(x/n)$ .**

Proof: Point-wise convergence is trivial. Let  $\varepsilon = |\varphi(1)| > 0$ , then for any  $n$  there exists  $x = 1/n > 0$  such that  $|f_n(x)| = \varepsilon$ , hence  $\{f_n(x)\}_{n \geq 1}$  is not uniformly convergent. Likewise  $\{g_n(x)\}_{n \geq 1}$  is not uniformly continuous.

**A8)  $f \in C([a, b])$ . For  $n \geq 1$ , let  $a_k = a + (k-1)(b-a)/n$ . Define**

$$S_n = \sum_{k=1}^n \frac{b-a}{n} f(a_k).$$

Prove that  $\{S_n\}_{n \geq 1}$  converges, and denote this limit by  $\int_a^b f$ . Further show that the mapping

$$\int_a^b : C([a, b]) \rightarrow \mathbb{R}, f \mapsto \int_a^b f$$

is linear and continuous with metric  $d_\infty$  on  $C([a, b])$ .

Proof: For any  $n, m \in \mathbb{N}$ , note that  $|S_n - S_m| \leq |S_n - S_{nm}| + |S_{nm} - S_m|$ , and

$$|S_n - S_{nm}| \leq \sum_{k=1}^n \frac{b-a}{n} \left| f(a_k^{(n)}) - \frac{1}{m} \sum_{j=1}^m f(a_{n(k-1)+j}^{(nm)}) \right| \leq (b-a) \sup_{|x-y| < 1/n} |f(x) - f(y)|.$$

Since  $f$  is uniformly continuous on  $[a, b]$ , the sequence  $\{S_n\}_{n \geq 1}$  is Cauchy.

Obviously  $\int_a^b \cdot$  is linear, and for  $f, g \in C([a, b])$ ,

$$\left| \int_a^b f - \int_a^b g \right| = \lim_{n \rightarrow \infty} |S_n(f) - S_n(g)| \leq (b-a) \|f - g\|_\infty.$$

Hence  $\int_a^b \cdot$  is continuous on  $C([a, b])$  with metric  $d_\infty$ .

**A9) For any  $f : [a, \infty) \rightarrow \mathbb{R}$ , suppose  $f$  is bounded on any closed interval  $[a, b]$ , then when the limits in RHS exist,**

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{f(x)}{x} &= \lim_{x \rightarrow \infty} f(x+1) - f(x). \\ \lim_{x \rightarrow \infty} f(x)^{1/x} &= \lim_{x \rightarrow \infty} \frac{f(x+1)}{f(x)}, \text{ if for any } x \in [a, \infty), f(x) \geq c > 0. \end{aligned}$$

Proof: Suppose  $\lim_{x \rightarrow \infty} f(x+1) - f(x) = A$ , then for any  $\varepsilon > 0$  there exists  $M$  such that for any  $x > M$ ,  $|f(x+1) - f(x) - A| < \varepsilon$ , so for any  $n \geq 1$ ,  $|f(x+n) - f(x) - nA| < n\varepsilon$ . Hence

$$\left| \frac{f(n+x)}{n+x} - A \right| \leq \left| \frac{f(n+x) - f(x) - nA}{n+x} \right| + \left| \frac{f(x) - xA}{n+x} \right| \leq \varepsilon A + \frac{|f(x) - xA|}{n} \rightarrow 0.$$

For any  $x > M$ . Therefore (since  $f$  is bounded on any closed interval) there exists  $N$  such that for any  $x > N$ ,  $|f(x)/x - A| < 2\varepsilon A$ , and hence

$$\lim_{x \rightarrow \infty} \frac{f(x)}{x} = A = \lim_{x \rightarrow \infty} f(x+1) - f(x).$$

Substitute  $f$  by  $\log f$  and we obtain the second identity.

## 5.2 PSB: Uniform Continuity

Determine whether the following functions  $f$  are uniformly continuous on  $I$ :

**B1)**  $f(x) = x^{1/3}$ ,  $I = (0, \infty)$

For any  $\varepsilon > 0$  and  $x - y \in (0, \varepsilon)$ ,

$$x^{1/3} - y^{1/3} = \frac{x - y}{x^{2/3} + x^{1/3}y^{1/3} + y^{2/3}} \leq \frac{\varepsilon}{\varepsilon^{2/3}} = \varepsilon^{1/3}.$$

Hence  $f(x)$  is uniformly continuous on  $I$ .

**B2)**  $f(x) = \log x$ ,  $I = (0, 1)$

For any  $\varepsilon > 0$  and  $x - y \in (0, \varepsilon)$ ,

$$\log x - \log y = \log \left( 1 + \frac{x - y}{y} \right).$$

When  $y \rightarrow 0$  and  $x - y$  is constant,  $\log x - \log y \rightarrow \infty$ , hence  $\log x$  is not uniformly continuous on  $I$ .

**B3)**  $f(x) = \cos x^{-1}$ ,  $I = (0, 1)$

Note that for  $x_n = 1/(2n\pi)$  and  $y_n = 1/(2n\pi + \pi)$ ,  $f(x_n) = 1$  and  $f(y_n) = -1$ . Hence for  $\varepsilon = 1$  and any  $\delta > 0$ , there exists  $n$  such that  $|x_n - y_n| < \delta$  but  $|f(x_n) - f(y_n)| = 2 > \varepsilon$ . Therefore  $f$  is not uniformly continuous on  $I$ .

**B4)**  $f(x) = x \cos x^{-1}$ ,  $I = (0, \infty)$

For  $x > y > 1$  and  $|x - y| < \varepsilon$ ,

$$\begin{aligned} |x \cos x^{-1} - y \cos y^{-1}| &\leq |x - y| |\cos x^{-1}| + |y| \cdot |\cos x^{-1} - \cos y^{-1}| \\ &\leq \varepsilon + 2|y| \cdot |\sin(x^{-1} + y^{-1})/2 \sin(x^{-1} - y^{-1})/2| \leq \varepsilon + \frac{y}{2} \left( \frac{1}{y^2} - \frac{1}{x^2} \right) \leq 2\varepsilon. \end{aligned}$$

For  $1 > x > y$  and  $|x - y| < \varepsilon$ ,

$$|x \cos x^{-1} - y \cos y^{-1}| \leq |x| + |y| < 2\varepsilon.$$

Hence  $f$  is uniformly continuous on  $I$ .

### 5.3 PSC: Existence of Limits

**C1)**  $\alpha > 0$ ,

$$\lim_{x \rightarrow 1} \frac{\log x}{(x - 1)^\alpha} = \lim_{t \rightarrow 0} \frac{\log(1 + t)}{t^\alpha} = \lim_{t \rightarrow 0} t^{1-\alpha}$$

exists iff  $\alpha \leq 1$ .

**C2)**  $\alpha > 0$ ,

$$\lim_{x \rightarrow 1} \frac{e^x - e}{(x - 1)^\alpha} = e \lim_{t \rightarrow 0} \frac{e^t - 1}{t^\alpha} = \lim_{t \rightarrow 0} e t^{1-\alpha}.$$

exists iff  $\alpha \leq 1$ .

**C3)**  $\alpha > 0$ ,

$$\lim_{x \rightarrow 1} \frac{x^x - 1}{(x - 1)^\alpha} = \lim_{x \rightarrow 1} \frac{x^x (\log x + 1)}{\alpha (x - 1)^{\alpha-1}}$$

exists iff  $\alpha \leq 1$ .

**C4)**  $\alpha > 0$ ,

$$\lim_{x \rightarrow 1} \frac{\sqrt[3]{1 - \sqrt{x}}}{(x - 1)^\alpha}$$

exists iff  $\alpha \leq 1/3$ .



C5)

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x^2} - 1}{1 - \cos x} = 1.$$

C6)

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x^4} - 1}{1 - \cos^2 x} = 0.$$

C7)  $\alpha > 0$ ,

$$\lim_{x \rightarrow 1} \frac{(x-1)^\alpha}{\sin(\pi x)}$$

exists iff  $\alpha \geq 1$ .

## 5.4 PSD: Problems on Uniform Continuity

**D1) If  $f$  is continuous, monotonic and bounded on the open interval  $I$ , then  $f$  is uniformly continuous on  $I$ .**

Proof: Otherwise if there exists  $\varepsilon > 0$  such that for any  $\delta > 0$  there exists  $|x - y| < \delta$  such that  $|f(x) - f(y)| > \varepsilon$ . We define  $x_n, y_n$  inductively as follows: Let  $L = \min\{x_1, \dots, x_{n-1}\}$ ,  $R = \max\{y_1, \dots, y_{n-1}\}$ . Since  $f$  is uniformly continuous on  $[L, R]$ , there exists  $\delta > 0$  such that for any  $|s - t| < \delta$ ,  $|f(s) - f(t)| < \varepsilon$ . Hence there exists  $x < y$  such that  $x, y \notin [L, R]$ ,  $|x - y| < \delta$  and  $|f(x) - f(y)| > \varepsilon$ . Let  $x_n = x, y_n = y$ , then  $(x_n, y_n)$  are disjoint intervals and  $|f(x_n) - f(y_n)| > \varepsilon$ . Which contradicts the fact that  $f$  is monotonic and bounded. Therefore  $f$  is uniformly continuous on  $I$ .

**D2)  $I$  is an interval with finite length. Prove that the function  $f$  on  $I$  is uniformly continuous iff for any Cauchy sequence  $\{x_n\}_{n \geq 1} \subset I$ ,  $\{f(x_n)\}_{n \geq 1}$  is also a Cauchy sequence.**

( $f$  should be continuous, otherwise after changing the value of  $f$  at one point,  $\{f(x_n)\}$  remains a Cauchy sequence.)

Proof:  $\implies$  If  $\{x_n\}_{n \geq 1}$  is a Cauchy sequence, then for any  $\varepsilon > 0$  there exists  $\delta > 0$  such that for all  $|x - y| < \delta$ ,  $|f(x) - f(y)| < \varepsilon$ . There exists  $N$  such that for all  $n, m > N$ ,  $|x_n - x_m| < \delta$ , hence  $|f(x_n) - f(x_m)| < \varepsilon$ , so  $\{f(x_n)\}_{n \geq 1}$  is a Cauchy sequence.

$\impliedby$  If  $I = (a, b)$  is open we can take  $x_n \rightarrow a$  and define  $f(a) = \lim_{n \rightarrow \infty} f(x_n)$ , hence we can assume that  $I$  is closed. Therefore  $f$  is uniformly continuous.

**D3)  $f$  is uniformly continuous on  $\mathbb{R}$ . Prove that there exists  $a, b \in \mathbb{R}_{>0}$  such that for any  $x \in \mathbb{R}$ ,**

$$|f(x)| \leq a|x| + b.$$

Proof: For  $\varepsilon = 1$ , there exists  $\delta > 0$  such that  $|x - y| < \delta \implies |f(x) - f(y)| < 1$ . Hence let  $C = \sup_{x \in [0, \delta]} |f(x)|$ , then  $|f(x)| \leq C + |x| \cdot (\frac{1}{\delta} + 1)$ .

**D4) Suppose  $f$  is uniformly continuous on  $[0, \infty)$  and for any  $x \in [0, 1]$ ,  $\lim_{n \rightarrow \infty} f(x+n) = 0$ . Prove that**

$$\lim_{x \rightarrow \infty} f(x) = 0.$$

If we change the condition to  $f$  is continuous, will the statement still hold?

Proof: For any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $|x - y| < \delta \implies |f(x) - f(y)| < \varepsilon$ . Let  $N = \lfloor 1/\delta \rfloor + 1$ , then for any  $1 \leq n \leq N$ , there exists  $M_n$  such that for all  $m > M_n$ ,  $|f(m + n/N)| < \varepsilon$ . Let  $M = \max\{M_1, \dots, M_N\}$ , then for all  $x > M$ , there exists  $m \in \mathbb{Z}_{>M}$  and  $1 \leq n \leq N$  such that  $|x - m - n/N| < \delta$ . Hence

$$|f(x)| \leq \varepsilon + |f(m + n/N)| < 2\varepsilon.$$

Therefore  $\lim_{x \rightarrow \infty} f(x) = 0$ .

**D5) Suppose  $X$  is an interval,  $f : X \rightarrow \mathbb{R}$  is continuous. If there is a constant  $L > 0$  such that for any  $x, y \in X$ ,**

$$|f(x) - f(y)| \leq L|x - y|.$$

We say  $f$  satisfy the Lipschitz condition on  $X$ .

1. Prove that  $f$  satisfy the Lipschitz condition implies  $f$  is uniformly continuous.

Proof: For any  $\varepsilon > 0$ , let  $\delta = \varepsilon/L$ , then for any  $|x - y| < \delta$ ,  $|f(x) - f(y)| \leq L|x - y| < \varepsilon$ .

2. Determine whether the reversed statement holds.

Consider the function  $f(x) = x^{1/2}$ , then  $f$  is uniformly continuous but  $\frac{f(x)-f(y)}{x-y} = \frac{1}{\sqrt{x}+\sqrt{y}}$  is unbounded, hence does not satisfy the Lipschitz condition.

3. If  $f$  satisfy the Lipschitz condition on  $[a, \infty)$ , where  $a > 0$ , prove that  $f(x)/x$  is uniformly continuous on  $[a, \infty)$ .

Proof: Same as D3), there exists  $C$  such that  $|f(x)| \leq C|x|$  for  $x \in [a, \infty)$ , then for  $a < x < y$ ,

$$\begin{aligned} \left| \frac{f(x)}{x} - \frac{f(y)}{y} \right| &= \frac{|xf(y) - yf(x)|}{xy} \leq \frac{x|f(y) - f(x)| + |f(x)|(y - x)}{xy} \\ &\leq \frac{L + C}{y} \cdot |x - y|. \end{aligned}$$

Hence  $f(x)/x$  satisfy the Lipschitz condition.

## 5.5 PSE:

Exactly the same as PSC in HW4?

## 5.6 PSF: Calculate Limits

**F1)**

$$\lim_{x \rightarrow \pi} \frac{\sin mx}{\sin nx} = \frac{m(-1)^m}{n(-1)^n}.$$

**F2)**

$$\lim_{x \rightarrow 0} \frac{1 - \cos x \sqrt{\cos 2x} \sqrt[3]{\cos 3x}}{x^2} = 3.$$

**F3)**

$$\lim_{x \rightarrow \infty} \sin \sqrt{1+x} - \sin \sqrt{x} = 0.$$

Since the function  $\sin x$  is uniformly continuous and  $\lim_{x \rightarrow \infty} \sqrt{1+x} - \sqrt{x} = 0$ .

**F4)**

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x \sin x} - 1}{e^{x^2} - 1} = \frac{1}{2}.$$

Since  $\lim_{x \rightarrow 0} x^2/(e^{x^2} - 1) = 1$ ,  $\lim_{x \rightarrow 0} x \sin x/x^2 = 1$  and  $\lim_{x \rightarrow 0} 1/(1 + \sqrt{1+x \sin x}) = 1/2$ .

**F5)**

$$\lim_{n \rightarrow \infty} \sin^{(n)}(x) = 0.$$

Since the sequence  $\{a_n = \sin^{(n)}(x)\}_{n \geq 1}$  is decreasing and bounded by 0, and its limit  $A$  satisfy  $A = \sin A$ . Therefore  $\lim_{n \rightarrow \infty} \sin^{(n)}(x) = 0$ .

## 5.7 Problem G

The continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$  satisfy the following property: for any  $\delta > 0$ ,

$$\lim_{n \rightarrow \infty} f(n\delta) = 0.$$

Prove that  $\lim_{x \rightarrow \infty} f(x) = 0$ .

Proof: Consider any  $\varepsilon > 0$ . For any  $N \in \mathbb{N}$ ,

$$A_N = \{\delta > 0 : \forall n \geq N, |f(n\delta)| < \varepsilon\}.$$

Then by the continuity of  $f$ ,  $A_N$  is closed, and by  $\lim_{n \rightarrow \infty} f(n\delta) = 0$  for any  $\delta > 0$ ,  $\bigcup_{N \geq 1} A_N = \mathbb{R}_{>0}$ . Hence by Baire Category Theorem, there exists an  $N > 0$  such that  $(a, b) \subset A_N$  for some interval  $(a, b)$ . Let  $X = \{x \in \mathbb{R}_{>0} : |f(x)| < \varepsilon\}$ , then since  $(a, b) \subset A_N$ , for any  $n \geq N$ ,  $(na, nb) \subset X$ . Note that when  $n > b/(b-a)$ ,  $nb > (n+1)a$ , hence there exists  $M > 0$  such that  $(M, \infty) \subset X$ . Therefore  $\lim_{x \rightarrow \infty} f(x) = 0$ .

## 5.8 Problem H

The continuous function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  satisfy the following properties:

1.  $\lim_{x \rightarrow \infty} (\varphi(x) - x) = \infty$ .
2.  $\{x \in \mathbb{R} : \varphi(x) = x\}$  is a non-empty finite set.

Prove that if  $f : \mathbb{R} \rightarrow \mathbb{R}$  is continuous and  $f \circ \varphi = f$ , then  $f$  is constant.

(Probably need the condition  $\lim_{x \rightarrow -\infty} \varphi(x) - x = -\infty$ ).

Proof: Suppose  $\{x \in \mathbb{R} : \varphi(x) = x\} = \{a_1, \dots, a_n\}$  where  $a_1 < \dots < a_n$ . For any  $x \in \mathbb{R}$ , we will show that  $f(x) \in \{f(a_1), \dots, f(a_n)\}$  hence  $f$  is constant.

If  $a_i < x < a_{i+1}$ . Suppose  $\varphi(x) > x$ , then let  $x_0 = x$ , and inductively define  $x_k$  as a point in  $(a_k, x_{k-1})$  such that  $\varphi(a_i) = a_i < \varphi(x_k) = x_{k-1} < \varphi(x_{k-1})$ . Since  $\varphi$  is continuous and  $a_1, \dots, a_n$  are all the roots of  $\varphi(x) = x$ , we know that  $\varphi(x_k) > x_k$  for all  $k \geq 0$ . The sequence  $\{x_k\}_{k \geq 0}$  is decreasing and bounded by  $a_i$ , hence has a limit  $A$ . From  $\varphi(x_k) = x_{k-1}$  we know that  $\varphi(A) = A$ , so  $A = a_i$ . Note that  $f(x_k) = f(\varphi(x_k)) = f(x_{k-1})$ , hence  $f(x) = f(x_k) = \lim_{k \rightarrow \infty} f(x_k) = f(a_i)$ . The case  $\varphi(x) < x$  is the same, by constructing a sequence which tends to  $a_{i+1}$ .

If  $x > a_n$ , then  $\varphi(x) > x$ , likewise we can construct a sequence  $x_k$  such that  $x_{k-1} = \varphi(x_k)$  and  $\lim_{k \rightarrow \infty} x_k = a_n$ . The case  $x < a_1$  is the same.  
Hence for all  $x \in \mathbb{R}$ ,  $f(x) \in \{f(a_1), \dots, f(a_n)\}$ .

### 5.9 Problem I

The continuous function  $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$  satisfy  $\lim_{x \rightarrow \infty} f(x)/x = 0$ . Suppose  $\{a_n\}_{n \geq 1}$  is a sequence of non-negative real numbers and the sequence  $\{a_n/n\}_{n \geq 1}$  is bounded. Prove that  $\lim_{n \rightarrow \infty} f(a_n)/n = 0$ .  
Proof: Suppose  $\{a_n/n\}$  is bounded by  $M$ .

For any  $\varepsilon > 0$ , we need to find  $N$  such that  $n \geq N \implies |f(a_n)| < \varepsilon n$ . For  $C > 0$ , we can divide  $n$  into two parts: If  $a_n \leq C$ , then  $|f(a_n)| \leq \sup_{x \in [0, C]} |f(x)|$ , otherwise  $a_n \geq C$ , then  $|f(a_n)| \leq \sup_{x \geq C} |f(x)/x| \cdot Mn$ . Therefore, if we choose  $C > 0$  such that  $\sup_{x \geq C} |f(x)/x| < \varepsilon/M$ , and choose  $N$  such that  $N > \sup_{x \in [0, C]} |f(x)|/\varepsilon$ , then for any  $n \geq N$ ,  $|f(a_n)| < \varepsilon n$ , hence

$$\lim_{n \rightarrow \infty} \frac{f(a_n)}{n} = 0.$$

### 5.10 Ex: Proof of the infinity of primes using topology

Proof: Assume otherwise that the set  $\mathcal{P}$  of primes is finite. Let  $L_{a,b} = \{at + b : t \in \mathbb{Z}\}, \forall (a, b) \in I = \mathbb{Z}_{>0} \times \mathbb{Z}$ . Then

$$\mathbb{Z} \subset \bigcup_{b \in \mathbb{Z}} L_{1,b} \subset \bigcup_{(a,b) \in I} L_{a,b} \subset \mathbb{Z} \implies \bigcup_{(a,b) \in I} L_{a,b} = \mathbb{Z}.$$

and for any  $x \in \bigcap_{i=1}^n L_{a_i, b_i}$ , let  $a = \text{lcm}(a_1, \dots, a_n)$ , then

$$x \in L_{a,x} \subset \bigcap_{i=1}^n L_{a_i, b_i}.$$

Hence  $L_{a,b}$  form a base. Consider the topology  $\mathcal{T}$  on  $\mathbb{Z}$  generated by the base  $\{L_{a,b} : (a,b) \in I\}$ . Note that

$$L_{a,b} = \mathbb{Z} \setminus \bigcup_{r=1}^{a-1} L_{a, b+r}$$

so  $L_{a,b}$  is also closed. Since  $\mathcal{P}$  is finite, the set

$$\bigcup_{p \in \mathcal{P}} L_{p,0} = \mathbb{Z} \setminus \{-1, 1\}$$

is closed, hence  $\{-1, 1\}$  is open. However, an open set  $G$  is the union of  $L_{a,b}$  which is infinite, so  $G$  is infinite, leading to contradiction.

Quote:

As for everything else, so for a mathematical theory: beauty can be perceived but not explained.

—A. Cayley

## 6 Homework 6: Takagi Function

### 6.1 PSA: Calculating Derivatives

A1) Consider the function

$$f : \mathbb{R} \rightarrow \mathbb{R}^n, x \mapsto f(x) = (f_1(x), \dots, f_n(x)).$$

Prove that  $f$  is differentiable at  $x_0$  iff every  $f_k$  is differentiable at  $x_0$  and

$$f'(x) = (f'_1(x), \dots, f'_n(x)).$$

Proof: For any  $h \in \mathbb{R}$ ,

$$\left\| \frac{f(x+h) - f(x)}{h} - (f'_1(x), \dots, f'_n(x)) \right\|_2 \leq n \max_{1 \leq k \leq n} \left\| \frac{f_k(x+h) - f_k(x)}{h} - f'_k(x) \right\| \rightarrow 0.$$

Therefore  $f'(x) = (f'_1(x), \dots, f'_n(x))$ .

### A2) Consider the function

$$f : \mathbb{R} \rightarrow \mathbb{C}, x \mapsto e^{ix}.$$

Prove by definition,  $f'(0) = i$  and  $(e^{ix})' = ie^{ix}$ .

Proof: For any  $h \in \mathbb{R}$ ,

$$\left| \frac{f(h) - f(0)}{h} - i \right| = \left| \frac{e^{ih} - ih - 1}{h} \right| \leq \sum_{n=2}^{\infty} \left| \frac{1}{h} \frac{(ih)^n}{n!} \right| \leq |h| \rightarrow 0.$$

Therefore  $f'(0) = i$ . For any  $x \in \mathbb{R}$ ,

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} e^{ix} \frac{f(h) - f(0)}{h} = ie^{ix}.$$

Hence  $(e^{ix})' = ie^{ix}$ .

### A3) Calculate the derivatives of $\sin x$ and $\cos x$ .

Solution:  $\sin x = (e^{ix} - e^{-ix})/2i$ , so  $(\sin x)' = (e^{ix} + e^{-ix})/2 = \cos x$ . Likewise  $(\cos x)' = -\sin x$ .

### A4) Prove Faà di Bruno's formula for $n = 3$ .

Proof:

$$\begin{aligned} \frac{d}{dx}(f \circ g) &= f'(g) \cdot g'. \\ \frac{d^2}{dx^2}(f \circ g) &= f'(g) \cdot g'' + f''(g) \cdot (g')^2. \\ \frac{d^3}{dx^3}(f \circ g) &= f'(g) \cdot g''' + f''(g) \cdot g'' \cdot g' + f'''(g) \cdot (g')^3 + f''(g) \cdot 2g'g''. \end{aligned}$$

### A5) Define the map

$$E : \mathbb{R} \rightarrow \mathbb{C} = \mathbb{R}^2, \theta \mapsto (\cos \theta, \sin \theta).$$

Prove that the points in  $\mathbf{S}^1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$  can be written in the form  $(\sin \theta, \cos \theta)$ , i.e.  $E(\mathbb{R}) = \mathbf{S}^1$ . Calculate  $E'(\theta)$  and show that Rolle's mean-value theorem is invalid for  $E$ .

Proof: Obviously  $E(\mathbb{R}) \subset \mathbf{S}^1$ . Consider any  $(x, y) \in \mathbf{S}^1$ , then  $x \in [-1, 1]$ . Note that  $\cos 0 = 1, \cos \pi = -1$ , hence there exists  $\theta \in [0, \pi]$  such that  $\cos \theta = x$ , and  $|\sin \theta| = |y|$ . If  $\sin \theta = y$  then  $(x, y) = (\cos \theta, \sin \theta) \in E(\mathbb{R})$ . Otherwise  $(x, y) = (\cos(-\theta), \sin(-\theta)) \in E(\mathbb{R})$ , therefore  $E(\mathbb{R}) = \mathbf{S}^1$ .

By A1) and A3),  $E'(\theta) = (-\sin \theta, \cos \theta)$ . Since  $E'(\theta) \neq 0$  for all  $\theta \in \mathbb{R}$  and  $E'(\theta) = E'(\theta + 2\pi)$ , Rolle's mean-value theorem is invalid.

**A6) Calculate the derivatives of the following functions:**

(1)  $f(x) = a^x, a > 0.$

$$f'(x) = (e^{x \log a})' = a^x \log a.$$

(2)  $f(x) = \arcsin x.$

Let  $y = \arcsin x$ , then  $x = \sin y$ , so  $1 = \cos y \cdot y'$ , hence

$$f'(x) = \frac{1}{\cos(\arcsin x)} = \frac{1}{\sqrt{1-x^2}}.$$

(3)  $f(x) = \arctan x.$

Let  $y = \arctan x$ , then  $x = \tan y$ , so  $1 = \sec^2 y \cdot y'$ , hence

$$f'(x) = \cos^2 y = \frac{1}{1+x^2}.$$

(4)  $f(x) = x^{x^x}, x > 0.$

Let  $y = x^x, z = x^y$ , then  $\log y = x \log x$ , so  $y'/y = \log x + 1, y' = x^x(1 + \log x)$ .  $\log z = y \log x$ , so  $z'/z = y' \log x + y/x = x^x \log x(1 + \log x) + x^{x-1}$ . Therefore

$$f'(x) = x^{x^x} \cdot x^x \cdot (\log x + \log^2 x + x^{-1}).$$

(5)  $f(x) = \log(\log(\log x)).$

$$f'(x) = \frac{(\log \log x)'}{\log \log x} = \frac{1}{x \log x \log \log x}.$$

(6)  $f(x) = \sqrt{x + \sqrt{x + \sqrt{x}}}.$

$$\begin{aligned} f'(x) &= \frac{(x + \sqrt{x + \sqrt{x}})'}{2\sqrt{x + \sqrt{x + \sqrt{x}}}} = \left(1 + \frac{1 + \frac{1}{2\sqrt{x}}}{2\sqrt{x + \sqrt{x}}}\right) / 2\sqrt{x + \sqrt{x + \sqrt{x}}} \\ &= \frac{2\sqrt{x + \sqrt{x}} + 1 + 1/2\sqrt{x}}{4\sqrt{x + \sqrt{x}}\sqrt{x + \sqrt{x + \sqrt{x}}}}. \end{aligned}$$

(7)  $f(x) = |x|.$

If  $x > 0$ ,  $f'(x) = (x)' = 1$ . If  $x < 0$ ,  $f'(x) = (-x)' = -1$ . If  $x = 0$ ,  $f$  is not differentiable at  $x$ .

(8)  $f(x) = \log|x|.$

If  $x > 0$ ,  $f'(x) = \frac{1}{x}$ . If  $x < 0$ ,  $f'(x) = -\frac{1}{x}$ . If  $x = 0$ ,  $f$  is not differentiable at  $x$ .

(9)

$$f(x) = \begin{cases} x^n \sin \frac{1}{x}, & x \neq 0, \\ 0, & x = 0. \end{cases} \quad n = 1, 2, \dots$$

For  $x \neq 0$ ,  $f'(x) = nx^{n-1} \sin \frac{1}{x} - x^{n-2} \cos \frac{1}{x}$ . When  $x = 0$ ,

$$f'(0) = \lim_{h \rightarrow 0} h^{n-1} \sin \frac{1}{h} = \begin{cases} 0, & n \geq 2; \\ \text{diverges}, & n = 1. \end{cases}$$

**A7) Calculate  $f^{(3)}(x)$ :**

(1)  $f(x) = \log(x+1)$ .

$$\frac{d^3}{dx^3} \log(x+1) = \frac{2}{(x+1)^3}.$$

(2)  $f(x) = x^{-1} \log x$ .

$$\frac{d^3}{dx^3} \frac{\log x}{x} = \frac{11 - 6 \log x}{x^4}.$$

(3)  $f(x) = \frac{x^m}{1-x}$ ,  $m \in \mathbb{Z}_{\geq 0}$ .

$$\frac{d^3}{dx^3} \frac{x^m}{1-x} = \frac{(m-2)(m-1)mx^{m-3}}{1-x} + \frac{3(m-1)mx^{m-2}}{(1-x)^2} + \frac{6mx^{m-1}}{(1-x)^3} + \frac{6x^m}{(1-x)^4}.$$

(4)  $f(x) = x^m e^x$ ,  $m \in \mathbb{Z}_{\geq 0}$ .

$$\frac{d^3}{dx^3} (x^m e^x) = e^x x^{m-3} (m^3 + 3m^2(x-1) + m(3x^2 - 3x + 2) + x^3).$$

(5)  $f(x) = e^{ax} \sin(bx)$ ,  $a, b \in \mathbb{R}$ .

$$\frac{d^3}{dx^3} (e^{ax} \sin(bx)) = e^{ax} ((3a^2b - b^3) \cos(bx) + a(a^2 - 3b^2) \sin(bx)).$$

(6)  $f(x) = e^{-x^2}$ .

$$\frac{d^3}{dx^3} e^{-x^2} = -4e^{-x^2} x(2x^2 - 3).$$

**A8)  $f'(x_0) > 0$  does not imply  $f$  is increasing in a neighborhood of  $x_0$ : consider**

$$f(x) = \begin{cases} x + 2x^2 \sin \frac{1}{x}, & x \neq 0; \\ 0, & x = 0. \end{cases}$$

Prove that  $f'(0) > 0$  but for any  $\varepsilon > 0$ ,  $f$  is not monotonic on  $(-\varepsilon, \varepsilon)$ .

Proof:

$$f'(0) = \lim_{h \rightarrow 0} 1 + 2h \sin \frac{1}{h} = 1 > 0.$$

However, for any  $n \in \mathbb{N}$ , let  $x_n = \frac{1}{(2n+1/2)\pi}$ ,  $y_n = \frac{1}{(2n-1/2)\pi}$ , then

$$f(x_n) = x_n + 2x_n^2, f(y_n) = y_n - 2y_n^2.$$

Note that  $0 < x_n < y_n$ , but

$$f(x_n) - f(y_n) = 2x_n^2 + 2y_n^2 - \pi x_n y_n > 0,$$

i.e.  $f(x_n) > f(0), f(y_n)$ , therefore  $f$  is not monotonic on any  $(-\varepsilon, \varepsilon)$ .

**A9)**  $A \in \mathbf{M}_n(\mathbb{R})$ , calculate

$$\left. \frac{d}{dx} \right|_{x=0} \det(\mathbf{I}_n + xA).$$

Solution: Let  $\Phi(x) = I_n + xA$ , then  $\Phi(0) = I_n$ . Denote  $\Phi(t) = (\varphi_1(t), \dots, \varphi_n(t))$ . Note that  $\det$  is a multi-linear function for  $n$  rows, hence by Euler's formula:

$$\frac{d}{dt} \det \Phi(t) = \det(\varphi'_1(t), \varphi_2(t), \dots, \varphi_n(t)) + \dots + \det(\varphi_1(t), \varphi_2(t), \dots, \varphi'_n(t)).$$

When  $t = 0$ , the formula becomes

$$\left. \frac{d}{dt} \right|_{t=0} \det \Phi(t) = \varphi'_{1,1} + \dots + \varphi'_{n,n} = \text{tr } \Phi'(0) = \text{tr } A.$$

**A10)** Prove that the derivation of odd functions are even, and that of even functions are odd.

Proof: If  $f$  is an odd function then

$$f'(-x) = \lim_{h \rightarrow 0} \frac{f(-x+h) - f(-x)}{h} = \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h} = f'(x),$$

so  $f'$  is even. If  $f$  is an even function then

$$f'(-x) = \lim_{h \rightarrow 0} \frac{f(-x+h) - f(-x)}{h} = - \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h} = -f'(x),$$

so  $f'$  is odd.

**A11)** Prove that

$$f(x) = \begin{cases} 1/q, & x = \frac{p}{q} \in \mathbb{Q}, q \geq 1, \gcd(p, q) = 1; \\ 0, & x \in \mathbb{Q}^C. \end{cases}$$

is nowhere differentiable on  $\mathbb{R}$ .

Proof: For any  $x \in \mathbb{Q}$ ,  $f(x) \neq 0$ , but for any  $\varepsilon > 0$ , there exists  $y \in (x - \varepsilon, x + \varepsilon) \cap \mathbb{Q}^C$ , such that  $f(y) = 0$ . Therefore  $f$  is not continuous at  $x$ , and clearly not differentiable.

Consider any  $x \in \mathbb{Q}^C$ , there is a sequence of irrational numbers  $\{y_n\}_{n \geq 1}$  that converges to  $x$ , then

$$\lim_{n \rightarrow \infty} \frac{f(x) - f(y_n)}{x - y_n} = 0.$$

Choose any sequence of rational numbers  $\{r_n = p_n/q_n\}_{n \geq 1}$  that converges to  $x$ , then

$$\lim_{n \rightarrow \infty} \frac{f(x) - f(r_n)}{x - r_n} = \lim_{n \rightarrow \infty} \frac{1}{xq_n - p_n} = \infty.$$

Therefore  $f$  is nowhere differentiable on  $\mathbb{R}$ .



## 6.2 PSB

**B1) Define the hyperbolic functions:**

$$\sinh x = \frac{e^x - e^{-x}}{2}, \cosh x = \frac{e^x + e^{-x}}{2}, \tanh x = \frac{\sinh x}{\cosh x}.$$

1. Prove that

$$(1) \cosh^2 x - \sinh^2 x = 1$$

$$\text{Proof: } \cosh^2 x - \sinh^2 x = \frac{e^{2x} + e^{-2x} + 2}{4} - \frac{e^{2x} + e^{-2x} - 2}{4} = 1.$$

$$(2) \sinh(x + y) = \sinh x \cosh y + \cosh x \sinh y.$$

$$\text{Proof: } \sinh x \cosh y + \cosh x \sinh y = \frac{e^{x+y} - e^{y-x} + e^{x-y} - e^{-x-y}}{4} + \frac{e^{x+y} - e^{x-y} + e^{y-x} - e^{-x-y}}{4} = \sinh(x + y)$$

$$(3) \cosh(x + y) = \cosh x \cosh y + \sinh x \sinh y.$$

Proof: Same as (2).

$$(4) \tanh(x + y) = \frac{\tanh x + \tanh y}{1 + \tanh x \tanh y}.$$

$$\text{Proof: } \tanh(x + y) = \frac{\sinh x \cosh y + \cosh x \sinh y}{\cosh x \cosh y + \sinh x \sinh y} = \frac{\tanh x + \tanh y}{1 + \tanh x \tanh y}.$$

2. Calculate  $\sinh'(x)$ ,  $\cosh'(x)$  and  $\tanh'(x)$ .

$$\text{Solution: } \sinh'(x) = \cosh x, \cosh'(x) = \sinh x, \tanh'(x) = \frac{1}{\cosh^2 x}.$$

3. Prove that  $\sinh : \mathbb{R} \rightarrow \mathbb{R}$  has an inverse  $\operatorname{arcsinh} : \mathbb{R} \rightarrow \mathbb{R}$  and calculate  $\operatorname{arcsinh}'(x)$ .

Proof:  $\sinh'(x) = \cosh x > 0$ , so  $\sinh$  is monotonically increasing over  $\mathbb{R}$ . Also  $\lim_{x \rightarrow \infty} \sinh x = \infty$ ,  $\lim_{x \rightarrow -\infty} \sinh x = -\infty$ , therefore  $\sinh : \mathbb{R} \rightarrow \mathbb{R}$  is a bijection and hence has an inverse.

$$\operatorname{arcsinh}'(x) = \frac{1}{\sqrt{1+x^2}}.$$

**B2)  $a, b \in \mathbb{R}$ ,  $a > 0$ . Consider  $f : [-1, 1] \rightarrow \mathbb{R}$ , where**

$$f(x) = \begin{cases} x^a \sin(x^{-b}), & x \neq 0, \\ 0, & x = 0. \end{cases}$$

Prove that

1.  $f \in C([-1, 1])$  iff  $a > 0$ ;

Proof:  $f \in C([-1, 1])$  iff  $\lim_{x \rightarrow 0} x^a \sin(x^{-b}) = 0$ . If  $a > 0$  then  $|x^a \sin(x^{-b})| \leq |x|^a \rightarrow 0$ . If  $a < 0$  then let  $x = ((2n + 1/2)\pi)^{-1/b}$ , when  $n \rightarrow \infty$ ,  $x \rightarrow 0$  but  $|x^a \sin(x^{-b})| \rightarrow \infty$ . If  $a = 0$ , then let  $x = ((2n + 1/2)\pi)^{-1/b}$ ,  $|x^a \sin(x^{-b})| = 1$ . Therefore  $f \in C([-1, 1])$  iff  $a > 0$ .

2.  $f$  is differentiable at 0 iff  $a > 1$ ;

Proof:  $f$  is differentiable at 0 iff  $\lim_{x \rightarrow 0} x^{-a} \sin(x^{-b})$  exists. By 1 we know that  $a > 1$ . ( $a = 1$  is invalid since  $x = (2n\pi)^{-1/b}$  and  $x = ((2n + 1/2)\pi)^{-1/b}$  converge to different values.)

3.  $f'$  is bounded on  $[-1, 1]$  iff  $a \geq 1 + b$ ;

Proof:  $f'(x) = ax^{a-1} \sin(x^{-b}) + x^a \cos(x^{-b})(-b)x^{-b-1}$  is bounded iff  $x^{a-1}$  and  $x^{a-b-1}$  are bounded, i.e.  $a \geq 1 + b$ .

4.  $f \in C^1([-1, 1])$  iff  $a > 1 + b$ ;

Proof:  $f \in C^1([-1, 1])$  iff  $f'(0) = 0 = \lim_{x \rightarrow 0} f'(x)$ . By 1 we know it is equivalent to  $a > 1 + b$ .

5.  $f'$  is differentiable at 0 iff  $a > 2 + b$ ;

6.  $f''$  is bounded on  $[-1, 1]$  iff  $a \geq 2 + 2b$ ;

7.  $f \in C^2([-1, 1])$  iff  $a > 2 + 2b$ .

Proof: 5,6,7 are exactly the same as 2,3,4.

### 6.3 PSC

If  $f$  satisfy  $\lim_{x \rightarrow x_0} f(x) = 0$  near  $x_0$ , we call  $f$  an infinitesimal when  $x \rightarrow x_0$ . Likewise when  $\lim_{x \rightarrow x_0} f(x) = +\infty$  or  $\lim_{x \rightarrow x_0} f(x) = -\infty$ , we call  $f$  an infinite quantity when  $x \rightarrow x_0$ .

Suppose  $f, g$  are both infinitesimal when  $x \rightarrow x_0$ , and  $g(x)$  does not vanish near  $x_0$ . We introduce the notations

- if  $\lim_{\substack{x \rightarrow x_0 \\ x \rightarrow x_0}} \frac{f(x)}{g(x)} = 0$ , we say  $f$  is an infinitesimal of higher order than  $g$ , and denote  $f(x) = o(g(x))$ ,
- If  $\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \neq 0$ , we say  $f$  and  $g$  are of the same order;
- If  $= 1$ , denote  $f \sim g, x \rightarrow x_0$ ;
- If  $\limsup_{x \rightarrow x_0} \left| \frac{f(x)}{g(x)} \right| < +\infty$ , denote  $f(x) = O(g(x)), x \rightarrow x_0$ .

**C1) Suppose  $a(x) = o(1)$  when  $x \rightarrow x_0$ , prove that:**

(1)  $o(a) + o(a) = o(a)$

Proof: If  $f, g = o(a)$ , then

$$\lim_{x \rightarrow x_0} \frac{f(x) + g(x)}{a(x)} = \lim_{x \rightarrow x_0} \frac{f(x)}{a(x)} + \lim_{x \rightarrow x_0} \frac{g(x)}{a(x)} = 0,$$

hence  $f + g = o(a)$ .

(2)  $co(a) = o(ca), c \in \mathbb{R}$

Proof: If  $f = o(a)$ , then

$$\lim_{x \rightarrow x_0} \frac{cf(x)}{a(x)} = c \lim_{x \rightarrow x_0} \frac{f(x)}{a(x)} = 0,$$

hence  $cf = o(a) = o(ca)$ .

(3)  $o(a)^k = o(a^k)$

Proof: If  $f = o(a)$  then

$$\lim_{x \rightarrow x_0} \frac{f(x)^k}{a(x)^k} = \left( \lim_{x \rightarrow x_0} \frac{f(x)}{a(x)} \right)^k = 0,$$

hence  $f^k = o(a^k)$ .

(4)  $1/(1+a) = 1-a+o(a)$

Proof:

$$\lim_{x \rightarrow x_0} \frac{1/(1+a) - 1 + a}{a(x)} = \lim_{x \rightarrow x_0} \frac{a(x)}{1+a(x)} = 0,$$

hence  $1/(1+a) = 1-a+o(a)$ .

**C2) Suppose  $f, g$  are infinitesimals when  $x \rightarrow x_0$ , then**

1. Prove that  $f \sim g \iff f(x) - g(x) = o(g(x)), x \rightarrow x_0$ .

Proof:  $f \sim g \iff \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 1 \iff \lim_{x \rightarrow x_0} \frac{f(x) - g(x)}{g(x)} = 0$ , i.e.  $f(x) - g(x) = o(g(x))$ .

2. If  $f \sim cg^k$ , we call  $cg^k$  the leading term of  $f$ . Find the leading terms of the following functions, compared to  $x - x_0$  or  $x$ :
- (1)  $1/\sin \pi x, x \rightarrow 1$ .  
 $\frac{1}{\sin \pi x} = -\frac{1}{\pi(x-1)} + o(1)$ .
  - (2)  $\sqrt{1+x} - \sqrt{1-x}, x \rightarrow 0$ .  
 $\sqrt{1+x} - \sqrt{1-x} = x + o(x)$ .
  - (3)  $\sin(\sqrt{1+\sqrt{1+\sqrt{x}}} - \sqrt{2}), x \rightarrow 0^+$ .  
 $= \frac{\sqrt{2}x^{1/2}}{8} + o(x^{1/2})$ .
  - (4)  $\sqrt{1+\tan x} - \sqrt{1-\sin x}, x \rightarrow 0$ .  
 $= x + o(x)$ .
  - (5)  $\sqrt{x + \sqrt{x + \sqrt{x}}}, x \rightarrow 0^+$ .  
 $= x^{1/8} + o(x^{1/8})$ .
  - (6)  $\sqrt{x + \sqrt{x + \sqrt{x}}}, x \rightarrow \infty$ .  
 $= \sqrt{x} + o(\sqrt{x})$ .
3. Suppose  $f \sim cx^k, x \rightarrow 0$ , i.e.  $f(x) = cx^k + o(x^k)$ . If  $f(x) - c^k$  has a leading term  $c'x^{k'}$ , we denote  $f(x) = cx^k + c'x^{k'} + o(x^{k'})$ . Expand the following terms to  $o(x^2)$ :
- (1)  $\sqrt{1+x} - 1$ .  
 $\sqrt{1+x} - 1 = \frac{1}{2}x - \frac{1}{8}x^2 + o(x^2)$ .
  - (2)  $(1+x)^{1/m} - 1, m \in \mathbb{Z}_{\geq 1}$ .  
 $(1+x)^{1/m} - 1 = \frac{1}{m}x - \frac{m-1}{2m^2}x^2 + o(x^2)$ .

## 6.4 PST: Takagi Function

Define  $\psi : [0, 1] \rightarrow \mathbb{R}$  as

$$\psi(x) = \begin{cases} x, & 0 \leq x < \frac{1}{2}; \\ 1-x, & \frac{1}{2} \leq x \leq 1. \end{cases}$$

For  $x \in \mathbb{R}$ , let  $\psi(x) = \psi(\{x\})$ , then  $\psi \in C(\mathbb{R})$ .

Define the Takagi function  $T : \mathbb{R} \rightarrow \mathbb{R}$  as follows:

$$T(x) = \sum_{k=0}^{\infty} \frac{1}{2^k} \psi(2^k x),$$

and the partial sum  $T_n(x) = \sum_{k=0}^n \frac{1}{2^k} \psi(2^k x)$ .

**T1) Prove that  $T(x)$  is a well-defined bounded continuous function on  $\mathbb{R}$ .**

Proof: Note that  $\psi(x) \in [0, 1/2]$  so the series  $\sum_{k=0}^{\infty} 2^{-k} \psi(2^k x)$  converges absolutely, and hence  $T(x)$  is well-defined and bounded by  $T(x) \in [0, 1]$ .

**T2) For  $x \in [0, 1]$ , suppose  $x = \sum_{n=1}^{\infty} a_n 2^{-n}$  is the binary form of  $x$ . Let  $v_n = \sum_{k=1}^n a_k$ , and  $\sigma_n(y) = a_n + (1 - 2a_n)y$ , where  $y \in \{0, 1\}$ . Prove that**

$$\psi(2^m x) = \sum_{n=1}^{\infty} \frac{\sigma_{m+1}(a_{m+n})}{2^n}.$$

Proof:

$$\psi(2^m x) = \psi\left(\sum_{n=1}^{\infty} a_n 2^{m-n}\right) = \psi\left(\sum_{n=m+1}^{\infty} a_n 2^{m-n}\right) = \begin{cases} \sum_{n=1}^{\infty} a_{m+n} 2^{-n}, & a_{m+1} = 0; \\ 1 - \sum_{n=1}^{\infty} a_{m+n} 2^{-n}, & a_{m+1} = 1. \end{cases}$$

Therefore

$$\psi(2^m x) = \sigma_n \left( \sum_{n=1}^{\infty} a_{m+n} 2^{-n} \right) = \sum_{n=1}^{\infty} \sigma_{m+1}(a_{m+n}) 2^{-n}.$$

**T3)**  $x = \sum_{n=1}^{\infty} a_n 2^{-n} \in [0, 1]$ , **prove that**

$$T(x) = \sum_{n=1}^{\infty} \frac{(1 - a_n)v_n + a_n(n - v_n)}{2^n}.$$

Proof: By T2),

$$T(x) = \sum_{m=0}^{\infty} \sum_{n=1}^{\infty} \sigma_{m+1}(a_{m+n}) 2^{-m-n} = \sum_{n=1}^{\infty} \sum_{m=0}^{n-1} \sigma_{m+1}(a_n) 2^{-n} = \sum_{n=1}^{\infty} \frac{(1 - a_n)v_n + a_n(n - v_n)}{2^n}.$$

**T4)** Suppose  $x_0 = k_0 2^{-m_0} \in [0, 1]$ , where  $k_0 \in \mathbb{Z}_{\geq 1}$  is odd,  $m_0 \in \mathbb{Z}_{\geq 0}$ . Let  $h_n = 2^{-n}$ , where  $n \in \mathbb{Z}_{\geq m_0}$ . **Prove that the sequence  $\left\{ \frac{T(x+h_n) - T(x)}{h_n} \right\}_{n \geq m_0}$  does not converge.**

Proof: By T3),

$$\frac{T(x+h_n) - T(x)}{h_n} = \frac{1}{h_n} \left( \frac{n - v_n}{2^n} - \frac{v_n}{2^n} \right) = n - 2 \sum_{k=1}^n a_k = n - 2 - 2S_2(k_0) \rightarrow \infty.$$

**T5)**  $f : I \rightarrow \mathbb{R}$ , where  $I$  is an open interval. If  $f$  is differentiable at  $a$ , **prove that**

$$\lim_{(h, h') \rightarrow (0, 0), h, h' > 0} \frac{f(a+h) - f(a-h')}{h+h'} = f'(a).$$

i.e. it converges for any sequence  $(h_n, h'_n) \rightarrow (0, 0)$ ,  $h_n, h'_n > 0$ .

Proof: Consider any sequence  $(h_n, h'_n) \rightarrow (0, 0)$ , then

$$\frac{f(a+h) - f(a-h')}{h+h'} = \frac{f(a+h) - f(a)}{h} \cdot \frac{h}{h+h'} + \frac{f(a) - f(a-h')}{h'} \cdot \frac{h'}{h+h'} \rightarrow f'(a).$$

**T6)** Same as T5), if  $f \in C^1(I)$ ,  $a \in I$ , **prove that**

$$\lim_{(h, h') \rightarrow (0, 0), h+h' \neq 0} \frac{f(a+h) - f(a-h')}{h+h'} = f'(a).$$

Proof: For any  $h+h' \neq 0$ , there exists  $\xi \in [a, a+h]$  and  $\eta \in [a-h', a]$  such that  $f(a+h) = f(a) + hf'(\xi)$  and  $f(a-h') = f(a) - h'f'(\eta)$ , then

$$\left| \frac{f(a+h) - f(a-h')}{h+h'} - f'(a) \right| \leq \frac{h}{h+h'} |f'(\xi) - f'(a)| + \frac{h'}{h+h'} |f'(\eta) - f'(a)| \rightarrow 0.$$

Hence

$$\lim_{(h, h') \rightarrow (0, 0), h+h' \neq 0} \frac{f(a+h) - f(a-h')}{h+h'} = f'(a).$$

**T7)** Suppose  $x \in [0, 1]$ , such that for any  $n \in \mathbb{N}$ ,  $2^n x \notin \mathbb{Z}$ . For every  $n \in \mathbb{N}$ , define  $\{h_n\}_{n \geq 1}$  and  $\{h'_n\}_{n \geq 1}$  as follows:

$$\lfloor 2^n x \rfloor = 2^n(x - h'_n), \quad \lfloor 2^n x \rfloor + 1 = 2^n(x + h_n).$$

Prove that for an arbitrary  $n$ ,  $h_n + h'_n = 2^{-n}$  and for every integer  $1 \leq n-1$ , the interval  $(2(x - h'_n), 2(x + h_n))$  does not include integers or half-integers.

Proof:  $1 = 2^n(x + h_n) - 2^n(x - h'_n) = 2^n(h_n + h'_n)$ , hence  $h_n + h'_n = 2^{-n}$ . For any integer  $1 \leq n-1$ ,  $2(x - h'_n) = \lfloor 2^n x \rfloor \cdot 2^{-n}$  and  $2(x + h_n) = (\lfloor 2^n x \rfloor + 1)2^{-n}$ . Since  $n-1 \geq 1$ , the interval does not include integers or half-integers.

**T8)** Follow the notations of T7), prove that the sequence  $\left\{ \frac{T(x+h_n) - T(x-h'_n)}{h_n + h'_n} \right\}_{n \geq 1}$  diverges.

Proof: Let  $t = \lfloor 2^n x \rfloor$ , then

$$a_n = \frac{T(x + h_n) - T(x - h'_n)}{h_n + h'_n} = \sum_{k=0}^{n-1} 2^{n-k} \left( \psi\left(\frac{t+1}{2^{n-k}}\right) - \psi\left(\frac{t}{2^{n-k}}\right) \right).$$

Since the interval  $(2^{k-n}(t+1), 2^{k-n}t)$  does not contain any integers or half-integers,  $2^{n-k}(\psi(2^{k-n}(t+1)) - \psi(2^{k-n}t)) \in \{-1, 1\}$ , so  $a_n \in \mathbb{Z}$  and  $n, a_n$  have the same parity. Therefore the sequence  $\{a_n\}_{n \geq 1}$  diverges.

**T9)** Prove that  $T(x)$  is continuous but nowhere differentiable on  $\mathbb{R}$ .

Proof: For any  $x \in [0, 1]$ , if  $x = k_0 \cdot 2^{-m_0}$  as in T4), by T4) the sequence  $\left\{ \frac{T(x + h_n) - T(x)}{h_n} \right\}$  diverges, hence  $T$  is not differentiable at  $x$ . Otherwise for any  $n \in \mathbb{N}$ ,  $2^n x \notin \mathbb{Z}$ . Define  $\{h_n\}_{n \geq 1}$  and  $\{h'_n\}_{n \geq 1}$  as in T7), then by T8), the sequence  $\left\{ \frac{T(x + h_n) - T(x - h'_n)}{h_n + h'_n} \right\}_{n \geq 1}$  diverges. Combined with T5) we know that  $T$  is not differentiable at  $x$ . Therefore  $T$  is nowhere differentiable on  $\mathbb{R}$ , since  $T$  is periodic.

For any  $x, y$  in  $\mathbb{R}$ ,

$$|T(x) - T(y)| \leq \sum_{k=0}^N 2^{-k} |T(2^k x) - T(2^k y)| + \sum_{k=N+1}^{\infty} 2^{-k} \leq 2 \max_{0 \leq k \leq N} |T(2^k x) - T(2^k y)| + 2^{-N}.$$

Hence for any  $N > 0$ , when  $\varepsilon \rightarrow 0$ ,  $|T(x) - T(x + \varepsilon)| \leq 2^{1-N} \rightarrow 0$ , so  $T$  is (uniformly) continuous on  $\mathbb{R}$ .

**T10)** Prove that

$$T(x) = \begin{cases} 2x + \frac{T(4x)}{4}, & 0 \leq x < \frac{1}{4}; \\ \frac{1}{2} + \frac{T(4x-1)}{4}, & \frac{1}{4} \leq x < \frac{1}{2}; \\ \frac{1}{2} + \frac{T(4x-2)}{4}, & \frac{1}{2} \leq x < \frac{3}{4}; \\ 2 - 2x + \frac{T(4x-3)}{4}, & \frac{3}{4} \leq x \leq 1. \end{cases}$$

Proof: If  $0 \leq x < 1/4$ , then

$$T(x) = \psi(x) + \psi(2x)/2 + \sum_{k=2}^{\infty} \psi(2^k x) 2^{-k} = 2x + \frac{T(4x)}{4}.$$

The other cases are exactly the same.

**T11)** Let  $\Gamma = \{(x, T(x)) : 0 \leq x \leq 1\} \subset \mathbb{R}^2$ . Define  $\Phi_i : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$\begin{aligned}\Phi_0 \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 1/4 & 0 \\ 1/2 & 1/4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}, & \Phi_1 \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 1/4 & 0 \\ 0 & 1/4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1/4 \\ 1/2 \end{pmatrix}, \\ \Phi_2 \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 1/4 & 0 \\ 0 & 1/4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix}, & \Phi_3 \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 1/4 & 0 \\ 0 & 1/4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 3/4 \\ 1/2 \end{pmatrix}.\end{aligned}$$

Prove that  $\Phi_i$  maps  $\Gamma$  to  $\{(x, T(x)) : x \in [\frac{i}{4}, \frac{i+1}{4}]\}$ .

Proof: Consider  $(x, T(x)) \in \Gamma$ , then by T10),

$$\Phi_0 \begin{pmatrix} x \\ T(x) \end{pmatrix} = \begin{pmatrix} x/4 \\ x/2 + T(x)/4 \end{pmatrix} = \begin{pmatrix} x/4 \\ T(x/4) \end{pmatrix}.$$

Hence  $\Phi_0(\Gamma) = \{(x, T(x)) : x \in [0, 1/4]\}$ . The cases  $i = 1, 2, 3$  are similar.

**T12)** Let  $S_0 = \{(x, y) \in \mathbb{R}^2 : 0 \leq x \leq 1, 0 \leq y \leq 1\}$ . For every  $n \geq 0$ , define  $S_{n+1} = \bigcup_{k=0}^3 \Phi_k(S_n)$ . Prove that  $S_n$  is a sequence of monotonically decreasing compact sets and  $\Gamma = \bigcap_{n \geq 0} S_n$ .

Proof: Let  $S_n(x) = \{y \in [0, 1] : (x, y) \in S_n\}$ . We prove by induction that  $S_n \subset S_{n-1}$  and  $S_n(x)$  is a closed interval containing  $T(x)$  for any  $x \in [0, 1]$ . The base  $n = 0$  is trivial. Suppose  $S_n \subset S_{n-1}$  and  $S_n(x)$  is a closed interval containing  $T(x)$ , then consider  $S_{n+1}$ . Note that  $\Phi_k(S_n)$  are disjoint, since for any  $(x, y) \in \Phi_k(S_n)$ ,  $x \in [k/4, (k+1)/4]$ . Hence for any  $x \in [0, 1/4]$ ,  $S_{n+1}(x) = \{y : (x, y) = \Phi_0(4x, z), z \in S_n(4x)\} = \{2x + z/4 : z \in S_n(4x)\}$  is a closed interval containing  $T(x) = 2x + T(4x)/4$ . By the induction hypothesis  $S_n(x) = \{2x + z/4 : z \in S_{n-1}(4x)\}$  and  $S_n(4x) \subset S_{n-1}(4x)$  so  $S_{n+1}(x) \subset S_n(x)$ . The case  $x \in [1/4, 1]$  is similar. Therefore  $S_{n+1} \subset S_n$  and  $S_{n+1}$  is compact, so by induction  $S_n \subset S_{n-1}$  for all  $n > 0$  and  $S_n$  is compact.

Clearly  $\Gamma \subset \bigcap_{n \geq 0} S_n$ , so it suffices to show that  $|S_n(x)| \rightarrow 0$  for all  $x \in [0, 1]$ . From the proof above we get  $\sup_{x \in [0, 1]} |S_n(x)| \leq \sup_{x \in [0, 1]} |S_{n-1}(x)|/4$ , hence  $|S_n(x)| \rightarrow 0$ , therefore

$$\Gamma = \bigcap_{n \geq 0} S_n.$$

**T13)** Prove that  $\sup_{x \in \mathbb{R}} T(x) \geq \frac{2}{3}$ .

Proof: For any  $(x, y) \in \Gamma$ , by T11) we know that  $(x/4 + 1/4, y/4 + 1/2) \in \Gamma$ , hence if  $a = \sup_{x \in \mathbb{R}} T(x)$  then  $a \geq a/4 + 1/2$ , i.e.  $a \geq 2/3$ .

**T14)** Find a  $c \in [0, 1]$  such that  $T(c) = \frac{2}{3}$ .

Solution: Consider  $c = 1/3$ , then by T10),  $T(c) = T(c)/4 + 1/2$ , hence  $T(c) = \frac{2}{3}$ .

**T15)** For  $x \in [0, 1]$ , suppose  $x = \sum_{n=1}^{\infty} b_n 4^{-n}$ , where  $b_n \in \{0, 1, 2, 3\}$ . Prove that

$$\left\{x \in [0, 1] : T(x) = \frac{2}{3}\right\} = \left\{x \in [0, 1] : x = \sum_{n=1}^{\infty} b_n 4^{-n}, b_n \in \{1, 2\}\right\}.$$

Proof: If  $x = \sum_{n=1}^{\infty} b_n 4^{-n}$ , where  $b_n \in \{1, 2\}$ , then by T10),

$$T(x) = \frac{1}{2} + \frac{1}{4}T\left(\sum_{n=1}^{\infty} b_{n+1} 4^{-n}\right) = \cdots = \frac{1}{2}\left(1 + \frac{1}{4} + \frac{1}{4^2} + \cdots\right) = \frac{2}{3}.$$

Otherwise take the least  $n$  such that  $b_n \in \{0, 3\}$ , denote  $y = \sum_{k=1}^{\infty} b_{n+k-1} 4^{-n}$ , then

$$T(x) = \frac{1}{2} \left( 1 + \frac{1}{4} + \cdots + \frac{1}{4^{n-2}} \right) + \frac{\min\{2y, 2-2y\}}{4^{n-1}} + \frac{1}{4^n} T(4y - b_n) < \frac{2}{3},$$

since  $T(4y - b_n) \leq 2/3$  and  $\min\{2y, 2-2y\} < 1/2$ . Therefore

$$\left\{ x \in [0, 1] : T(x) = \frac{2}{3} \right\} = \left\{ x \in [0, 1] : x = \sum_{n=1}^{\infty} b_n 4^{-n}, b_n \in \{1, 2\} \right\}.$$

**T16) As in T11), study the self-similarity of  $\Phi_1, \Phi_2$  on  $\{(x, T(x)) : x \in [0, 1], T(x) = \frac{2}{3}\}$ , which is a cantor set of Hausdorff dimension  $\frac{1}{2}$ .**

Solution: Same as T11), denote  $\Gamma' = \{(x, T(x)) : x \in [0, 1], T(x) = \frac{2}{3}\}$ , then

$$\Phi_1(\Gamma') = \left\{ (x, T(x)) : x \in \left[0, \frac{1}{2}\right], T(x) = \frac{2}{3} \right\}, \Phi_2(\Gamma') = \left\{ (x, T(x)) : x \in \left[\frac{1}{2}, 1\right], T(x) = \frac{2}{3} \right\}.$$

## 7 Homework 7: Émile Borel Lemma

### 7.1 PSA

$f$  is a function on the interval  $I$ .

**A1) Suppose  $f$  is twice-differentiable at  $x$ , prove that**

$$f''(x) = \lim_{h \rightarrow 0} \frac{f(x+h) + f(x-h) - 2f(x)}{h^2}.$$

Proof: For any  $h > 0$ , consider the function  $g(t) = f(t) - f(t-h)$ , then there exists  $\xi \in [0, h]$  such that  $g(x+h) = g(x) + hg'(\xi)$ , and there exists  $\eta \in [\xi-h, \xi] \subset [-h, h]$  such that  $f'(\xi) - f'(\xi-h) = hf''(\eta)$

$$\frac{f(x+h) + f(x-h) - 2f(x)}{h^2} = \frac{f'(\xi) - f'(\xi-h)}{h} = f''(\eta),$$

therefore

$$f''(x) = \lim_{h \rightarrow 0} \frac{f(x+h) + f(x-h) - 2f(x)}{h^2}.$$

**A2) Suppose  $x_0 \in I$ , and**

$$\begin{aligned} f(x) &= a_0 + a_1(x-x_0) + \cdots + a_n(x-x_0)^n + o(|x-x_0|^n) \\ &= b_0 + b_1(x-x_0) + \cdots + b_n(x-x_0)^n + o(|x-x_0|^n). \end{aligned}$$

when  $x \rightarrow x_0$ , then for any  $i = 0, 1, \dots, n$ ,  $a_i = b_i$ .

Proof: Otherwise let  $c_i = a_i - b_i$  and take the least  $k$  such that  $c_k \neq 0$ , then

$$c_k(x-x_0)^k + \cdots + c_n(x-x_0)^n + o(|x-x_0|^n) = 0 \implies c_k = -c_{k+1}(x-x_0) - \cdots - c_n(x-x_0)^{n-k} + o(|x-x_0|^{n-k}),$$

which leads to contradiction when  $x \rightarrow x_0$ .

**A3) Suppose  $f$  is  $n$ -times differentiable at 0. Prove that if  $f$  is an even (odd) function, then the Taylor expansion of  $f$  at 0 has only even (odd) terms.**

Proof: Use the fact that if  $f$  is even (odd) then  $f'$  is odd (even).

**A4) If  $f$  is differentiable on  $(a, b)$  and  $\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow b^-} f(x)$  prove that exists  $x_0 \in (a, b)$  such that  $f'(x_0) = 0$ .**

Proof: Otherwise if  $f'(x) \neq 0$  for all  $x \in (a, b)$ , by Darboux's theorem  $f'(x)$  have the same sign over  $(a, b)$ , hence  $f$  is monotonic and non-constant on  $(a, b)$ , contradicting  $\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow b^-} f(x)$ .

**A5) Suppose  $f \in C([a, b])$  and is differentiable on  $(a, b)$ . Prove that  $f$  is strictly increasing on  $[a, b]$  iff for any  $x \in (a, b)$ ,  $f'(x) \geq 0$  and on any sub-interval  $(c, d) \subset (a, b)$ ,  $f'(x)$  does not vanish.**

Proof:  $\implies$  For any  $x \in (a, b)$ ,  $(f(x+h) - f(x))/h \geq 0$  so

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \geq 0.$$

If  $f'(x)$  vanish on some sub-interval  $(c, d)$  then  $f(c) = f(d)$ , a contradiction.

$\Leftarrow$  For any  $a \leq x < y \leq b$ , there exists  $\xi \in (a, b)$  such that  $f(y) - f(x) = (y - x)f'(\xi)$ , hence  $f(y) \geq f(x)$  and  $f$  is increasing. If  $f(x) = f(y)$  for some  $x < y$  then  $f(t)$  is constant on  $[x, y]$  and hence  $f'$  vanish on  $(x, y)$ , a contradiction.

## 7.2 PSB

Use L'Hôpital theorem to calculate limits:

**B1)  $a > 0$ , then**

$$\lim_{x \rightarrow \infty} \frac{\log x}{x^a} = \lim_{x \rightarrow \infty} \frac{x^{-1}}{ax^{a-1}} = 0.$$

**B2)  $a > 0, b > 1$  then**

$$\lim_{x \rightarrow \infty} \frac{x^a}{b^x} = \lim_{x \rightarrow \infty} \frac{ax^{a-1}}{b^x \ln b} = \dots = \lim_{x \rightarrow \infty} \frac{a(a-1) \cdots \{a\}}{b^x (\ln b)^{[a]} x^{1-\{a\}}} = 0.$$

**B3)**

$$\lim_{x \rightarrow 0} \frac{e^{ax} - e^{bx}}{\sin ax - \sin bx} = \lim_{x \rightarrow 0} \frac{ae^{ax} - be^{bx}}{a \cos ax - b \cos bx} = 1.$$

**B4)**

$$\lim_{x \rightarrow 0} \frac{\tan x - x}{x - \sin x} = \lim_{x \rightarrow 0} \frac{\sec^2 x - 1}{1 - \cos x} = \lim_{x \rightarrow 0} \frac{1 + \cos x}{\cos^2 x} = 2.$$



B5)

$$\lim_{x \rightarrow 0} \frac{1 - \cos x^2}{x^2 \sin x^2} = \lim_{x \rightarrow 0} \frac{1 - \cos x}{x \sin x} = \lim_{x \rightarrow 0} \frac{\sin x}{\sin x + x \cos x} = \lim_{x \rightarrow 0} \frac{\cos x}{2 \cos x - x \sin x} = \frac{1}{2}.$$

B6)

$$\lim_{x \rightarrow 1} \frac{\sqrt{2x - x^4} - \sqrt[3]{x}}{1 - x^{4/3}} = \lim_{x \rightarrow 1} \frac{(2x - x^4)^{-1/2}(1 - 2x^3) - x^{-2/3}/3}{-\frac{4}{3}x^{1/3}} = 1.$$

B7)

$$\lim_{x \rightarrow 1^-} (\log x)(\log(1 - x)) = \lim_{x \rightarrow 1^-} \frac{\log(1 - x)}{1/\log x} = \lim_{x \rightarrow 1^-} \frac{x \log^2 x}{1 - x} = 0.$$

B8)

$$\lim_{x \rightarrow 0^+} \frac{\log \sin ax}{\log \sin bx} = \lim_{x \rightarrow 0^+} \frac{\sin bx}{\sin ax} \cdot \frac{a \cos ax}{b \cos bx} = 1.$$

B9)

$$\lim_{x \rightarrow 0^+} x^x = \exp \lim_{x \rightarrow 0^+} \frac{\log x}{x^{-1}} = \exp \lim_{x \rightarrow 0^+} -x = 1.$$

B10)

$$\lim_{x \rightarrow 1} x^{1/(1-x)} = \exp \lim_{x \rightarrow 1} \frac{\log x}{1 - x} = e^{-1}.$$

B11)

$$\begin{aligned} \lim_{x \rightarrow 1} \left( \frac{1}{\log x} - \frac{1}{x - 1} \right) &= \lim_{x \rightarrow 1} \frac{x - 1 - \log x}{(x - 1) \log x} = \lim_{x \rightarrow 1} \frac{1 - x^{-1}}{1 - x^{-1} + \log x} \\ &= \lim_{x \rightarrow 1} \frac{x - 1}{x - 1 + x \log x} = \frac{1}{2}. \end{aligned}$$

B12)

$$\lim_{x \rightarrow 0^+} (\sin x)^x = \exp \lim_{x \rightarrow 0^+} \frac{\log \sin x}{x^{-1}} = \exp \lim_{x \rightarrow 0^+} -\frac{x^2}{\tan x} = 1.$$

**B13)**

$$\begin{aligned}\lim_{x \rightarrow 0} \left( \frac{\sin x}{x} \right)^{1/(1-\cos x)} &= \exp \lim_{x \rightarrow 0} \frac{\log \sin x - \log x}{1 - \cos x} = \exp \lim_{x \rightarrow 0} \frac{\cot x - x^{-1}}{\sin x} \\ &= \exp \lim_{x \rightarrow 0} \frac{x \cos x - \sin x}{x \sin^2 x} = \exp \lim_{x \rightarrow 0} \frac{-x \sin x}{\sin^2 x + x \sin 2x} \\ &= e^{-1/3}.\end{aligned}$$

**B14)**

$$\lim_{x \rightarrow a} \frac{a^x - x^a}{x - a} = \lim_{x \rightarrow a} \frac{a^x \log a - a x^{a-1}}{1} = a^a (\log a - 1).$$

**B15)**

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{(1 + 1/x)^x - e}{1/x} &= \lim_{x \rightarrow 0} \frac{(1 + x)^{1/x} - e}{x} = \lim_{x \rightarrow 0} (1 + x)^{1/x} \cdot \frac{x/(x+1) - \log(1+x)}{x^2} \\ &= e \lim_{x \rightarrow 0} \frac{(x+1)^{-2} - (x+1)^{-1}}{2x} = \frac{e}{2}.\end{aligned}$$

**B16)**

$$\lim_{x \rightarrow \infty} \frac{x^{\log x}}{(\log x)^x} = \exp \lim_{x \rightarrow \infty} (\log x)^2 - x \log \log x = 0.$$

**B17)**

$$\begin{aligned}\lim_{x \rightarrow \infty} (x+a)^{1+1/x} - x^{1+1/(x+a)} &= \lim_{x \rightarrow \infty} \frac{(x+a)^{1+1/x} x^{-1} - x^{1/(x+a)}}{x^{-1}} \\ &= a.\end{aligned}$$

**B18)**

$$\lim_{x \rightarrow \infty} \sqrt[3]{x^3 + x^2 + x + 1} - \sqrt{x^2 + x + 1} \cdot \frac{\log(e^x + x)}{x} = -\frac{1}{6}.$$

(Using WolframAlpha)

### 7.3 PSC

Calculate the maximum and minimum values of the following functions:

1.  $f(x) = x^4 - 2x^2 + 5, x \in [-2, 2]$ .

$f(x) = (x^2 - 1)^2 + 4 \in [4, 13]$ .

2.  $f(x) = \frac{2x}{1+x^2}, x \in \mathbb{R}$

$1 - f(x) = (1 + x^2)^{-1}(x - 1)^2 \geq 0, f(x) + 1 = (1 + x^2)^{-1}(x + 1)^2 \geq 0$ , therefore  $f(x) \in [-1, 1]$ .

3.  $f(x) = \arctan x - \frac{1}{2} \log(1 + x^2), x \in \mathbb{R}$ .

$f'(x) = \frac{1-x}{x^2+1}$ , hence  $\sup_{x \in \mathbb{R}} f(x) = f(1) = \frac{\pi}{4} - \frac{\log 2}{2}$ , and  $f$  has no minimum.

4.  $f(x) = x \log x, x \in (0, \infty)$ .

$f'(x) = \log x + 1$ , hence  $\inf_{x \in (0, \infty)} f(x) = f(e^{-1}) = -e^{-1}$ , and  $f$  has no maximum.

5.  $f(x) = \sqrt{x} \log x, x \in (0, \infty)$ .

$f'(x) = x^{-1/2} \left(1 + \frac{\log x}{2}\right)$ , hence  $\inf_{x \in (0, \infty)} f(x) = f(e^2) = -2e^{-1}$ .

6.  $f(x) = 2 \tan x - \tan^2 x, x \in [0, \pi/2]$ .

$f(x) = 1 - (1 - \tan x)^2 \in (-\infty, 1]$ .

### 7.4 PSD

$f$  is differentiable on  $(a, b)$ . Suppose  $x_0 \in (a, b)$  and  $f'(x_0) = 0$ .

**D1) Prove that  $f(x_0)$  is a local maximum if there exists  $(x_0 - \delta, x_0 + \delta) \subset (a, b)$  such that**

$$f'(x) = \begin{cases} > 0, & \forall x \in (x_0 - \delta, x_0), \\ < 0, & \forall x \in (x_0, x_0 + \delta). \end{cases}$$

Proof: Trivial by Lagrange mean-value theorem.

**D2) Prove that if  $f''(x_0)$  exists and  $f''(x_0) < 0$  then  $f(x_0)$  is a local maximum.**

Proof:  $f''(x_0) < 0$  and  $f'(x_0) = 0$  implies for some  $\delta > 0$ ,  $f'(x) < 0$  for  $x \in (x_0, x_0 + \delta)$  and  $f'(x) > 0$  for  $x \in (x_0 - \delta, x_0)$ . Hence by D1),  $f(x_0)$  is a local maximum.

**D3) Suppose  $f$  is  $n$ -times differentiable at  $x_0$ ,  $f'(x_0) = \dots = f^{(n-1)}(x_0) = 0$  and  $f^{(n)}(x_0) \neq 0$ . Determine the conditions that  $f(x_0)$  is a local maximum.**

Solution:  $n$  is even and  $f^{(n)}(x_0) < 0$ .

### 7.5 PSE: Roots of Polynomials

**E1) Prove that if all the roots of the polynomial  $P_n(x) \in \mathbb{R}[x]$  are real numbers, then so are the polynomials  $P'_n(x), P''_n(x), \dots, P_n^{n-1}(x)$ , where  $n = \deg P_n$ .**

Proof: We only need to prove that  $P'_n$  has  $n - 1$  real roots. By Rolle's mean-value theorem, between any two roots of  $P_n$  there is a root of  $P'_n$  hence  $P'_n$  has  $n - 1$  real roots.

**E2) Prove that the Legendre polynomial  $P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$  has  $n$  different roots in the interval  $(-1, 1)$ .**

Proof: We know that the polynomials  $\sqrt{(2n+1)/2} P_n(x)$  form a set of orthogonal base on the space  $L^2([-1, 1])$ , hence it must have  $n$  different roots in the interval  $(-1, 1)$ .

**E3) Prove that the Laguerre polynomial  $L_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n}(e^{-x}x^n)$  has  $n$  different real roots.**

Proof: We know that the Laguerre polynomials are orthogonal on the space  $L^2([0, \infty))$  with weight  $e^{-x}$ , hence it must have  $n$  distinct roots.

Or note that  $f(x) = x^n e^{-x}$  has a root with multiplicity  $n$  at 0 and it vanishes at  $\infty$ , hence use Rolle's theorem and induction we can show that  $f^{(k)}(x)$  has a root with multiplicity  $n - k$  at 0 and  $k$  roots between 0 and  $\infty$ .

**E4) Prove that the Hermite polynomial  $H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n}(e^{-x^2})$  has  $n$  different real roots.**

Proof: We know that the polynomials  $H_n(x)/\sqrt{2^n n! \sqrt{n}}$  form a set of orthogonal base on the Hilbert space  $L^2(\mu)$  where  $\mu(dx) = e^{-x^2} dx$ , hence it must have  $n$  distinct real roots.

## 7.6 PSF: Émile Borel's Lemma

**Part 1:**

**F1) Define  $\phi : \mathbb{R} \rightarrow \mathbb{R}$ :**

$$\phi(x) = \begin{cases} e^{-1/x^2}, & x > 0, \\ 0, & x \leq 0. \end{cases}$$

Prove that  $\phi \in C^\infty(\mathbb{R})$ .

Proof: We prove by induction that for any  $n \in \mathbb{Z}_{\geq 0}$ , there is a polynomial  $P_n \in \mathbb{R}[x]$  such that

$$\phi^{(n)}(x) = \begin{cases} P_n(1/x) \cdot e^{-1/x^2}, & x > 0, \\ 0, & x \leq 0. \end{cases}$$

(Which implies  $\phi^{(n)}$  is continuous.)

The case  $n = 0$  is trivial. Suppose it holds for  $n$ , then for any  $x > 0$ ,

$$\phi^{(n+1)}(x) = e^{-1/x^2} \left( P_n(1/x) \frac{2}{x^3} - P'_n(1/x) \frac{1}{x^2} \right),$$

for any  $x < 0$ ,  $\phi^{(n+1)}(x) = 0$ , and for  $x = 0$ ,

$$\phi_+^{(n+1)}(0) = \lim_{x \rightarrow 0^+} e^{-1/x^2} P_n(1/x) \frac{1}{x} = 0.$$

Hence the claim holds for  $n + 1$  too.

Therefore  $\phi \in C^\infty(\mathbb{R})$ .

**F2) Define  $\chi : \mathbb{R} \rightarrow \mathbb{R}$ :**

$$\chi(x) = \frac{\phi(2 - |x|)}{\phi(2 - |x|) + \phi(|x| - 1)}.$$

Prove that  $\chi(x) \in C^\infty(\mathbb{R})$  and  $\chi|_{[-1,1]} \equiv 1$ ,  $\chi|_{(-\infty, -2] \cup [2, \infty)} \equiv 0$ ,  $0 \leq \chi(x) \leq 1$  and  $\chi$  is an even function.

Proof:  $2 - |x|$  and  $|x| - 1$  cannot be both negative, hence the denominator is always positive, so  $\chi \in C^\infty(\mathbb{R})$ . The fact that  $\chi|_{[-1,1]} \equiv 1$ ,  $\chi|_{(-\infty, -2] \cup [2, \infty)} \equiv 0$ ,  $\chi(x) \in [0, 1]$  and  $\chi$  is even is trivial.

**F3) Prove that for any  $0 < a < b$ , there exists a smooth function  $\rho(x) \in C^\infty(\mathbb{R})$  such that  $\rho|_{[-a,a]} \equiv 1$ ,  $\rho|_{(-\infty,-b] \cup [b,\infty)} \equiv 0$ , and  $0 \leq \rho(x) \leq 1$ .**

Proof: Same as F2), define

$$\rho(x) = \frac{\phi(b - |x|)}{\phi(b - |x|) + \phi(|x| - a)}.$$

**F4) Prove that there exists an even function  $\psi \in C^\infty(\mathbb{R}^n)$  such that  $\psi|_{\{x:|x| \leq 1\}} \equiv 1$ ,  $\psi|_{\{x:|x| \geq 2\}} \equiv 0$ , and  $0 \leq \psi(x) \leq 1$ .**

Proof: (A special case of Urysohn's lemma)

Define  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  as  $f(\mathbf{x}) = \phi(1 - |\mathbf{x}|^2)$  and  $g: \mathbb{R}^n \rightarrow \mathbb{R}$  as  $g(\mathbf{x}) = \phi(|x^2|/4 - 1)$ , then  $f$  vanishes on  $B(0,1)^C$  and  $g$  vanishes on  $\bar{B}(0,2)$ . Therefore

$$\psi(\mathbf{x}) = \frac{f(\mathbf{x})}{f(\mathbf{x}) + g(\mathbf{x})}$$

satisfy the requirements.

**Part 2: Interchanging  $\sum$  and  $\frac{d}{dx}$**

$I = [a, b]$  is a closed interval,  $\{f_k\}_{k \geq 0}$  is a sequence of functions in  $C^1(I)$ . Assume  $\sum_{k=0}^\infty f_k$  converges point-wise on  $I$ , and let  $f(x) = \sum_{k=0}^\infty f_k(x)$ .

**F5) Assume the series  $\sum_{k=0}^\infty f'_k(x)$  converges absolutely on  $I$ , i.e.  $\sum_{k=0}^\infty \|f'_k\|_\infty$  converges. Prove that  $f$  is differentiable and  $f'(x) = \sum_{k=0}^\infty f'_k(x)$ .**

Proof: Note that

$$\frac{f(x+h) - f(x)}{h} = \sum_{k=0}^\infty \frac{f_k(x+h) - f_k(x)}{h} = \sum_{k=0}^\infty f'_k(x + \xi_k).$$

Hence

$$\left| \frac{f(x+h) - f(x)}{h} - \sum_{k=0}^\infty f'_k(x) \right| \leq \sum_{n=0}^N |f'_k(x + \xi_n) - f'_k(x)| + 2 \sum_{n=N+1}^\infty \|f'_k\|$$

Note that  $f'_k$  is uniformly continuous, so

$$\lim_{h \rightarrow 0} \sum_{n=0}^N |f'_k(x + \xi_n) - f'_k(x)| = 0, \quad \lim_{N \rightarrow \infty} 2 \sum_{n=N+1}^\infty \|f'_k\| = 0.$$

Hence

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \sum_{k=0}^\infty f'_k(x).$$

**F6) Assume  $\sum_{k=0}^\infty f'_k(x)$  converges uniformly on  $I$ , then  $f$  is differentiable and  $f'(x) = \sum_{k=0}^\infty f'_k(x)$ .**

Proof: Let  $g(x) = \sum_{k=0}^\infty f'_k(x)$ , since the series converges uniformly,  $g(x)$  is continuous on  $I$ . By Lebesgue's Dominated Convergence Theorem,

$$\int_{x_0}^x g(t) dt = \sum_{k=0}^\infty f_k(t) \Big|_{x_0}^x = f(x) - f(x_0).$$

Hence  $f'(x) = g(x) = \sum_{k=0}^\infty f'_k(x)$ .

**F7) Calculate the derivative of  $e^x$  using F6).**

Solution: On any closed interval  $[-M, M]$ ,

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = e^x$$

converges uniformly. Hence

$$(e^x)' = \sum_{n=0}^{\infty} \left( \frac{x^n}{n!} \right)' = \sum_{n=0}^{\infty} \frac{x^n}{n!} = e^x.$$

### Part 3: Borel's Lemma

Given an arbitrary sequence  $\{a_k\}_{k \geq 0}$ .

**F8) For any  $t_k > 0$ , let  $f_k(x) = \frac{a_k}{k!} x^k \chi(t_k x)$ , determine the derivatives of any order of  $f_k$  at  $x = 0$ .**

Solution: Note that when  $x = 0$ ,  $\chi^{(m)}(t_k x) = 0$  for any  $m \geq 1$  and  $\chi(t_k x) = 1$ . Hence

$$f_k^{(n)}(0) = \frac{a_k}{k!} \sum_{j=0}^n \binom{n}{j} (x^k)^{(j)} \chi^{(n-j)}(t_k x) \Big|_{x=0} = \frac{a_k}{k!} (x^k)^{(n)} \Big|_{x=0} = a_k \delta_{n,k}.$$

**F9) Prove that when  $k \geq 2n$ ,**

$$f_k^{(n)}(x) = a_k \sum_{=0}^n \binom{n}{=0} \frac{t_k^{n-}}{(k-)!} x^{k-} \chi^{(n-)}(t_k x).$$

Proof: Leibniz's Formula.

**F10) (Borel's lemma) Prove that for any sequence  $\{a_k\}_{k \geq 0}$ , there exists a smooth function  $f$  on  $\mathbb{R}$ , such that for any  $k \geq 0$ ,  $f^{(k)}(0) = a_k$ .**

Proof: Let  $f_k(x) = \frac{a_k}{k!} x^k \chi(t_k x)$  where  $t_k$  is yet to be determined, and

$$f(x) = \sum_{k=0}^{\infty} f_k(x) = \sum_{k=0}^{\infty} \frac{a_k x^k}{k!} \chi(t_k x).$$

For any  $n \geq 0$ , we want to show that  $\sum_{k=0}^{\infty} f_k^{(n)}(x)$  converges uniformly on  $\mathbb{R}$ . Suppose  $M_n = \sup_{x \in \mathbb{R}, m \leq n} |\chi^{(m)}(x)|$ , and

$$C_k = \sup_{n < k/2} \sum_{=0}^n \frac{\binom{n}{=0}}{(k-)!},$$

then for any  $x \in \mathbb{R}$ ,

$$|f_k^{(n)}(x)| \leq |a_k| C_k M_k t_k^{-k/2}.$$

Hence if we choose  $t_k$  such that

$$|a_k| C_k M_k t_k^{-k/2} < 2^{-k},$$

then the series

$$\sum_{k=0}^{\infty} f_k^{(n)}(x)$$

converges uniformly on  $\mathbb{R}$ . By F6) we know that  $f(x) = \sum_{k=0}^{\infty} f_k(x)$  is smooth, and by F8) we obtain  $f^{(n)}(0) = a_n$  for any  $n \geq 0$ ,

#### Part 4: Peano's Proof

**F11)  $\{c_k\}$  and  $\{b_k\}$  are two sequences, and  $b_k > 0$ . Prove that**

$$\left( \frac{c_k x^k}{1 + b_k x^2} \right)^{(n)}(0) = \begin{cases} n!(-1)^j c_{n-2j} b_{n-2j}^j, & \text{if } k = n - 2j, j \in \mathbb{Z}_{\geq 0}; \\ 0, & \text{otherwise.} \end{cases}$$

Proof: For  $x \rightarrow 0$ ,

$$\frac{c_k x^k}{1 + b_k x^2} = c_k \sum_{n=0}^{\infty} (-1)^n x^{2n+k} b_k^n.$$

Which converges absolutely on the interval  $[-b_k^{-1/2}/2, b_k^{-1/2}/2]$ , and so are its  $n$ -times derivations, hence by F5)

$$\left( \frac{c_k x^k}{1 + b_k x^2} \right)^{(n)}(0) = c_k \sum_{j=0}^{\infty} (-1)^j \frac{(2j+k)!}{(2j+k-n)!} x^{2j+k-n} b_k^j \Big|_{x=0} = \begin{cases} n!(-1)^j c_k b_k^j, & k = n - 2j, \\ 0, & \text{otherwise.} \end{cases}$$

**F12) Prove that there is a constant  $C$  such that for any  $k \geq n + 2$ , and any  $x$ ,**

$$\left| \left( \frac{c_k x^k}{1 + b_k x^2} \right)^{(n)}(x) \right| \leq C(n+1)! \frac{|c_k| k!}{b_k} |x|^{k-n-2}.$$

Proof: Use du Bois-Reymond, we can let  $C = 1$ .

**F13) Prove that for a given  $\{c_k\}$ , we can choose  $\{b_k\}$  such that  $b_k$  depends only on the value of  $c_k$ , and the function**

$$f(x) = \sum_{k=0}^{\infty} \frac{c_k x^k}{1 + b_k x^2}$$

is infinitely differentiable.

Proof: Let  $b_k = (k!)^2 c_k$ , then by F12),

$$\left| \sum_{k \geq n+2} \left( \frac{c_k x^k}{1 + b_k x^2} \right)^{(n)} \right| \leq (n+1)! \sum_{k \geq n+2} \frac{|x|^{k-n-2}}{k!}$$

hence the series

$$\sum_{k=0}^{\infty} \left( \frac{c_k x^k}{1 + b_k x^2} \right)^{(n)}$$

converges uniformly for any  $n \geq 1$ . By F6) the function  $f(x)$  is infinitely differentiable, and

$$f^{(n)}(x) = \sum_{k=0}^{\infty} \left( \frac{c_k x^k}{1 + b_k x^2} \right)^{(n)}.$$

**F14) Prove that  $f(0) = c_0, f'(0) = c_1$  and when  $n \geq 2$ ,**

$$\frac{f^{(n)}(0)}{n!} = c_n + \sum_{j=1}^{\lfloor n/2 \rfloor} (-1)^j c_{n-2j} b_{n-2j}^j.$$

Proof: Combine F11) and F13).

**F15) Prove that by carefully choosing  $\{c_k\}$  and  $\{b_k\}$ , we can prove Borel's lemma.**

Proof: Let  $b_k = (k!)^2 c_k$  and define  $c_k$  inductively such that

$$c_n = \frac{a_n}{n!} + \sum_{j=1}^{\lfloor n/2 \rfloor} (-1)^j c_{n-2j} b_{n-2j}^j$$

Then let  $f(x) = \sum_{k=0}^{\infty} \frac{c_k x^k}{1+b_k x^2}$

## 7.7 PSG: Midterm Test Part B

Consider  $f : \mathbb{R} \rightarrow \mathbb{R}$ .

- Let  $\mathcal{B}$  be all bounded function on  $\mathbb{R}$ .
- Let  $\mathcal{L}$  be all Lipschitz functions on  $\mathbb{R}$ .

Suppose  $a, \lambda \in \mathbb{R}$ ,  $f \in \mathcal{B} \cap \mathcal{L}$ , the goal is to find a function  $F \in \mathcal{L}$  to solve:

$$F(x) - \lambda F(x+a) = f(x), x \in \mathbb{R}. \quad (\star)$$

### Part 1: Basic Properties of Lipschitz Functions

**B1) Prove that if  $f, g \in \mathcal{B} \cap \mathcal{L}$ , then  $fg \in \mathcal{L}$ .**

Proof: Suppose  $|f(x) - f(y)|, |g(x) - g(y)| \leq A|x - y|$ , and  $|f(x)|, |g(x)| \leq C$ , then for any  $x, y \in \mathbb{R}$ ,

$$|f(x)g(x) - f(y)g(y)| \leq 2MA|x - y|.$$

Hence  $fg \in \mathcal{L}$ .

**B2) Prove that if  $f$  is differentiable and  $f \in \mathcal{L}$  then  $f' \in \mathcal{B}$ .**

Proof: If  $|f(x) - f(y)| \leq C|x - y|$  then for any  $x \in \mathbb{R}$ ,

$$|f'(x)| = \lim_{h \rightarrow 0} \left| \frac{f(x+h) - f(x)}{h} \right| \leq C.$$

Hence  $f' \in \mathcal{B}$ .

**B3) Prove that if  $f$  is differentiable and  $f' \in \mathcal{B}$  then  $f \in \mathcal{L}$ .**

Proof: For any  $x, y \in \mathbb{R}$ , there exists  $\xi \in (x, y)$  such that

$$|f(x) - f(y)| = |x - y| \cdot |f'(\xi)| \leq \sup_{t \in \mathbb{R}} |f'(t)| \cdot |x - y|.$$

Hence  $f \in \mathcal{L}$ .



**B4) If  $f \in \mathcal{B}$  and there exists  $B > 0$  such that for any  $x, y \in \mathbb{R}$ ,  $|x - y| \leq 1$  implies  $|f(x) - f(y)| \leq B|x - y|$ . Prove that  $f \in \mathcal{L}$ .**

Proof: Suppose  $M = \sup_{x \in \mathbb{R}} |f(x)|$ , then for any  $x, y \in \mathbb{R}$ ,

$$|f(x) - f(y)| \leq \max\{B, 2M\}|x - y|.$$

Hence  $f \in \mathcal{L}$ .

**Part 2: Solution of  $(\star)$  when  $|\lambda| < 1$ .**

Suppose  $f \in \mathcal{B} \cap \mathcal{L}$  and  $|\lambda| < 1$ .

**B5) Suppose  $F$  satisfy  $(\star)$ . Prove that for any  $x \in \mathbb{R}$  and  $n \in \mathbb{Z}_{\geq 1}$ ,**

$$\begin{aligned} F(x) &= \lambda^n F(x + na) + \sum_{k=0}^{n-1} \lambda^k f(x + ka), \\ F(x) &= \lambda^{-n} F(x - na) - \sum_{k=1}^n \lambda^{-k} f(x - ka). \end{aligned}$$

Proof: Use induction and apply  $(\star)$ .

(Let  $n \rightarrow \infty$  and we can obtain  $F$  formally.)

**B6) Prove that for any  $x \in \mathbb{R}$ ,  $\sum_{k \geq 0} \lambda^k f(x + ka)$  converges.**

Proof: Since  $f$  is bounded,

$$\left| \sum_{k=n}^{n+p} \lambda^k f(x + ka) \right| \leq \frac{M\lambda^n}{1 - \lambda}.$$

Hence the series converges.

**B7-8) Let  $F(x) = \sum_{k \geq 0} \lambda^k f(x + ka)$ . Prove that  $F \in \mathcal{L}$  and solve  $(\star)$ .**

Proof: For any  $x, y \in \mathbb{R}$ ,

$$|F(x) - F(y)| \leq \sum_{k=0}^{\infty} \lambda^k |f(x + ka) - f(y + ka)| \leq \sum_{k=0}^{\infty} \lambda^k C|x - y| = \frac{C}{1 - \lambda}|x - y|.$$

Hence  $F \in \mathcal{L}$ . For any  $x \in \mathbb{R}$ ,

$$F(x) - \lambda F(x + a) = \sum_{k \geq 0} \lambda^k f(x + ka) - \sum_{k \geq 1} \lambda^k f(x + ka) = f(x).$$

Therefore  $F$  solves  $(\star)$ .

If  $F'$  also solves  $(\star)$ , let  $G = F - F'$ , then  $G$  is bounded and

$$G(x) = \lambda G(x + a), \quad x \in \mathbb{R}.$$

Therefore for any  $x \in \mathbb{R}$ ,

$$|G(x)| = \lambda^n |G(x + na)| \leq M\lambda^n \rightarrow 0.$$

Hence  $G \equiv 0$  and  $F \equiv F'$ , so the solution to  $(\star)$  is  $F$ .

**B9) Solve  $(\star)$  when  $f(x) \equiv 1$  and  $f(x) = \cos x$ .**

Solution: When  $f(x) \equiv 1$ ,

$$F(x) = \sum_{k=0}^{\infty} \lambda^k f(x + ka) = \frac{1}{1 - \lambda}.$$

When  $f(x) = \cos x$ ,

$$\begin{aligned} F(x) &= \sum_{k=0}^{\infty} \lambda^k \cos(x + ka) = \sum_{k=0}^{\infty} \lambda^k \frac{e^{i(x+ka)} + e^{-i(x+ka)}}{2} = \frac{1}{2} \left( \frac{e^{ix}}{1 - \lambda e^{ia}} + \frac{e^{-ix}}{1 - \lambda e^{-ia}} \right) \\ &= \frac{\cos x - \lambda \cos(x - a)}{1 - 2\lambda \cos a + \lambda^2}. \end{aligned}$$

**Part 3: Solution of  $(\star)$  when  $|\lambda| > 1$ .**

**B10) Solve  $(\star)$  as in Part 2.**

Solution: By B5), the solution should be

$$F(x) = - \sum_{k=1}^{\infty} \lambda^{-k} f(x - ka).$$

$f \in \mathcal{B}$  implies the series converges. Same as B8) we can show that the solution to  $(\star)$  is unique, and like B7) we can show that  $F \in \mathcal{L}$  and  $F$  satisfy  $(\star)$ .

**B11) Solve  $(\star)$  for  $f(x) \equiv 1$  and  $f(x) = \cos x$ .**

Solution: When  $f(x) \equiv 1$ ,

$$F(x) = - \sum_{k=1}^{\infty} \lambda^{-k} f(x - ka) = \frac{1}{1 - \lambda}.$$

When  $f(x) = \cos x$ , same as B9) we have

$$F(x) = - \sum_{k=1}^{\infty} \lambda^{-k} f(x - ka) = \frac{\cos x - \lambda \cos(x - a)}{1 - 2\lambda \cos a + \lambda^2}.$$

**Part 4: The Case when  $|\lambda| = 1$ .**

**B12) Suppose  $\lambda = 1$ . Prove that there exists  $F \in \mathcal{L}$  not identically zero, such that for any  $x$ ,  $F(x) - F(x + a) = 0$ .**

Proof: Let  $F(x) = |\{x/a\} - 1/2|$ , then  $F(x) = F(x + a)$ , and  $F \in \mathcal{L} \cap \mathcal{B}$ .

**B13) Let  $f(x) = \cos x$  in  $(\star)$ . Prove that if  $\cos a \neq 1$ , then there exists  $F \in \mathcal{L}$  that solves  $(\star)$ . Determine whether the solution is unique.**

Proof: The equation  $(\star)$  becomes  $F(x) = F(x + a) + \cos x$ . Let

$$F(x) = \{x/a\} - \sum_{k=0}^{\lfloor x/a \rfloor - 1} \cos(k + \{x/a\})a,$$

(if  $x < 0$  the sum is viewed as from  $\lfloor x/a \rfloor - 1$  to 0) then clearly  $F(x) = F(x+a) + \cos x$ , and  $F$  is bounded since  $\cos a \neq 1$ .

For any  $x, y \in \mathbb{R}$ , if  $|x - y| < a/2$ , then suppose  $na \leq x < y < (n+1)a$ ,

$$\begin{aligned} |F(x) - F(y)| &\leq \left| \left\{ \frac{x}{a} \right\} - \left\{ \frac{y}{a} \right\} \right| + 2 \left| \sin \frac{\{x/a\} - \{y/a\}}{2} a \right| \cdot \left| \sum_{k=0}^{n-1} \sin(k + (\{x/a\} + \{y/a\})/2)a \right| \\ &\leq \frac{|x - y|}{a} + \frac{|x - y|}{|\sin a|}. \end{aligned}$$

Hence  $F \in \mathcal{L}$  by B4), so  $F$  solves  $(\star)$ .

The solution is clearly not unique since we can add any factor of the  $F$  in B12) to the solution.

**B14) Following B13), if  $a = 2\pi$ , then  $(\star)$  has no solution in  $\mathcal{L}$ .**

Proof: If  $a = 2\pi$  and  $F$  is a solution to  $(\star)$ , then for any  $x, y \in \mathbb{R}$ ,

$$|F(x + 2\pi n) - F(y + 2\pi n)| = n|\cos x - \cos y| \rightarrow \infty.$$

Hence  $F \notin \mathcal{L}$ .

**B15) Suppose  $\lambda = -1$ , Prove that there exists  $F \in \mathcal{L}$  not identically zero, such that for any  $x$ ,  $F(x) + F(x+a) = 0$ .**

Proof: Let  $F(x) = |2\{x/2a\} - 1| - 1/2$ , then  $F \in \mathcal{L}$  and  $F(x) + F(x+a) = 0$ .

**B16) Suppose  $\lambda = -1$ ,  $a = 1$ ,  $f \in \mathcal{L}$  is monotonically decreasing and  $\lim_{x \rightarrow \infty} f(x) = 0$ ,  $f$  is differentiable and  $f'$  is increasing. Prove that there exists  $F \in \mathcal{L}$  such that**

$$F(x) + F(x+1) = f(x), \quad x \in \mathbb{R}.$$

Further show that if we require  $F \in \mathcal{L}$  and  $\lim_{x \rightarrow \infty} F(x) = 0$ , then the solution is unique.

Proof: Since  $f$  is monotonically decreasing, for any  $x \in \mathbb{R}$ , the series

$$F(x) = \sum_{n=0}^{\infty} (-1)^n f(x+n)$$

converges.

For any  $x, y \in \mathbb{R}$ ,  $|x - y| < 1$ , there exists  $\xi_n \in (x+n, y+n)$  such that  $f(y+n) - f(x+n) = (y-x)f'(\xi_n)$ , hence (by B3)  $f'$  is bounded)

$$|F(x) - F(y)| = |y - x| \cdot \left| \sum_{n=0}^{\infty} (-1)^n f'(\xi_n) \right| \leq \sup_{t \in \mathbb{R}} |f'(t)| \cdot |y - x|.$$

so  $F \in \mathcal{L}$ . Clearly  $F(x) + F(x+1) = f(x)$ , so  $F$  solves  $(\star)$ , and  $0 < F(x) < f(x)$  so  $\lim_{x \rightarrow \infty} F(x) = 0$ . If  $F' \in \mathcal{L}$  also satisfy  $(\star)$  and  $\lim_{x \rightarrow \infty} F(x) = 0$ , let  $G = F - F'$ , then  $G(x) + G(x+1) = 0$  and  $\lim_{x \rightarrow \infty} G(x) = 0$ . Hence  $G(x) = \lim_{n \rightarrow \infty} (-1)^n G(x+n) = 0$  for any  $x \in \mathbb{R}$ , so  $G \equiv 0$ . Therefore  $F$  is the unique solution.

## 8 Homework 8; Strum-Liouville Theory

### 8.1 PSA: Convex functions

A1)

(1)  $f(x) = |x|$ ,  $I = \mathbb{R}$  is convex, since

$$|\lambda x + (1 - \lambda)y| \leq \lambda|x| + (1 - \lambda)|y|.$$

(2)  $f(x) = x^p$ ,  $p \in \mathbb{R}$ ,  $I = \mathbb{R}_{>0}$

$f''(x) = p(p-1)x^{p-2}$  so  $f$  is concave if  $p \in [0, 1]$  and convex if  $p \in (-\infty, 0] \cup [1, \infty)$ .

(3)  $f(x) = \sin x$ ,  $I = [0, \pi]$  is concave since  $f''(x) = -\sin x \leq 0$  when  $x \in [0, \pi]$ .

(4)  $f(x) = x \log x$ ,  $I = \mathbb{R}_{\geq 0}$  is (strictly) convex since  $f''(x) = 1/x > 0$ .

(5)  $f(x) = \mathbf{1}_{\{0,1\}}$ ,  $I = [0, 1]$  is convex since

$$f(\lambda x + (1 - \lambda)y) = 0 \leq \lambda f(x) + (1 - \lambda)f(y).$$

A2) Prove the following properties:

1. If  $f, g$  are convex on  $I$ , then  $f + g$  is convex on  $I$ .

Proof: By definition,  $(f + g)(\lambda x + (1 - \lambda)y) \leq \lambda(f + g)(x) + (1 - \lambda)(f + g)(y)$ , so  $f + g$  is convex.

2. If  $f, g$  are monotonically increasing, non-negative, convex functions on  $I$ , then  $fg$  is convex.

Proof: Note that

$$f(\lambda x + (1 - \lambda)y)g(\lambda x + (1 - \lambda)y) \leq (\lambda f(x) + (1 - \lambda)f(y)) \cdot (\lambda g(x) + (1 - \lambda)g(y))$$

and

$$\begin{aligned} & \lambda f(x)g(x) + (1 - \lambda)f(y)g(y) - (\lambda f(x) + (1 - \lambda)f(y))(\lambda g(x) + (1 - \lambda)g(y)) \\ &= \lambda(1 - \lambda)(f(x) - f(y))(g(x) - g(y)) \geq 0. \end{aligned}$$

hence

$$(fg)(\lambda x + (1 - \lambda)y) \leq \lambda(fg)(x) + (1 - \lambda)(fg)(y).$$

1. If  $f$  is convex on  $I$ ,  $g$  is a monotonically increasing convex function on  $J \supset f(I)$ , then  $g \circ f$  is convex.

Proof: Note that

$$g(f(\lambda x + (1 - \lambda)y)) \leq g(\lambda f(x) + (1 - \lambda)f(y)) \leq \lambda g(f(x)) + (1 - \lambda)g(f(y)).$$

hence  $g \circ f$  is convex.

1. If  $f, g$  are convex on  $I$ , then  $h(x) = \max\{f(x), g(x)\}$  is convex.

Proof: For any  $x, y, \lambda$  and  $t = \lambda x + (1 - \lambda)y$ , suppose  $h(t) = f(t)$ , then

$$h(t) \leq \lambda f(x) + (1 - \lambda)f(y) \leq \lambda h(x) + (1 - \lambda)h(y)$$

hence  $h$  is convex.

**A3) Suppose  $f \in C((a, b))$ . If for any  $x, y \in (a, b)$ ,  $f\left(\frac{x+y}{2}\right) \leq \frac{f(x)+f(y)}{2}$ , prove that  $f$  is convex.**

Proof: For any  $x, y \in (a, b)$  and  $\lambda \in [0, 1]$ , we need to prove that

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y).$$

Note that it holds for any dyadic number  $\lambda$ , since the cases  $\lambda = 0, 1, 1/2$  is trivial, and for  $\lambda = (2m+1)/2^t$ , let  $u = m/2^{t-1}, v = (m+1)/2^{t-1}$ , then

$$\begin{aligned} f(\lambda x + (1 - \lambda)y) &\leq \frac{f(ux + (1 - u)y) + f(vx + (1 - v)y)}{2} \\ &\leq \lambda f(x) + (1 - \lambda)f(y). \end{aligned}$$

Now since  $f \in C((a, b))$ , for any  $\lambda \in (0, 1)$  there is a sequence of dyadic numbers  $\lambda_n$  such that  $\lim_{n \rightarrow \infty} \lambda_n = \lambda$ , hence

$$\begin{aligned} f(\lambda x + (1 - \lambda)y) &= \lim_{n \rightarrow \infty} f(\lambda_n x + (1 - \lambda_n)y) \leq \lim_{n \rightarrow \infty} \lambda_n f(x) + (1 - \lambda_n)f(y) \\ &= \lambda f(x) + (1 - \lambda)f(y). \end{aligned}$$

**A4)  $f$  is a convex function on  $[a, b]$ . Prove that if there exists  $c \in (a, b)$  such that  $f(c) \geq \max\{f(a), f(b)\}$  then  $f$  is constant.**

Proof: For any  $t \in (a, b)$ , let  $\lambda = (t - a)/(b - a)$  then

$$f(t) = f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b) \leq \max\{f(a), f(b)\}.$$

By  $f(c) \geq \max\{f(a), f(b)\}$  we know that  $f(a) = f(b)$ . If for some  $t \in (a, b)$ ,  $f(t) \neq f(a)$ , suppose  $c \in (a, t)$ , then

$$f(c) = \lambda f(a) + (1 - \lambda)f(t) < f(a)$$

a contradiction. Hence  $f(t) = f(a)$  for all  $t \in [a, b]$ .

**A5)  $f$  is convex on  $\mathbb{R}$ . Prove that if  $f$  has an upper-bound, then  $f$  is constant.**

Proof: Otherwise suppose that  $f(a) < f(b)$ , where  $a < b$ . (If  $f(a) > f(b)$  let  $g(x) = f(-x)$ ). Let  $x_0 = a, x_1 = b, x_n = a + n(b - a)$ , then

$$f(x_{n+1}) - f(x_n) \geq f(x_n) - f(x_{n-1}) \geq f(b) - f(a),$$

hence  $f(x_n) \geq f(a) + n(f(b) - f(a)) \rightarrow \infty$ , leading to contradiction.

**A6)  $f$  is strictly convex on  $I$ . Suppose  $f(x_0)$  is a local minimum of  $f$ , prove that  $x_0$  is the unique global minimum point of  $f$ .**

Proof: Suppose there is another  $x_1 \neq x_0$  such that  $f(x_1) \leq f(x_0)$ , then let  $x_n = x_0 + n(x_1 - x_0)$ . Since  $f$  is strictly convex,  $f(x_n) < \max\{f(x_1), f(x_0)\} = f(x_0)$ , contradicting the fact that  $f(x_0)$  is a local minimum.

**A7)  $I$  is an open interval. Prove that  $f$  is convex on  $I$ , iff for any  $x_0 \in I$ , there exists  $a \in \mathbb{R}$ , such that for any  $x \in I$ ,  $f(x) \geq a(x - x_0) + f(x_0)$ .**

Proof: Suppose  $f$  is convex on  $I$ , then the any  $x_0 \in I$ , the function  $g(x) = \frac{f(x) - f(x_0)}{x - x_0}$  is monotonically increasing. Hence we can let  $a = \sup_{x < x_0} g(x) < \infty$ .

If for any  $x_0 \in I$ , and  $x \in I$ ,  $f(x) \geq g(x_0)(x - x_0) + f(x_0)$ , then for any  $x, y \in I$  and  $\lambda \in (0, 1)$ , let  $t = \lambda x + (1 - \lambda)y$ ,

$$\begin{aligned}\lambda f(x) + (1 - \lambda)f(y) &\geq \lambda(f(t) + (x - t)g(t)) + (1 - \lambda)(f(t) + (y - t)g(t)) \\ &= f(t) = f(\lambda x + (1 - \lambda)y).\end{aligned}$$

Hence  $f$  is convex.

## 8.2 PSB

**B1) Prove the following inequalities:**

(1)

$$x - \frac{x^2}{2} < \log(1 + x) < x, \quad x > 0.$$

Proof: If  $f(x) = \log(x + 1) - x$ , then  $f'(x) = \frac{1}{x+1} - 1 < 0$  hence  $f(x) < f(0) = 0$ . Let  $g(x) = \log(1 + x) - x + x^2/2$ , then  $g'(x) = \frac{1}{x+1} + (x + 1) - 2 \geq 0$ , hence  $g(x) > g(0) = 0$ .

(2)

$$(x^\alpha + y^\alpha)^{1/\alpha} > (x^\beta + y^\beta)^{1/\beta}, \quad x, y > 0, \beta > \alpha > 0.$$

Proof: Assume that  $x^\alpha + y^\alpha = 1$ , then  $0 < x, y < 1$ , so

$$x^\beta + y^\beta < x^\alpha + y^\alpha < 1 \implies (x^\beta + y^\beta)^{1/\beta} < (x^\alpha + y^\alpha)^{1/\alpha}.$$

(3)

$$x - \frac{x^3}{6} < \sin x < x, \quad x > 0.$$

Proof: Let  $f(x) = \sin x - x$ , then  $f'(x) = \cos x - 1 \leq 0$ , so  $f(x) < f(0) = 0$ . Let  $g(x) = \sin x - x + x^3/6$ , then  $g'(x) = \cos x - 1 + x^2/2$ ,  $g''(x) = x - \sin x > 0$ , so  $g'(x) > g'(0) = 0$  and  $g(x) > g(0) = 0$ .

(4)

$$\left(\frac{1+x}{2}\right)^p + \left(\frac{1-x}{2}\right)^p \leq \frac{1}{2}(1+x^p), \quad p \in [2, \infty), x \in [0, 1].$$

Proof: ???

**B2) Find all  $a > 0$  such that  $a^x \geq x^a$  for any  $x > 0$ .**

Solution:  $f(x) = x^{1/x}$  then  $f'(x) = x^{1/x} \frac{1 - \log x}{x^2}$  hence  $f$  has a unique minimum at  $e$ .

**B3) Prove that for any  $x_i, t_i, i = 1, 2, \dots, n$ ,  $\sum_{i=1}^n t_i = 1$ ,**

$$\left(\sum_{i=1}^n t_i x_i\right)^{\sum_{i=1}^n t_i x_i} \leq \prod_{i=1}^n x_i^{t_i x_i}.$$

Proof: Let  $f(x) = x \log x$ , then  $f''(x) = 1/x > 0$ , so  $f$  is convex. By Jensen's inequality,

$$\sum_{i=1}^n t_i f(x_i) \geq f\left(\sum_{i=1}^n t_i x_i\right)$$

hence

$$\left( \sum_{i=1}^n t_i x_i \right)^{\sum_{i=1}^n t_i x_i} \leq \prod_{i=1}^n x_i^{t_i x_i}.$$

and equality holds iff  $x_i = x_1$ .

**B4) Prove that for any  $a, b > 0$ ,  $1/p + 1/q = 1$ ,**

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}, \text{ if } p > 1; ab \geq \frac{a^p}{p} + \frac{b^q}{q}, \text{ if } p < 1.$$

Proof: The function  $-\log x$  is convex, so when  $p > 1$ ,  $q > 1$ , then

$$\log \left( \frac{a^p}{p} + \frac{b^q}{q} \right) \geq \frac{1}{p} \log a^p + \frac{1}{q} \log b^q$$

so  $ab \leq \frac{a^p}{p} + \frac{b^q}{q}$ .

When  $p < 1$ , then  $pq < 0$ , so likewise  $ab \geq \frac{a^p}{p} + \frac{b^q}{q}$ .

**B5) Prove that if  $x_i, y_i \geq 0, i = 1, 2, \dots, n$ ,  $1/p + 1/q = 1$ , then**

$$\sum_{i=1}^n x_i y_i \leq \left( \sum_{i=1}^n x_i^p \right)^{1/p} \left( \sum_{i=1}^n y_i^q \right)^{1/q}, \text{ if } p > 1;$$

and the inequality reverses when  $p < 1$ .

Proof: Assume that  $\sum_{i=1}^n x_i^p = \sum_{i=1}^n y_i^q = 1$ , then by B4), if  $p > 1$ ,

$$\sum_{i=1}^n x_i y_i \leq \sum_{i=1}^n \frac{x_i^p}{p} + \frac{y_i^q}{q} = \frac{1}{p} + \frac{1}{q} = 1.$$

The case  $p < 1$  is similar.

### 8.3 PSC

**C1) Suppose  $f \in C([0, 1])$ ,  $g$  is differentiable on  $[0, 1]$  and  $g(0) = 0$ . If there is a constant  $\lambda \neq 0$ , such that for any  $x \in [0, 1]$ ,  $|g(x)f(x) + \lambda g'(x)| \leq |g(x)|$ , prove that  $g(x) \equiv 0$ .**

Proof: Otherwise assume that  $\forall \varepsilon > 0 \exists t \in (0, \varepsilon)$ , such that  $g(t) \neq 0$ . Let  $C = (1 + \sup_{x \in [0, 1]} |f(x)|)/\lambda$ , then  $|g'(x)| \leq C|g(x)|, \forall x \in [0, 1]$ . For any  $t \in (0, 1)$ , there exists  $\xi \in [0, t]$  such that  $g(t) = g(0) + tg'(\xi)$ , hence

$$\frac{|g(t)|}{t} = |g'(\xi)| \leq C \sup_{\xi \in [0, t]} |g(\xi)|.$$

For any  $t > 0$  suppose  $|g(s)| = \sup_{\xi \in [0, t]} |g(\xi)| > 0$ , then  $|g(s)|/s \leq C|g(s)|$  hence  $t \geq s \geq \frac{1}{C}$ , a contradiction.

**C2)  $f$  is twice differentiable on  $(-1, 1)$ ,  $f(0) = f'(0) = 0$ . If for any  $x \in (-1, 1)$ ,  $|f''(x)| \leq |f(x)| + |f'(x)|$ , prove that  $f(x) \equiv 0$ .**

Proof: We prove that  $f''(x) \equiv 0$ . Otherwise suppose  $\forall \varepsilon > 0, \exists x \in [0, \varepsilon], f''(x) \neq 0$ . Note that

$$|f''(x)| \leq |f(x)| + |f'(x)| \leq \left( \frac{x^2}{2} + |x| \right) \sup_{y \in [0, x]} |f''(y)|.$$

Since  $f''(0) = 0$ , take  $x \in [0, 1/2]$  such that  $f''(x) \neq 0$ , and suppose  $|f''(t)| = \sup_{y \in [0, x]} |f''(y)|$ , then  $|f''(t)| \leq (t^2/2 + t)|f''(t)|$ , a contradiction.

**C3)**  $f$  is  $n$ -times differentiable on  $\mathbb{R}$ ,  $f(0) = f'(0) = \dots = f^{(n-1)}(0) = 0$ . If there exists  $C \in \mathbb{R}_{>0}$  and  $\varepsilon \in \mathbb{Z}_{\geq 0}$  such that for any  $x \in \mathbb{R}$ ,  $|f^{(n)}(x)| \leq C|f^{(\varepsilon)}(x)|$ . Prove that  $f(x) \equiv 0$ .

Proof: We can assume that  $f(x) \neq 0$ . Since  $f^{(k)}(x) = 0, \forall 0 \leq k < n$ , we have

$$|f^{(n)}(x)| \leq C|f(x)| \leq C \frac{x^n}{n!} \sup_{y \in [0, x]} |f^{(n)}(y)|.$$

Hence for any  $x \in [0, \varepsilon]$ ,  $\varepsilon = (n!/C)^{1/n}$ ,  $f^{(n)}(x) = 0$ , so  $f^{(k)}(x) = 0$  for all  $x \in [0, \varepsilon], 0 \leq k < n$ . Likewise we get  $f(x) \equiv 0$ .

**C4)**  $n \in \mathbb{Z}_{>0}$ , prove that the polynomial  $P(x) = \sum_{k=0}^{n+1} \binom{n+1}{k} (-1)^k (x-k)^n \equiv 0$ .

Proof: We know the identity

$$\sum_{k=0}^n \binom{n}{k} (-1)^k k^l = 0, \forall 0 \leq l \leq n-1.$$

Since  $\Delta^n x^l \equiv 0$

Likewise by considering  $f(t) = (x-t)^n$  we have  $P(x) \equiv 0$ .

(Or we can use C3)

**C5)**  $f \in C^\infty(\mathbb{R})$ . Assume there exists  $C > 0$  such that for any  $n \in \mathbb{Z}_{\geq 0}$  and  $x \in \mathbb{R}$ ,  $|f^{(n)}(x)| \leq C$ .

i. Prove that given an arbitrary  $x_0 \in \mathbb{R}$ ,

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x-x_0)^k, \forall x \in \mathbb{R}.$$

Proof: The Lagrange remainder

$$R_n(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} (x-x_0)^{n+1}$$

tends to zero as  $n \rightarrow \infty$ , hence the Taylor series

$$f(x) = \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x-x_0)^k + R_n(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x-x_0)^k.$$

ii.  $E \subset \mathbb{R}$  is an infinite bounded set. Prove that if  $f(E) = \{0\}$ , then  $f \equiv 0$ .

Proof: Suppose  $E \subset [-M, M]$ , then by Bolzano-Weierstrass theorem, there exists a sequence  $\{z_n\}_{n \geq 1} \subset E$  such that  $z = \lim_{n \rightarrow \infty} z_n$  exists. Since  $f \in C(\mathbb{R})$ ,  $f(z) = \lim_{n \rightarrow \infty} f(z_n) = 0$ , so

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(z)}{k!} (x-z)^k.$$

If  $f$  does not vanish on  $\mathbb{R}$ , then take the least  $m > 0$  such that  $f^{(m)}(z) \neq 0$ . When  $z_n \rightarrow z$ ,

$$0 = \frac{f^{(m)}(z)}{m!} + \sum_{k=m+1}^{\infty} \frac{f^{(k)}(z)}{k!} (x-z)^{k-m}$$

which leads to contradiction. Hence  $f$  vanishes on  $\mathbb{R}$ .



**C6)** Assume  $f \in C^2((0, 1))$ ,  $\lim_{x \rightarrow 1^-} f(x) = 0$ . If there exists  $C > 0$ , such that for any  $x \in (0, 1)$ ,  $(1 - x)^2 |f''(x)| \leq C$ . Prove that  $\lim_{x \rightarrow 1^-} (1 - x)f'(x) = 0$ .

Proof: For any  $0 < x < y < 1$ , there exists  $\xi \in (x, y)$  such that

$$f(y) = f(x) + (y - x)f'(x) + \frac{(y - x)^2}{2}f''(\xi).$$

For any  $\lambda > 0$ , let  $y = (\lambda + x)/(\lambda + 1) \in (x, 1)$ , then

$$|(y - x)f'(x)| \leq |f(y)| + |f(x)| + \frac{\lambda^2}{2}(1 - y)^2|f''(\xi)| \leq |f(y)| + |f(x)| + \frac{C\lambda^2}{2}.$$

Therefore

$$|(1 - x)f'(x)| \leq (|f(y)| + |f(x)|)\frac{\lambda + 1}{\lambda} + \frac{1}{2}\lambda(\lambda + 1)C$$

Hence for any  $\lambda > 0$ ,

$$\lim_{x \rightarrow 1^-} |(1 - x)f'(x)| \leq \frac{1}{2}\lambda(\lambda + 1)C \rightarrow 0,$$

so  $\lim_{x \rightarrow 1^-} (1 - x)f'(x) = 0$ .

## 8.4 PSD

Calculate  $\sup_{x \in I} f(x)$  and  $\inf_{x \in I} f(x)$ :

**D1)**  $f(x) = \frac{(\log x)^2}{x}$ ,  $I = \mathbb{R}_{>0}$

Solution: Let  $y = \log x \in \mathbb{R}$ , then  $f(x) = y^2 e^{-y}$ .

$$\frac{d}{dy} y^2 e^{-y} = y e^{-y} (2 - y).$$

Hence  $\sup_{x \in I} f(x) = f(e^2) = 4e^{-2}$ ,  $\inf_{x \in I} f(x) = \min\{f(0), f(\infty)\} = 0$ .

**D2)**  $f(x) = |x(x^2 - 1)|$ ,  $I = \mathbb{R}$

Solution:  $\sup = \infty$ ,  $\inf = 0$ .

**D3)**

$$f(x) = \frac{x(x^2 + 1)}{x^4 - x^2 + 1}, I = \mathbb{R}.$$

Solution: Note that

$$2(x^4 - x^2 + 1) - x(x^2 + 1) = (x^2 - 1)^2 + (x - 1)^2(x^2 + x + 1) \geq 0.$$

Therefore  $f(x) \leq 2$  where equality holds at  $x = 1$ . Since  $f(x) = f(-x)$ ,  $\sup = 2$ ,  $\inf = -2$ .

**D4)**

$$f(x) = x^{1/3}(1 - x)^{2/3}, I = (0, 1).$$

Solution: By AM-GM,  $f(x) \leq \frac{2^{2/3}}{3}$  where equality holds at  $x = 1/3$ . Hence  $\sup = \frac{2^{2/3}}{3}$ ,  $\inf = 0$ .

**D5)**

$$f(x) = \left(1 + x + \frac{x^2}{2!} + \cdots + \frac{x^n}{n!}\right) e^{-x}, \quad I = \mathbb{R}.$$

Solution:  $f'(x) = -e^{-x} \frac{x^n}{n!}$ , so if  $n$  is even,  $\sup = \infty, \inf = 0$ , and if  $n$  is odd,  $\sup = 1, \inf = -\infty$ .

**D6)**  $f(x) = \sin^{2m} x \cos^{2n} x, \quad I = \mathbb{R}.$ 

Solution: Let  $t = \sin^2 x \in [0, 1]$ , then  $f(x) = t^m(1-t)^n \in [0, n^m m^n / (n+m)^{n+m}]$ .

**8.5 PSE**

Compare the two functions (or real numbers).

**E1)**  $f(x) = e^x, g(x) = 1 + xe^x, x > 0.$ 

Solution: The case  $x \geq 1$  is trivial. If  $x \in (0, 1)$ , then

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \leq \sum_{n=0}^{\infty} x^n = \frac{1}{1-x},$$

hence  $f(x) \leq g(x)$ . Therefore  $f(x) \leq g(x)$  for all  $x > 0$ .

**E2)**  $f(x) = xe^{x/2}, g(x) = e^x - 1, x > 0.$ 

Solution: ( $x/2 \leq \sinh(x/2)$ ) Consider  $h(x) = e^{x/2} - e^{-x/2} - x$ , then  $h(0) = 0$  and

$$h'(x) = \frac{1}{2}(e^{x/2} + e^{-x/2} - 2) \geq 0.$$

Hence  $h(x) \geq 0$ , i.e.  $g(x) \geq f(x)$  for all  $x > 0$ .

**E3)**  $f(x) = \left(\frac{x+1}{2}\right)^{(x+1)}, g(x) = x^x, x > 0.$ 

Solution: Consider  $h(x) = x \log x - (x+1) \log \frac{x+1}{2}$ , then  $h(1) = 0$  and

$$h'(x) = \log \frac{2x}{x+1} \geq 0 \iff x \geq 1.$$

Hence  $f(x) \leq g(x)$  for all  $x > 0$ .

**E4)**  $2^{\sqrt{2}}$  and  $e$ .

Solution: Note that

$$\log 2 = \sum_{n=1}^{\infty} \frac{1}{2^n n} \leq \frac{2}{3} + \sum_{n=4}^{\infty} \frac{1}{2^n \cdot 4} = \frac{2}{3} + \frac{1}{32} < \frac{2}{3} + \frac{1}{30} = 0.7 < \frac{1}{\sqrt{2}},$$

hence  $2^{\sqrt{2}} < e$ .

**E5)**  $f(x) = \log(1 + \sqrt{1 + x^2})$ ,  $g(x) = 1/x + \log x$ ,  $x > 0$ .

Solution: Consider  $h(x) = \log x + 1/x - \log(1 + \sqrt{1 + x^2})$ , then

$$h'(x) = \frac{1}{x} - \frac{1}{x^2} - \frac{x}{(1 + \sqrt{1 + x^2})\sqrt{1 + x^2}} \leq 0.$$

$$(\iff (x-1)(\sqrt{1+x^2} + 1 + x^2) - x^3 \leq 0 \iff (x1)\sqrt{1+x^2} \leq x^2)$$

Therefore  $h(x) \geq \lim_{x \rightarrow \infty} h(x) = 0$ .

**E6)**  $\log 8$  and  $2$ .

Solution: Note that

$$\log 2 = \log \frac{1}{1 - \frac{1}{2}} = \sum_{n=1}^{\infty} \frac{1}{2^n n} \geq \sum_{n=1}^3 \frac{1}{2^n n} = \frac{2}{3},$$

hence  $\log 8 > 2$ .

## 8.6 PSF

If  $f$  satisfy  $f(x) = (x - x_0)^r g(x)$  in a neighborhood of  $x_0$ , where  $r \in \mathbb{Z}_{\geq 0}$ ,  $g$  is continuous at  $x_0$  and  $g(x_0) \neq 0$ , then we call  $x_0$  an  $r$ -fold root of  $f$ .

**F1)** Suppose  $x_0$  is an  $r$ -fold root of  $f$  where  $r > 0$ . Prove that if  $g(x) = f(x)/(x - x_0)^r$  is continuous, then  $x_0$  is an  $(r - 1)$ -fold root of  $f'$ .

Proof: Suppose  $f(x) = (x - x_0)^r g(x)$  in the neighborhood  $O(x_0)$ , then  $f'(x) = (x - x_0)^r g'(x) + r(x - x_0)^{r-1} g(x)$  in  $O(x_0)$ . Therefore let  $h(x) = (x - x_0)g'(x) + g(x)$ , then  $f'(x) = (x - x_0)^{r-1} h(x)$  and  $h(x_0) = g(x_0) \neq 0$ , so  $x_0$  is an  $(r - 1)$ -fold root of  $f'$ .

**F2)** Suppose  $f$  is  $n$ -times differentiable on  $\mathbb{R}$ . Prove that if  $f(x) = 0$  has  $n + 1$  distinct real roots, then  $f^{(n)}(x) = 0$  has at least one root.

Proof: Use induction and Rolle's mean-value theorem to prove that  $f^{(n-k)}(x)$  has at least  $k + 1$  different real roots.

**F3)**  $f$  is differentiable on  $\mathbb{R}$ . Suppose  $f(x) = 0$  has  $r$  roots (counting multiplicity), then  $f'(x) = 0$  has at least  $r - 1$  roots (counting multiplicity).

Proof: Combine F1) and F2).

**F4)** Suppose  $f$  is  $n$ -times differentiable on  $\mathbb{R}$ . Prove that if  $f(x) = 0$  has exactly  $n + 1$  roots counting multiplicity, then  $f^{(n)}(x) = 0$  has at least one root.

Proof: Use F3) and induction.

## 8.7 PSG

Let  $a \in \mathbb{R}$ ,  $f : (a, \infty) \rightarrow \mathbb{R}$  twice differentiable on  $(a, \infty)$ , and

$$M_0 := \sup_{x \in (a, \infty)} |f(x)|, M_1 := \sup_{x \in (a, \infty)} |f'(x)|, M_2 := \sup_{x \in (a, \infty)} |f''(x)|,$$

are real numbers.

**G1) Prove that  $M_1^2 \leq 4M_0M_2$ .**

Proof: Let  $h = \sqrt{M_0/M_2}$ , then for any  $x \in (a, \infty)$ , there exists  $\xi \in (x, x+2h)$  such that

$$f(x+2h) = f(x) + 2hf'(x) + 2h^2f''(\xi) \implies f'(x) = hf''(\xi) + \frac{f(x+2h) - f(x)}{2h}.$$

Therefore  $f'(x) \leq M_0/h + M_2h = 2\sqrt{M_0M_2}$ , hence  $M_1^2 \leq 4M_0M_2$ .

**G2) Let  $a = -1$ , consider the function**

$$f(x) = \begin{cases} 2x^2 - 1, & x \in (-1, 0), \\ \frac{x^2-1}{x^2+1}, & x \in [0, \infty), \end{cases}$$

verify that  $f$  is twice differentiable, and  $M_0 = 1, M_1 = 4, M_2 = 4$ .

Proof: Note that  $\lim_{x \rightarrow 0^-} f(x) = -1 = f(0)$  so  $f$  is continuous, and

$$f'(x) = \begin{cases} 4x, & x \in (-1, 0), \\ \frac{4x}{(x^2+1)^2}, & x \in [0, \infty). \end{cases}$$

$f'$  is also continuous, so

$$f''(x) = \begin{cases} 4, & x \in (-1, 0), \\ 4\frac{1-3x^2}{(x^2+1)^2}, & x \in [0, \infty). \end{cases}$$

Therefore  $f \in C^2((-1, \infty))$  and  $M_0 = 1, M_1 = 4, M_2 = 4$ .

**G3) Suppose  $f : (a, \infty) \rightarrow \mathbb{R}^n$  is twice differentiable, Let  $M_0, M_1, M_2$  be the least upper bounds of  $|f|, |f'|, |f''|$ . Prove that  $M_1^2 \leq 4M_0M_2$ .**

Proof: Use G1) and Cauchy-Schwarz inequality.

## 8.8 Problem S: Strum-Liouville Theory

Assume the following uniqueness theorem holds:

Theorem:

Suppose  $a(t) \in C^1(\mathbb{R})$ ,  $t_0 \in \mathbb{R}$ . If  $x(t), y(t) \in C^2(\mathbb{R})$  both satisfy the equation

$$x''(t) + a(t)x(t) = 0, y''(t) + a(t)y(t) = 0,$$

and  $(x(t_0), x'(t_0)) = (y(t_0), y'(t_0))$ , then  $x(t) \equiv y(t)$ .

(Can be proved using Exercise C3?)

For any  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $t \geq 0$ , denote

$$Z_t(f) = |\{x \in [0, t] : f(x) = 0\}|.$$

### Part 1

Let  $a(t), b(t) \in C^1(\mathbb{R})$  and for any  $t \in \mathbb{R}$ ,  $a(t) \leq b(t)$ . Suppose  $x(t), y(t) \in C^2(\mathbb{R})$  satisfy the following equation

$$x''(t) + a(t)x(t) = 0, y''(t) + b(t)y(t) = 0.$$

Further assume that  $x(t), y(t)$  are not identically zero.

**S1) Assume  $x(t_1) = 0$ , if there exists  $t > t_1$ , such that  $x(t) = 0$ . Prove that there exists  $t_2 > t_1$  such that  $x(t_2) = 0$  and  $x$  has no roots in  $(t_1, t_2)$ . We call  $t_1, t_2$  neighboring roots.**

Proof: Consider the set  $S = \{t > t_1 : x(t) = 0\}$ , and let  $t_2 = \inf S$ . Note that  $|x''(t)| \leq |a(t)| \cdot |x(t)|$ , so by C3)  $x'(t_1) \neq 0$ . Assume  $x'(t_1) > 0$ , since  $x \in C^2(\mathbb{R})$ , there exists  $\varepsilon > 0$  such that  $x'(t) > 0$  for all  $t \in (t_1, t_1 + \varepsilon)$ , hence  $x(t) > 0$  for all  $t \in (t_1, t_1 + \varepsilon)$ . Therefore  $t_2 > t_1$ , so by  $x \in C(\mathbb{R})$ ,  $x(t_2) = 0$  and  $x$  has no roots in  $(t_1, t_2)$ .

**S2) If  $t_2 > t_1$  are two neighboring roots of  $x$ , prove that  $y$  has a root in  $(t_1, t_2]$ .**

Proof: Otherwise assume that  $x, y$  are positive on  $(t_1, t_2)$ , and  $y(t_2) \neq 0$ . Consider the function  $h(t) = x'y - xy'$ , then  $h'(t) = (b - a)xy \geq 0$ , so  $h(t_2) \geq h(t_1) = x'(t_1)y(t_1) \geq 0$ , but  $h(t_2) = x'(t_2)y(t_2) < 0$ , a contradiction.

**S3) Prove that for any  $t \geq 0$ ,  $Z_t(y) \geq Z_t(x) - 1$ .**

Proof: Use S2).

**S4) Suppose  $t_2 > t_1$  and  $x(t_1) = x'(t_2) = 0$ , prove that**

- If  $y(t_1) = 0$ , then there exists  $t_3 \in [t_1, t_2]$ , such that  $y'(t_3) = 0$ .  
Proof: We can assume that  $t_2 = \inf \{t > t_1 : x'(t) = 0\}$  ( $t_2 > t_1$  since  $x'(t_1) \neq 0$ ). If there is no such  $t_3$ , we can further assume that  $x'(t), y'(t), x(t), y(t) > 0$  for all  $t \in (t_1, t_2)$ . Again consider  $h(t) = x'y - xy'$ , then  $h(t_1) = 0$ ,  $h(t_2) = -x(t_2)y'(t_2) < 0$ , but  $h'(t) = (b - a)xy \geq 0$ , leading to contradiction.
- If  $y'(t_2) = 0$ , then there exists  $t_4 \in [t_1, t_2]$  such that  $y(t_4) = 0$ .  
(The two theorems are similar.)

## Part 2

In this section,  $p(t) \in C^1(\mathbb{R})$  is a positive function.  $x(t), y(t) \in C^2(\mathbb{R})$  are not identically zero and satisfy

$$x''(t) + p(t)x(t) = 0, \quad y''(t) + p(t)y(t) = 0.$$

**S5) Prove that for any  $t \geq 0$ ,  $|Z_t(x) - Z_t(y)| \leq 1$ .**

Proof: Use S3).

**S6) Prove that**

- If  $t_1, t_2$  are neighboring roots of  $x$ , then there exists a unique  $t_3 \in [t_1, t_2]$  such that  $x'(t_3) = 0$ .  
Proof: The existence of  $t_3$  is given by Rolle's mean-value theorem. If there exists  $t_3 < t_4 \in [t_1, t_2]$  such that  $x'(t_3) = x'(t_4) = 0$ , then  $t_3, t_4 \neq t_1, t_2$  and there exists  $t_5 \in [t_3, t_4]$  such that  $x''(t_5) = 0$ . Hence  $x(t_5) = 0$ , which contradicts the fact that  $t_1, t_2$  are neighboring roots. Therefore  $t_3$  is unique.
- If  $t'_1, t'_2$  are neighboring roots of  $x'$ , then there exists a unique  $t'_3 \in [t'_1, t'_2]$  such that  $x(t'_3) = 0$ .  
Proof: Exactly the same.

**S7) Prove that**

- $t_0$  is a local maximum of  $|x(t)|$  iff  $x'(t_0) = 0$ .  
Proof: Trivial?
- $t'_0$  is a local maximum of  $|x'(t)|$  iff  $x(t'_0) = 0$ .

**Part 3**

In this section,  $p(t) \in C^1(\mathbb{R})$  is monotonically decreasing and  $\lim_{t \rightarrow \infty} p(t) > 0$ . Denote

$$p(\infty) := \lim_{t \rightarrow \infty} p(t).$$

$x(t) \in C^2(\mathbb{R})$  is not identically zero and

$$x''(t) + p(t)x(t) = 0.$$

**\*S8) Calculate**

$$\lim_{t \rightarrow \infty} \frac{Z_t(x)}{t}.$$

Solution: By S5) we can ignore initial conditions. First consider the ODE  $y''(t) + p(\infty)y(t) = 0$ , where one solution is  $y = \sin(t\sqrt{p(\infty)})$ , so  $\lim_{t \rightarrow \infty} Z_t(y)/t = \sqrt{p(\infty)}/\pi$ .

Since  $p(t) \geq p(\infty)$ , by S3) we know  $\lim_{t \rightarrow \infty} Z_t(x)/t \geq \lim_{t \rightarrow \infty} Z_t(y)/t = \sqrt{p(\infty)}/\pi$ . For any  $\varepsilon > 0$ , there exists  $M > 0$  such that for any  $t > M$ ,  $p(t) < p(\infty) + \varepsilon$ . By S3),  $\lim_{t \rightarrow \infty} Z_t(x)/t \leq \sqrt{p(\infty) + \varepsilon}/\pi$ . Therefore

$$\lim_{t \rightarrow \infty} \frac{Z_t(x)}{t} = \frac{\sqrt{p(\infty)}}{\pi}.$$

**S9) Suppose  $0 \leq t_1 < t_2 < t_3 < \dots$  are all the roots of  $x(t)$  on  $[0, \infty)$ ,  $0 \leq t'_1 < t'_2 < \dots$  are all the roots of  $x'(t)$  on  $[0, \infty)$ . Prove that the sequence  $\{|x'(t_k)|\}_{k \geq 1}$  is monotonically decreasing and the sequence  $\{|x(t'_k)|\}_{k \geq 1}$  is monotonically increasing, and**

$$\lim_{k \rightarrow \infty} |x'(t_k)| = \sqrt{p(\infty)} \lim_{k \rightarrow \infty} |x(t'_k)|.$$

Proof: Consider the (energy) function  $E(t) = p(t)x^2(t) + x'(t)^2$ , then  $E'(t) = p'(t)x^2(t) \leq 0$  so  $E$  is monotonically decreasing. For  $k \geq 1$ ,  $E(t_k) = x'(t_k)^2$  is decreasing, so  $\{|x'(t_k)|\}_{k \geq 1}$  is decreasing. Likewise, consider  $F(t) = x(t)^2 + x'(t)^2/p(t)$ , then  $F'(t) = -p'(t)x(t)^2/p(t)^2 \geq 0$ , so  $F(t'_k) = x(t'_k)^2$  is increasing, and

$$\lim_{k \rightarrow \infty} |x'(t_k)| = \sqrt{\lim_{k \rightarrow \infty} E(t_k)} = \sqrt{p(\infty) \lim_{k \rightarrow \infty} F(t_k)} = \sqrt{p(\infty)} \lim_{k \rightarrow \infty} |x(t'_k)|.$$

**\*S10) Suppose  $0 \leq \tilde{t}_1 < \tilde{t}_2 < \dots$  are all the roots of  $x(t)x'(t)$  on  $[0, \infty)$ . Prove that the sequence  $\{\tilde{t}_{k+1} - \tilde{t}_k\}_{k \geq 1}$  is monotonically increasing and calculate its limit.**

Proof: By S6), the roots of  $x$  and  $x'$  appear alternating in  $\{\tilde{t}_k\}$ . Since  $t$  is a root of  $x$  iff  $t$  is a root of  $x''$ , we only need to prove that if  $t_1, t_2$  are neighboring roots of  $x$ , and  $t_3 \in [t_1, t_2]$  satisfy  $x'(t_3) = 0$ , then  $t_3 - t_1 \leq t_2 - t_3$ .

Same as before we can prove that, for  $p(t), q(t), x(t), y(t)$  such that  $p(0) = q(0)$ ,  $p(t) \leq q(t)$ ,  $x'(0) = y'(0) = 0$ ,  $x(0) = y(0)$  and

$$x''(t) + p(t)x(t) = 0, y''(t) + q(t)y(t) = 0,$$

then the first roots  $a, b$  of  $x, y$  satisfy  $a \leq b$ .

Since the sequence is increasing, by S8) we know that  $\lim_{k \rightarrow \infty} \tilde{t}_{k+1} - \tilde{t}_k = \frac{1}{2} \lim_{t \rightarrow \infty} Z_t(x)/t = \sqrt{p(\infty)}/2\pi$ .

## 9 Homework 9: Stone-Weierstrass Theorem

### 9.1 PSA

Assume  $I = [a, b] \subset \mathbb{R}$ ,  $V$  is a normed linear space.

**A1)**  $\sigma_1, \sigma_2 \in \mathcal{S}$  are two partitions. Prove that for any  $\varepsilon > 0$ , there exists a partition  $\sigma$  such that  $\sigma \prec \sigma_1, \sigma \prec \sigma_2$  and  $|\sigma| < \varepsilon$ .

Proof: Take  $n > 1/\varepsilon$ , and let

$$\sigma = \sigma_1 \cup \sigma_2 \cup \left\{ \frac{k}{n}a + \frac{n-k}{n}b : 0 \leq k \leq n \right\}.$$

**A2)** Consider the space of simple functions  $\mathcal{E}(I)$  with range  $V$ . Prove that it is a linear space on  $\mathbb{R}$ , and the integration operator  $\int_a^b : \mathcal{E}(I) \rightarrow V$  is well-defined and is linear. Use this to define Riemann integrable functions with range  $V$ .

Proof: For any simple function  $f = \sum_{i=1}^n c_i \mathbf{1}_{A_i}$  (where  $A_i$  are disjoint), let

$$\int_a^b f = \sum_{i=1}^n c_i \mu(A_i)$$

For any function  $f : I \rightarrow V$ , partition  $\mathcal{C} = \{x_0, x_1, \dots, x_n\}$  and  $\xi_i \in [x_{i-1}, x_i]$ , define

$$\mathcal{R}(f; \mathcal{C}, \xi) = \sum_{k=0}^n f(\xi_k)(x_k - x_{k-1}).$$

Then  $f$  is Riemann integrable iff  $\lim_{|\mathcal{C}| \rightarrow 0} \mathcal{R}(f; \mathcal{C}, \xi)$  exists.

**A3)** Suppose  $f : I \rightarrow \mathbb{R}^n$  and  $f_i$  be the components of  $f$ , then  $f \in \mathcal{R}(I)$  iff for every  $i$ ,  $f_i \in \mathcal{R}(I)$ .

Proof: Note that

$$\max\{|x_k|\} \leq |(x_1, \dots, x_n)|_{\mathbb{R}^n} \leq |x_1| + \dots + |x_n|.$$

Hence the limit  $|\underline{S}(f; \sigma) - \overline{S}(f; \sigma)| = 0$  iff the components of  $f$  are all Riemann integrable.

**A4) Assume  $a < c < b$ , then for any  $f \in \mathcal{R}(I)$ ,  $f|_{[a,c]}$  and  $f|_{[c,b]}$  are both Riemann integrable, and**

$$\int_a^b f = \int_a^c f + \int_c^b f.$$

Proof: They are both obviously Riemann integrable, and for any partition  $\sigma$ , let  $\sigma' = \sigma \cup \{c\} = \sigma_1 \cup \sigma_2$ , where  $\sigma_1$  and  $\sigma_2$  are partitions of  $[a, c]$  and  $[c, b]$ , then

$$\underline{S}(f; \sigma) \leq \underline{S}(f; \sigma') = \underline{S}(f|_{[a,c]}; \sigma_1) + \underline{S}(f|_{[c,b]}; \sigma_2),$$

and the other side is the same. Hence

$$\int_a^b f = \int_a^c f + \int_c^b f.$$

**A5) Prove that for any two partition  $\sigma$  and  $\sigma'$ ,  $\underline{S}(f; \sigma) \leq \overline{S}(f; \sigma')$ . Use this to prove that if  $f \in \mathcal{R}(I)$ , then  $\lim_{|\sigma| \rightarrow 0} |\underline{S}(f; \sigma) - \overline{S}(f; \sigma)| = 0$ .**

Proof: Let  $\sigma'' = \sigma \cup \sigma'$ , then

$$\underline{S}(f; \sigma) \leq \underline{S}(f; \sigma'') \leq \overline{S}(f; \sigma'') \leq \overline{S}(f; \sigma').$$

If  $f \in \mathcal{R}(I)$ , then  $\sup_{\sigma} \underline{S}(f; \sigma) = \inf_{\sigma} \overline{S}(f; \sigma)$  hence

$$\lim_{|\sigma| \rightarrow 0} |\underline{S}(f; \sigma) - \overline{S}(f; \sigma)| = 0.$$

**A6)  $f \in \mathcal{R}(I)$ . Prove that if we change the value of  $f$  at a finite number of points to  $g$ , then  $g$  is Riemann integrable and  $\int_I g = \int_I f$ .**

Proof: We can assume that  $f$  and  $g$  differ only at the point  $c \in (a, b)$ . Let  $M = \sup_{x \in I} |f(x)|$ . For any  $\varepsilon > 0$ , and any partition  $\sigma$ , let  $\sigma' = \sigma \cup \{c - \varepsilon, c + \varepsilon\}$ , then  $|\underline{S}(f; \sigma') - \underline{S}(f; \sigma)| \leq 4\varepsilon M \rightarrow 0$ .

**A7)  $f \in C([a, b])$ . Assume for any  $x \in I$ ,  $f(x) \geq 0$  and there exists  $x_0 \in I$  such that  $f(x_0) > 0$ . Prove that  $\int_a^b f > 0$ .**

Proof: Since  $f$  is continuous and  $f(x_0) > 0$ , there is an  $\varepsilon > 0$  such that for all  $y \in (x_0 - \varepsilon, x_0 + \varepsilon)$ ,  $f(y) > 0$ . Hence for any partition  $\sigma = \{x_0, x_1, \dots, x_n\}$  such that  $|\sigma| < \varepsilon/2$ , there is a  $k \in \{1, \dots, n\}$  such that  $(x_{k-1}, x_k) \subset (x_0 - \varepsilon, x_0 + \varepsilon)$ . Hence  $\mathcal{R}(f; \sigma, \xi) > 0$  whenever  $|\sigma| < \varepsilon/2$ , so  $\int_a^b f(x) dx > 0$ .

**A8) Suppose  $f, g \in C^1(I)$ , then**

$$\int f' \cdot g = f \cdot g - \int f \cdot g'.$$

Proof:

$$d(f \cdot g) = df \cdot g + f \cdot dg.$$



**A9)** Suppose  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  is differentiable,  $f$  is a continuous function, then

$$\int (f \circ \Phi) \Phi' = \int f.$$

Proof:

$$(f(\Phi(x)))' = f'(\Phi(x))\Phi'(x).$$

## 9.2 PSB: Calculating Integrals

(1)

$$\begin{aligned} \int \frac{x^5}{1+x} dx &= \int x^4 - x^3 + x^2 - x + 1 - \frac{1}{1+x} dx \\ &= \frac{x^5}{5} - \frac{x^4}{4} + \frac{x^3}{3} - \frac{x^2}{2} + x - \log(x+1) + C. \end{aligned}$$

(2)

$$\int \sqrt{x\sqrt{x\sqrt{x}}} dx = \int x^{7/8} dx = \frac{8}{15} x^{15/8}.$$

(3)

$$\begin{aligned} \int \left( \frac{1+x}{1-x} + \frac{1-x}{1+x} \right) dx &= \int \left( \frac{2}{1-x} + \frac{2}{1+x} - 2 \right) dx \\ &= -2x + 2 \log \frac{1+x}{1-x} + C. \end{aligned}$$

(4)

$$\int \frac{e^{3x} + 1}{1 + e^x} dx = \int 1 - e^x + e^{2x} dx = x - e^x + \frac{e^{2x}}{2} + C.$$

(5)

$$\int \sqrt{1 - \sin(2x)} dx = \int \sqrt{2} \sin \left( x - \frac{\pi}{4} \right) dx = -\sqrt{2} \cos \left( x - \frac{\pi}{4} \right) + C.$$

(6)

$$\int \frac{\cos(2x)}{\cos x - \sin x} dx = \int \cos x + \sin x dx = \sin x - \cos x + C.$$

(7)

$$\int \tan^2 x \, dx = -x + \tan x + C.$$

(8)

$$\int |x| \, dx = \frac{x|x|}{2} + C.$$

(9)

$$\int e^{-|x|} \, dx = -\operatorname{sgn}(x)e^{-|x|} + C.$$

(10)

$$\int \frac{x^2}{(1-x)^{2018}} \, dx = \frac{1}{2017(1-x)^{2017}} - \frac{1}{1013(1-x)^{2016}} + \frac{1}{2015(1-x)^{2015}}.$$

(11)

$$\int |x-1| \, dx = \frac{(x-1)|x-1|}{2} + C.$$

(12)

$$\int \frac{1}{\sqrt{b^2+x^2}} \, dx = \frac{1}{b} \log \frac{1 + \tan \frac{\arctan \frac{x}{b}}{2}}{1 - \tan \frac{\arctan \frac{x}{b}}{2}} + C.$$

(13)

Let  $x = t^2$ , then

$$\int \frac{dx}{\sqrt{x}(1+x)} = 2 \arctan \sqrt{x} + C.$$

(14)

$$\int \frac{x^4}{(1-x^5)^4} \, dx = \frac{1}{5} \int \frac{dx^5}{(1-x^5)^4} = \frac{1}{15(1-x^5)^3} + C.$$

(15)

$$\int \left( \frac{1}{\sqrt{3-x^2}} + \frac{1}{1-3x^2} \right) dx = \arcsin \frac{x}{\sqrt{3}} + \frac{1}{2\sqrt{3}} \log \frac{1+\sqrt{3}x}{1-\sqrt{3}x} + C.$$

(16)

$$\int \frac{2x-3}{x^2-3x+8} dx = \log(x^2-3x+8) + C.$$

(17)

$$\int \frac{dx}{\sin^2(2x + \frac{\pi}{4})} = \frac{\tan(2x - \pi/4)}{2} + C.$$

(18)

$$\int \frac{dx}{1 + \cos x} = \tan \frac{x}{2} + C.$$

(19)

$$\int \frac{1}{x^2} \sin \frac{1}{x} dx = \cos \frac{1}{x} + C.$$

(20)

$$\int \cos^5 x dx = \frac{\sin^5 x}{5} - \frac{2 \sin^3 x}{3} + \sin x + C.$$

(21)

$$\int \cos(ax) \sin(bx) dx = \frac{\cos(a-b)x}{2(a-b)} - \frac{\cos(a+b)x}{2(a+b)} + C.$$

(22)

$$\int \frac{dx}{a \cos x + b \sin x} = \frac{2}{\sqrt{a^2 + b^2}} \tanh^{-1} \frac{a \tan(x/2) - b}{\sqrt{a^2 + b^2}} + C.$$

(23)

$$\int \frac{\sin(2x)}{a^2 \cos^2 x + b^2 \sin^2 x} dx = \frac{\log((b^2 - a^2) \sin^2 x + a^2)}{b^2 - a^2} + C.$$

(24)

$$\int \frac{dx}{2 - \sin^2 x} = \frac{1}{\sqrt{2}} \arctan \left( \frac{\tan x}{\sqrt{2}} \right) + C.$$

(25)

$$\int \frac{dx}{x \ln x \ln \ln x} = \ln \ln \ln x + C.$$

(26)

$$\int \frac{\log x}{x\sqrt{1+\log x}} dx = \frac{2}{3}(1+\log x)^{3/2} - 2\sqrt{1+\log x} + C.$$

(27)

$$\int \frac{\cos x + \sin x}{(\sin x - \cos x)^{1/3}} dx = \frac{3}{2}(\sin x - \cos x)^{2/3} + C.$$

(28)

$$\int e^{\sqrt{x}} dx = 2\sqrt{x}e^{\sqrt{x}} - 2e^{\sqrt{x}} + C.$$

(29)

$$\int \frac{x^{n/2}}{1+x^{n+2}} dx = \frac{2}{n+2} \arctan x^{n/2+1} + C.$$

(30)

$$\int \frac{\sqrt{x}}{1-x^{1/3}} dx = 6 \arctan x^{1/6} - \frac{6}{5}x^{5/6} - \frac{6}{7}x^{7/6} - 2x^{1/2} - 6x^{1/6} + C.$$

(31)

$$\int \frac{dx}{(x^2+a^2)^{3/2}} = \frac{x}{a^2\sqrt{a^2+x^2}} + C.$$

(32)

$$\int \frac{dx}{\cos^4 x} = \frac{\sin x}{2 \cos^3 x} + \frac{\sin(3x)}{6 \cos^3 x} + C.$$

(33)

$$\int \arcsin^2 x dx = x \arcsin^2 x + 2\sqrt{1-x^2} \arcsin x - 2x + C.$$

(34)

$$\int x \arcsin x \, dx = \frac{x\sqrt{1-x^2}}{4} - \frac{1}{4} \arcsin x (1-2x^2) + C.$$

(35)

$$\int x \arctan x = \frac{1}{2}(x^2+1) \arctan x - \frac{1}{2}x + C.$$

(36)

$$\int \frac{\arctan x}{x^2} = \log x - \frac{\arctan x}{x} - \frac{1}{2} \log(1+x^2) + C.$$

(37)

$$\int x^2 \sin x = -x^2 \cos x + 2x \sin x + 2 \cos x + C.$$

(38)

$$\int \frac{x}{\cos^2 x} = x \tan x + \log \cos x + C.$$

(39)

$$\int \log(x + \sqrt{1+x^2}) = x \log(x + \sqrt{1+x^2}) - \sqrt{1+x^2} + C.$$

(40)

$$\int \sin \log x = \frac{x}{2} (\sin \log x - \cos \log x) + C.$$

(41)

$$\int \sqrt{x^2+a^2} = \frac{1}{2}x\sqrt{x^2+a^2} + \frac{a^2}{4} \log \frac{x+\sqrt{x^2+a^2}}{\sqrt{x^2+a^2}-x} + C.$$

(42)

$$\int \frac{x^2}{\sqrt{x^2-a^2}} = \frac{1}{2}x\sqrt{x^2-a^2} + \frac{a^2}{4} \log \frac{x+\sqrt{x^2-a^2}}{x-\sqrt{x^2-a^2}} + C.$$

(43)

$$\int \frac{x \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} = \sqrt{x^2+1} \log(x + \sqrt{1+x^2}) - x + C.$$

Let  $u = \sqrt{x^2+1} + x$  then  $du/dx = 1 + x/\sqrt{1+x^2}$ , so it becomes

$$\int \frac{(u^2-1) \log u}{2u^2} dx = -\frac{u}{2} + \frac{1}{2u} + \frac{1}{2} u \log u + \frac{\log u}{2u} + C.$$

(44)

$$\int \frac{1}{\sqrt{x^2+a^2}} = \log \frac{\sin t + \cos t}{\sin t - \cos t} + C = \tanh^{-1} \frac{x}{\sqrt{x^2+a^2}} + C.$$

where  $t = \frac{1}{2} \arctan(x/a)$ .

(45)

$$\int \frac{xe^x}{(1+x)^2} = \frac{e^x}{1+x} + C.$$

(46)

$$\int \arctan(1 + \sqrt{x}) = x \arctan(1 + \sqrt{x}) - \sqrt{x} + \log(2 + 2\sqrt{x} + x) + C.$$

(47)

$$\int \left(1 - \frac{2}{x}\right)^2 e^x = e^x - \frac{4e^x}{x} + C.$$

since  $\int e^x/x^2 dx = -e^x/x + \int e^x/x dx$ .

(48)

$$\int \sqrt{2 + \tan^2 x} = \theta + \log \frac{\sin \theta + \cos \theta}{\sin \theta - \cos \theta} + C.$$

where  $\theta = \arcsin(\sin x/\sqrt{2})$ .

(49)

$$\int \frac{1}{1+x^3} = -\frac{1}{6} \log(x^2-x+1) + \frac{1}{3} \log(x+1) + \frac{1}{\sqrt{3}} \arctan \frac{2x-1}{\sqrt{3}} + C.$$

(50)

$$\int \frac{x^7}{x^4+2} = \frac{x^4}{4} - \frac{1}{2} \log(2+x^4).$$

(51)

$$\int \frac{2x^2 + 1}{(x+3)(x-1)(x-4)} = -\frac{1}{4} \log(1-x) + \frac{11}{7} \log(4-x) + \frac{19}{28} \log(x+3) + C.$$

(52)

$$\int \frac{1+x^2}{1+x^4} = \frac{1}{\sqrt{2}} (\arctan(\sqrt{2}x+1) - \arctan(1-\sqrt{2}x)) + C.$$

Note that

$$\frac{1+x^2}{1+x^4} = \frac{1}{2(x^2+\sqrt{2}x+1)} - \frac{1}{2(-x^2+\sqrt{2}x+1)}.$$

(53)

Let  $x = y^6 - 1$  then

$$\begin{aligned} \int \frac{x}{\sqrt{x+1} + (x+1)^{1/3}} &= \int (y^3 - 1)(1 - y + y^2) 6y^3 dy \\ &= \frac{2x\sqrt{x+1}}{3} - \frac{3x(x+1)^{1/3}}{4} + \frac{6x(x+1)^{1/6}}{7} - x + \frac{6}{5}(x+1)^{5/6} \\ &\quad - \frac{3}{2}(x+1)^{2/3} + \frac{2\sqrt{x+1}}{3} - \frac{3(x+1)^{1/3}}{4} + \frac{6(x+1)^{1/6}}{7} + C. \end{aligned}$$

(54)

Let  $x = y^2$ , then

$$\int \frac{1}{\sqrt{x+x^2}} = \int \frac{dy}{\sqrt{y^2+1}} = \tanh^{-1} \left( \sqrt{\frac{x}{x+1}} \right) + C.$$

(55)

The Poisson kernel

$$\int \frac{1-r^2}{1-2r \cos x + r^2} = 2 \arctan \left( \frac{1+r}{1-r} \tan \frac{x}{2} \right) + C.$$

(56)

Let  $x = \tan \theta$  then

$$\int \frac{1}{x\sqrt{1+x^2}} = \int \frac{d\theta}{\sin \theta} = \log \tan \frac{\arctan x}{2} + C.$$

(57)

Let  $t = \tan x/2$  then

$$\int \frac{1}{5-3 \cos x} = \frac{1}{2} \arctan \left( 2 \tan \frac{x}{2} \right) + C.$$

(58)

Let  $t = \tan x$ , then

$$\int \frac{1}{2 + \sin^2 x} = \frac{1}{\sqrt{6}} \arctan\left(\sqrt{\frac{3}{2}} \tan x\right) + C.$$

(59)

$$\int \frac{\sin^3 x}{\cos^4 x} = \frac{1}{3 \cos^3 x} - \frac{1}{\cos x} + C.$$

(60)

Let  $t = \cos x$  then

$$\int \frac{1}{\sin x \cos^4 x} = - \int t^{-4} + t^{-2} + \frac{1}{1-t^2} = \frac{1}{3 \cos^3 x} + \frac{1}{\cos x} + \frac{1}{2} \log \frac{1 + \cos x}{1 - \cos x} + C.$$

### 9.3 Problem W: Stone-WeierstraSS Theorem

#### Part 1: Approximating $|x|$

**W1) (Dini)** Suppose  $K \subset \mathbb{R}^n$  is compact,  $f_n : K \rightarrow \mathbb{R}$  is a sequence of continuous functions, which converges point-wise to  $f : K \rightarrow \mathbb{R}$ . If  $f$  is continuous and  $f_n \leq f_{n+1}$ , then  $f_n$  converges uniformly to  $f$ .

Proof: For any  $\varepsilon > 0$ , and any  $x \in K$ , there is an integer  $n_x > 0$  such that  $|f_{n_x}(x) - f(x)| < \varepsilon/4$ . There exists  $\delta > 0$ , such that  $\forall y \in B(x, \delta) \cap K$ ,  $|f(x) - f(y)| < \varepsilon/4$  and  $|f_{n_x}(x) - f_{n_x}(y)| < \varepsilon/4$ , then  $|f_{n_x}(y) - f(y)| < \varepsilon/4$ . Note that  $K \subset \bigcup_{x \in K} B(x, \delta_x)$  hence we can choose a finite set of  $x_1, x_2, \dots, x_N$  such that  $K \subset \bigcup_{i=1}^N B(x_i, \delta_{x_i})$ . Let  $M = \max\{n_{x_i} : i = 1, 2, \dots, N\}$  then for any  $m \geq M$  and  $x \in K$ ,  $|f_m(x) - f(x)| < \varepsilon$ . Hence  $f_n$  converges uniformly to  $f$ .

**W2) Consider the interval  $[-1, 1]$ . Define inductively a sequence of polynomials:**

$$P_0(x) = 0, P_{n+1}(x) = P_n(x) + \frac{1}{2}(x^2 - P_n^2(x)).$$

Prove that for any  $n, x$ ,  $0 \leq P_n(x) \leq P_{n+1}(x) \leq |x|$ .

Proof: Assume  $x > 0$ , we prove by induction. If  $t = P_n(x) \in [0, x]$ , then

$$P_{n+1}(x) = \frac{1}{2}x^2 - \frac{1}{2}(t-1)^2 + \frac{1}{2} \leq \frac{1}{2}(x^2 - (1-x)^2 + 1) = x,$$

and  $P_{n+1}(x) \geq P_n(x) = t$ , hence  $P_{n+1}(x) \in [0, x]$ .

**W3) Prove that  $|x|$  can be uniformly approximated by polynomials on the interval  $[-1, 1]$ , i.e. for any  $\varepsilon > 0$ , there exists a polynomial  $P_\varepsilon(x)$  such that  $\sup_{x \in [-1, 1]} ||x| - P_\varepsilon(x)| < \varepsilon$ .**

Proof: By W2), the sequence of polynomials  $\{P_n\}$  converge point-wise to  $|x|$ , hence by W1)  $P_n$  converge uniformly to  $|x|$ .



**Part 3: Bernstein Polynomial**

Assume  $I = [0, 1]$ , and  $n$  is an integer.

**W4) For any  $0 \leq k \leq n$ , define  $p_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}$ . Prove that**

$$\sum_{0 \leq k \leq n} p_{n,k}(x) \left(x - \frac{k}{n}\right)^2 = \frac{x(1-x)}{n}.$$

Proof:

Note that

$$\sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} x^2 = x^2,$$

$$\sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} \frac{k}{n} = \sum_{k=1}^n \binom{n-1}{k-1} x^k (1-x)^{n-k} = x,$$

$$\sum_{k=0}^n p_{n,k}(x) k(k-1) = n(n-1) \sum_{k=2}^n \binom{n-2}{k-2} x^k (1-x)^{n-k} = n(n-1)x^2.$$

Therefore

$$\sum_{k=0}^n p_{n,k}(x) \left(x - \frac{k}{n}\right)^2 = \frac{x(1-x)}{n}.$$

**W5) For any  $f \in C([0, 1])$ , define**

$$B_{f,n} = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}.$$

For  $x \in [0, 1]$ , prove that

$$|f(x) - B_{f,n}(x)| \leq \sum_{k=0}^n \left|f(x) - f\left(\frac{k}{n}\right)\right| p_{n,k}(x).$$

Proof: Note that

$$\sum_{k=0}^n f(x) \binom{n}{k} x^k (1-x)^{n-k} = f(x).$$

**W6) For any  $f \in C([0, 1])$ , prove that for any  $\varepsilon > 0$ , there exists  $n$  such that  $\|f - B_{f,n}\|_\infty < \varepsilon$ .**

Proof:

Let

$$\begin{aligned} \text{I} &= \sum_{|m-nx| < n^{3/4}} \left(f(x) - f\left(\frac{m}{n}\right)\right) p_{n,m}(x), \\ \text{II} &= \sum_{|m-nx| > n^{3/4}} \left(f(x) - f\left(\frac{m}{n}\right)\right) p_{n,m}(x). \end{aligned}$$

Then  $|f - B_{f,n}| \leq |I| + |II|$ .

For any  $\varepsilon > 0$ ,  $\exists N \in \mathbb{N}$  such that  $\forall x \in [0, 1]$ ,  $n \geq N \implies |I| < \varepsilon$ , since

$$|I| \leq \sup_{|x - m/n| < n^{-1/4}} |f(x) - f(m/n)| \rightarrow 0.$$

Suppose  $M = \sup_{x \in [0, 1]} |f(x)|$ , then

$$|II| \leq 2M \sum_{|m - nx| > n^{3/4}} p_{n,m}(x) \leq 2M \sqrt{n} \sum_{m=0}^n (x - m/n)^2 p_{n,m}(x) = \frac{2Mx(1-x)}{\sqrt{n}}.$$

Hence  $\|f - B_{f,n}\|_{\infty} \rightarrow 0$ .

### Part 3: Stone-Weierstrass Theorem

#### W7-14):

Let  $X$  be a compact Hausdorff space,  $\mathcal{A} \subset C(X, \mathbb{R})$  satisfy the following properties:

- (a)  $\forall c \in \mathbb{R}$ ,  $c \cdot 1_X \in \mathcal{A}$ , (b)  $\forall f, g \in \mathcal{A}$ ,  $f + g, f - g, fg \in \mathcal{A}$ ,
- (c)  $\mathcal{A}$  can separate any pair of points in  $X$ .

Then  $\bar{\mathcal{A}} = C(X, \mathbb{R})$ .

#### Lemma 1

There is a list of polynomials  $\{P_n(x)\}$  that converges uniformly to  $|x|$  on  $[-1, 1]$ .

#### Lemma 2

If  $\mathcal{A}$  is a subspace of  $C(X, \mathbb{R})$ , such that (a)  $\mathcal{A}$  is a lattice, (b)  $1_X \in \mathcal{A}$ , and (c)  $\mathcal{A}$  can separate any pair of points, then  $\bar{\mathcal{A}} = C(X, \mathbb{R})$ .

#### Proof of main theorem

Assume WLOG  $\mathcal{A}$  is closed, then by Lemma 1,  $\forall f \in \mathcal{A}$ ,  $P_n(f) \in \mathcal{A}$ , hence  $|f| \in \mathcal{A}$ . (Since  $X$  is compact,  $|f|$  is bounded.) Note that

$$\max\{f, g\} = \frac{1}{2}(|f + g| + |f - g|), \min\{f, g\} = \frac{1}{2}(|f + g| - |f - g|).$$

Hence  $\mathcal{A}$  is a lattice, by Lemma 2  $\mathcal{A} = C(X, \mathbb{R})$ .

#### Proof of Lemma 1

Proof 1: Let

$$Q_n(x) = \int_0^x (1 - t^2)^n dt / \int_0^1 (1 - t^2)^n dt.$$

$$P_n(x) = \int_0^x Q_n(t) dt.$$

Note that

$$\int_{\varepsilon}^1 (1 - t^2)^n dt \leq (1 - \varepsilon^2)^n (1 - \varepsilon) \rightarrow 0$$

Hence (combined with Wallis's formula),  $P_n(x)$  converges uniformly to  $|x|$  on  $[a, b]$ .

Proof 2: WLOG change the interval to  $[-1/2, 1/2]$ . The series

$$(1-t)^{1/2} = 1 + \sum_{n=1}^{\infty} (-t)^n \binom{\frac{1}{2}}{n} = 1 - \sum_{n=1}^{\infty} c_n t^n.$$

converges when  $|t| < 1$ . Hence  $\forall \varepsilon > 0$ , there exists  $Q \in \mathbb{R}[x]$  such that  $\sup_{|t| \leq 1/2} |Q(t) - (1-t)^{1/2}| < \varepsilon/2$ .

Let  $t = 1 - x^2$ , then  $|Q(1 - x^2) - |x|| < \varepsilon/2$ , so  $P(x) = Q(1 - x^2) - Q(1)$  converges to  $|x|$  uniformly on  $[-1/2, 1/2]$ .

### Proof of Lemme 2

Step 1: Take any  $f \in C(X, \mathbb{R})$ , and any  $x, y \in X$ , we can find  $g_{xy} \in \mathcal{A}$ , such that  $g_{xy}(x) = f(x)$ ,  $g_{xy}(y) = f(y)$ . Since there exists  $u \in \mathcal{A}$  such that  $u(x) \neq u(y)$ ,

$$\begin{pmatrix} u(x), 1 \\ u(y), 1 \end{pmatrix} \begin{pmatrix} \lambda \\ \mu \end{pmatrix} = \begin{pmatrix} f(x) \\ f(y) \end{pmatrix}$$

has a solution. (If  $x = y$  it is trivial.)

Step 2:

For all  $\varepsilon > 0$ ,  $x, y \in X$ , there is an open neighborhood  $O_{x,y}$  of  $y$ , such that  $\forall z \in O_{x,y}$ ,  $f(z) - g_{xy}(z) \leq \varepsilon$ . Note that  $\bigcup_{y \in X} O_{x,y} = X$ , so by  $X$  is compact, there is a list  $y_1, \dots, y_N$  such that  $\bigcup_{k \leq N} O_{x,y_k} = X$ . Let  $h_x = \max \{g_{xy_k} : k \leq N\}$ , then  $h_x(y) - f(y) \geq -\varepsilon$ , and  $f(x) = h_x(x)$ .

Step 3:

For all  $x \in X$ , there is an open neighborhood  $G_x$  of  $x$ , such that  $\forall z \in G_x$ ,  $h_x(z) - f(z) \leq \varepsilon$ . Note that  $\bigcup_{x \in X} G_x = X$ , so by  $X$  is compact, there is a list  $x_1, \dots, x_M$  such that  $\bigcup_{k \leq M} G_{x_k} = X$ . Let  $F = \min \{h_{x_k} : k \leq M\}$ , then  $|F(x) - f(x)| \leq \varepsilon, \forall x \in X$ .

Therefore  $\bar{\mathcal{A}} = C(X, \mathbb{R})$ .

For complex numbers, there is an additional requirement: for any  $f \in \mathcal{A}$ ,  $\bar{f} \in \mathcal{A}$ .

### W15-16):

It is easy to see that polynomials and trigonometric polynomials both satisfy the requirements of the theorem.

## 10 Homework 10: Irrationality of $\pi$

### 10.1 PSA

**A1) Construct continuous functions  $f_n, f \in C([0, 1])$ , such that for every  $x \in [0, 1]$ , when  $n \rightarrow \infty$ ,  $f_n(x) \rightarrow f(x)$ , but**

$$\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx \neq \int_0^1 f(x) dx.$$

Solution: Let  $f_n(x) = nxe^{-nx^2}$ ,  $f(x) = \lim_{n \rightarrow \infty} f_n(x) = 0$ . Then

$$\lim_{n \rightarrow \infty} \int_0^1 f_n(x) dx = \lim_{n \rightarrow \infty} \frac{n}{2} \int_0^1 e^{-nx^2} d(x^2) = \lim_{n \rightarrow \infty} \frac{n}{2} \left( \frac{1}{n} - \frac{1}{ne^n} \right) = \frac{1}{2}$$

Hence  $\lim_{n \rightarrow \infty} \int_0^1 f_n = 1/2 \neq 0 = \int_0^1 f$ .

**A2)**  $\alpha \in \mathbb{R}_{\geq 0}$ . **Prove that**  $\int_{100}^{\infty} \frac{dx}{x \log^{\alpha}(x)}$  **converges iff**  $\alpha > 1$ .

Proof: Substitute  $y = \log x$ , then

$$\int_{100}^{\infty} \frac{dx}{x \log^{\alpha}(x)} = \int_{\log 100}^{\infty} \frac{dy}{y^{\alpha}}$$

which converges iff  $\alpha > 1$ .

**A3)**  $f, F$  are defined on  $I$ , and for every bounded closed interval  $J \subset I$ ,  $f, F$  are both Riemann integrable on  $J$ . Assume for all  $x \in I$ ,  $|f(x)| \leq F(x)$ . Then if the improper integral of  $F$  on  $I$  converges, so does  $f$ .

Proof: This is because

$$\int_I f(x) dx \text{ converges} \iff \forall \varepsilon > 0 \exists N \forall u, v \in I, N < u < v, \left| \int_u^v f(x) dx \right| < \varepsilon.$$

**A4)** Prove the integrals below converge:

(1)  $\int_0^{\infty} e^{-x^2} dx$  (2)  $\int_0^1 \frac{dx}{\sqrt{1-x^3}}$  (3)  $\int_1^{\infty} \frac{(\log x)^2}{1+x(\log x)^5} dx$   
(1):

$$\int_0^{\infty} e^{-x^2} dx \leq 1 + \int_1^{\infty} e^{-x} dx \leq 1 + \frac{1}{e}.$$

(2):

$$\int_0^1 \frac{dx}{\sqrt{1-x^3}} \leq \int_0^1 \frac{dx}{\sqrt{1-x}} = 2.$$

(3):

$$\int_1^{\infty} \frac{(\log x)^2}{1+x(\log x)^5} dx \leq 5000 + \int_{100}^{\infty} \frac{1}{x(\log x)^3} dx, \text{ which converges by A2.}$$

**A5)** Prove the series below converge:

(1)  $\sum_{n=1}^{\infty} e^{-n}(n^2 + \log n)$  (2)  $\sum_{n=1}^{\infty} \frac{\log n}{1+n(\log n)^3}$   
(1):

$$\sum_{n=1}^{\infty} e^{-n}(n^2 + \log n) \leq \sum_{n=1}^{\infty} \frac{2n^2}{e^n} \leq 2 \int_0^{\infty} x^2 e^{-x} dx = 4.$$

(2):

$$\sum_{n=1}^{\infty} \frac{\log n}{1+n(\log n)^3} \leq \sum_{n=2}^{\infty} \frac{1}{n(\log n)^2} \leq \frac{1}{2(\log 2)^2} + \int_2^{\infty} \frac{1}{x(\log x)^2} dx \leq 3.$$

**A6)** Calculate

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{k^{\alpha}}{n^{\alpha+1}}, \alpha > -1.$$

Solution:

$$\sum_{k=1}^n k^\alpha \leq \int_1^{n+1} x^\alpha dx = \frac{1}{\alpha+1}((n+1)^{\alpha+1} - 1).$$

$$\sum_{k=1}^n k^\alpha \geq 1 + \int_1^n x^\alpha dx = 1 + \frac{1}{\alpha+1}n^{\alpha+1}.$$

Therefore

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{k^\alpha}{n^{\alpha+1}} = \frac{1}{\alpha+1}.$$

**A7) Calculate  $\int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx$ , to show that  $\pi = 3.14 \dots$ .**

Solution:

$$\int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx = \int_0^1 x^6 - 4x^5 + 5x^4 - 4x^2 + 4 - \frac{4}{1+x^2} dx = \frac{22}{7} - \pi.$$

$$\int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx \leq \int_0^1 \frac{x^3(1-x)^4}{2} dx = \frac{1}{560} < 0.02, \frac{22}{7} > 3.1428.$$

**A8) Assume  $a, b, n \in \mathbb{Z}$ , let**

$$f_{a,b;n} = \frac{x^n(a-bx)^n}{n!}.$$

- Prove that for  $k = 0, 1, \dots, 2n$ ,  $f_{a,b;n}^{(k)}(x) \in \mathbb{Z}$  when  $x = 0, \frac{a}{b}$ .  
See B10)
- If  $\pi = \frac{a}{b} \in \mathbb{Q}$ , then for every  $n \in \mathbb{N}$ ,

$$\int_0^\pi f_{a,b;n}(x) \sin x dx$$

is an integer.

Proof: By Darboux's formula of integration of parts

$$\int_0^\pi f_{a,b;n}(x) \sin x dx = \sum_{k=0}^{2n} f_{a,b;n}^{(k)}(x) \sin \left( x - \frac{(k+1)\pi}{2} \right) \Big|_0^\pi \in \mathbb{Z}.$$

- Prove that  $\pi \notin \mathbb{Q}$ .  
Proof: Let  $n = 2a^4 + 10$ , then  $\forall 0 \leq x \leq a/b$ ,

$$f_{a,b;n} \leq \frac{a^{2n}}{n!} < \frac{1}{2} \frac{(a^4)^{n/2}}{n \cdot (n-1) \cdots \left(\frac{n}{2}\right)} < \frac{1}{2}.$$

Hence

$$0 < \int_0^\pi f_{a,b;n}(x) \sin x dx < \frac{1}{2} \int_0^\pi \sin x dx = 1,$$

leading to contradiction.

**A9) Let  $I_n = \int_0^{\pi/2} \sin^n x \, dx$ , prove that  $I_n \sim \sqrt{\frac{\pi}{2n}}$ .**

Proof: Since  $I_n = \frac{n-1}{n} I_{n-2}$ ,

$$I_n = \begin{cases} \frac{(n-1)!!}{n!!}, & n \text{ is even,} \\ \frac{(n-1)!!}{n!!} \cdot \frac{\pi}{2}, & n \text{ is odd.} \end{cases}$$

Combined with  $I_{2n+1} < I_{2n} < I_{2n-1}$ , we get

$$\left[ \frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n+1} < \frac{\pi}{2} < \left[ \frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n},$$

where

$$0 < - \left[ \frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n+1} + \left[ \frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n} = \left[ \frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n(2n+1)} < \frac{\pi}{4n}.$$

Therefore

$$\lim_{n \rightarrow \infty} \left[ \frac{(2n)!!}{(2n-1)!!} \right]^2 \frac{1}{2n} = \frac{\pi}{2}.$$

Hence  $I_n \sim \sqrt{\frac{\pi}{2n}}$ .

**A10) Assume  $f : [0, 1] \rightarrow [0, 1]$  is monotonously increasing,  $g = f^{-1} : [0, 1] \rightarrow [0, 1]$  is its inverse, and  $f, g$  are both continuously differentiable, then**

$$\int_0^1 f(x) \, dx + \int_0^1 g(x) \, dx = 1.$$

Proof: We show that

$$\int_0^x f(t) \, dt + \int_0^{f(x)} g(t) \, dt = xf(x), \forall 0 \leq x \leq 1. (1)$$

$x = 0$  is trivial, hence it suffices to show that the derivatives of the two sides match.

$$\frac{d}{dx} \int_0^x f(t) \, dt = f(x), \frac{d}{dx} \int_0^{f(x)} g(t) \, dt = f'(x) \cdot g(f(x)) = xf'(x).$$

Hence (1) holds.

**A11) Prove that**

$$\lim_{\varepsilon \rightarrow 0} \sum_{k=0}^{\infty} \frac{(-1)^k (1-\varepsilon)^{2k+1}}{2k+1} = \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1}.$$

Proof: By Dirichlet's test,  $\sum_{k=0}^{\infty} (-1)^k (1-\varepsilon)^{2k+1} / (2k+1)$  converges uniformly.

Hence for any  $\delta > 0$ , there exists an  $N \in \mathbb{Z}$  such that

$$\left| \sum_{k=N}^{\infty} \frac{(-1)^k x^{2k}}{2k+1} \right| < \delta, \forall x \in [0, 1].$$

Then  $\forall \varepsilon < \frac{\delta}{N}$ ,

$$\begin{aligned} & \left| \sum_{k=0}^{\infty} \frac{(-1)^k (1-\varepsilon)^{2k+1}}{2k+1} - \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} \right| \\ & \leq \sum_{k=0}^{N-1} \frac{|(1-\varepsilon)^{2k+1} - 1|}{2k+1} + \left| \sum_{k=N}^{\infty} \frac{(-1)^k (1-\varepsilon)^{2k+1}}{2k+1} \right| + \left| \sum_{k=N}^{\infty} \frac{(-1)^k}{2k+1} \right| < 3\delta. \end{aligned}$$

Hence

$$\lim_{\varepsilon \rightarrow 0} \sum_{k=0}^{\infty} \frac{(-1)^k (1-\varepsilon)^{2k+1}}{2k+1} = \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1}.$$

**A12) For any continuous function  $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ ,  $(x, y) \mapsto f(x, y)$ , where  $a, b, c, d \in \mathbb{R}$ , show that  $f$  is uniformly continuous on  $[a, b] \times [c, d]$ .**

Proof:  $K = [a, b] \times [c, d]$  is a compact set. Consider an arbitrary  $\varepsilon > 0$ .

For any  $x \in K$ , there is an open ball  $B(x, 2r_x)$  with center  $x$  such that  $\forall y \in B(x, 2r_x), |f(x) - f(y)| < \varepsilon/2$ . Let  $O_x = B(x, r_x)$ . Note that  $\bigcup_{x \in K} O_x = K$  and  $K$  is compact, hence we can find  $x_1, \dots, x_n$  such that  $\bigcup_{k \leq n} O_{x_k} = K$ .

Let  $\delta = \min\{r_{x_k} : k \leq n\}$ , then  $\forall |u - v| < \delta$ , suppose  $u \in O_{x_1}$ , then

$$|v - x_1| \leq |v - u| + |u - x_1| < 2r_{x_1} \implies v \in B(x_1, 2r_{x_1}).$$

Hence

$$|f(u) - f(v)| \leq |f(u) - f(x_1)| + |f(v) - f(x_1)| < \varepsilon.$$

Therefore  $f$  is uniformly continuous on  $K$ .

## 10.2 PSB: On $\zeta(2)$

**Part 1: The sequence  $\{\sum_{k=1}^n 1/k^p\}$**

Define the sequence  $S_n(p) = \sum_{k=1}^n 1/k^p$  where  $p \in \mathbb{Z}_{\geq 1}$ .

**B1) Prove that for any  $k \in \mathbb{Z}_{\geq 1}$ , we have**

$$\frac{1}{(k+1)^p} \leq \int_k^{k+1} \frac{1}{x^p} dx \leq \frac{1}{k^p}.$$

Proof:  $\frac{1}{(k+1)^p} \leq \frac{1}{x^p} \leq \frac{1}{k^p}, \forall k \leq x \leq k+1$ .

**B2) Prove that for any  $n \in \mathbb{Z}_{\geq 2}$ , we have**

$$S_n(p) - 1 \leq \int_1^n \frac{1}{x^p} dx \leq S_{n-1}(p).$$

Proof:

$$S_n(p) - 1 = \sum_{k=1}^{n-1} \frac{1}{(k+1)^p} \leq \sum_{k=1}^{n-1} \int_k^{k+1} \frac{1}{x^p} dx = \int_1^n \frac{1}{x^p} dx.$$

Likewise we have  $\int_1^n \frac{1}{x^p} dx \leq S_{n-1}(p)$ .

**B3) Let  $p \in \mathbb{Z}_{\geq 1}$ . Prove that  $x \mapsto \frac{1}{x^p}$  is integrable on  $[1, \infty)$  iff  $p \geq 2$ .**

Proof: For  $p \geq 2$ ,

$$\lim_{n \rightarrow \infty} \int_1^n \frac{1}{x^p} dx = \lim_{n \rightarrow \infty} \frac{1}{1-p} x^{1-p} \Big|_1^n = \frac{1}{1-p}.$$

If  $p = 1$ ,  $\lim_{n \rightarrow \infty} \int_1^n \frac{1}{x} dx = \lim_{n \rightarrow \infty} \log x \Big|_1^n = \infty$ .

**B4) Prove that  $\{S_n(p)\}_{n \geq 1}$  converges iff  $p \geq 2$ . For  $p \geq 2$  let**

$$\zeta(p) = \lim_{n \rightarrow \infty} S_n(p) = \sum_{k=1}^{\infty} \frac{1}{k^p}.$$

Proof: If  $p = 1$ ,  $S_n(p) \geq \int_1^{n+1} \frac{1}{x} dx \rightarrow \infty$ .

For  $p \geq 2$ ,  $S_n(p) \leq S_{n+1}(p)$ , and  $S_n(p) \leq 1 + \int_1^n \frac{1}{x^p} dx \leq 1 + \int_1^{\infty} \frac{1}{x^p} dx$ . Hence  $\lim_{n \rightarrow \infty} S_n(p)$  exists.

**Part 2: Calculate  $\zeta(2)$**

(We can also use Bernolli numbers and the Taylor expansion of  $\tan x$ ).

Let  $h(t) = \frac{t^2}{2\pi} - t$ ,  $\varphi : [0, \pi] \rightarrow \mathbb{R}$  :

$$\varphi(x) = \begin{cases} -1, & x = 0; \\ \frac{h(x)}{2 \sin(\frac{x}{2})}, & 0 < x \leq \pi. \end{cases}$$

**B5) Prove that  $\varphi \in C^1([0, \pi])$ .**

Proof:

$$\lim_{x \rightarrow 0} \frac{h(x)}{2 \sin(\frac{x}{2})} = \lim_{x \rightarrow 0} \frac{-x + o(x)}{2 \sin(\frac{x}{2})} = -1 = \varphi(0).$$

Hence  $\varphi \in C^1([0, \pi])$ .

**B6) For all  $k \geq 1$ , calculate**

$$\int_0^{\pi} h(x) \cos(kx) dx.$$

Solution:

$$\begin{aligned} \int_0^{\pi} \left( \frac{x^2}{2\pi} - x \right) \cos(kx) dx &= \frac{1}{k} \int_0^{\pi} \left( \frac{x^2}{2\pi} - x \right) d \sin(kx) \\ &= -\frac{1}{k} \int_0^{\pi} \sin(kx) \left( \frac{x}{\pi} - 1 \right) dx \\ &= \frac{1}{k^2} \int_0^{\pi} \left( \frac{x}{\pi} - 1 \right) d \cos(kx) \\ &= \frac{1}{k^2} - \frac{1}{\pi k^2} \int_0^{\pi} \cos(kx) dx = \frac{1}{k^2}. \end{aligned}$$

**B7) Prove that there is a constant  $\lambda$ , such that for any  $x \in (0, \pi)$ ,**

$$\sum_{k=1}^n \cos(kx) = \frac{\sin\left(n + \frac{1}{2}\right)x}{2 \sin\left(\frac{x}{2}\right)} - \lambda.$$

Proof: Note that  $2 \cos(kx) \sin(\frac{x}{2}) = \sin(k + 1/2)x - \sin(k - 1/2)x$ , hence

$$\sum_{k=1}^n \cos(kx) \cdot 2 \sin \frac{x}{2} = \sin \left( n + \frac{1}{2} \right) x - \sin \frac{x}{2}, \lambda = \frac{1}{2}.$$



**B8) Prove that for any  $\psi \in C^1([0, \pi])$ ,**

$$\lim_{n \rightarrow \infty} \int_0^\pi \psi(x) \sin(n + 1/2)x \, dx = 0.$$

Proof: Since  $\sin(n + 1/2)x = c_1 \sin nx + c_2 \cos nx$ , where  $c_1, c_2$  are constant, it suffices to show that

$$\lim_{n \rightarrow \infty} \int_0^\pi \psi(x) \sin(2nx) \, dx = \lim_{n \rightarrow \infty} \int_0^\pi \psi(x) \cos(2nx) \, dx = 0.$$

Note that

$$\begin{aligned} \int_0^\pi \psi(x) \sin(2nx) \, dx &= \sum_{k=1}^n \int_{(k-1)\pi/n}^{k\pi/n} \psi(x) \sin(2nx) \, dx \\ &= \sum_{k=1}^n \frac{1}{2n} \int_0^{2\pi} \psi\left(\frac{x}{2n} + \frac{(k-1)\pi}{n}\right) \sin x \, dx \\ \left(t = \frac{(k-1)\pi}{n}\right) &\leq \sum_{k=1}^n \frac{\pi}{n} \sup_{x \leq \pi} \left| \psi\left(\frac{x+\pi}{2n} + t\right) - \psi\left(\frac{x}{2n} + t\right) \right| \\ &\leq \pi \sup_{0 \leq x \leq \pi - \pi/2n} \left| \psi\left(x + \frac{\pi}{2n}\right) - \psi(x) \right| \rightarrow 0. \end{aligned}$$

since  $\psi$  is uniformly continuous on  $[0, \pi]$ .

**B9) Prove that  $\zeta(2) = \frac{\pi^2}{6}$ .**

Proof:

$$\begin{aligned} \zeta(2) &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{k^2} = \lim_{n \rightarrow \infty} \sum_{k=1}^n \int_0^\pi h(x) \cos(kx) \, dx \\ &= \lim_{n \rightarrow \infty} \int_0^\pi \psi(x) \sin(n + 1/2)x - \frac{1}{2} \left( \frac{x^2}{2\pi} - x \right) \, dx \\ \text{(B8)} &= \frac{1}{2} \int_0^\pi \left( x - \frac{x^2}{2\pi} \right) \, dx = \frac{\pi^2}{6}. \end{aligned}$$

**Part 3:  $\zeta(2)$  is irrational**

Otherwise assume  $\pi^2 = \frac{a}{b}$  where  $a, b \in \mathbb{Z}$ .

**B10) Define a sequence of polynomials  $f_n(x) = \frac{x^n(1-x)^n}{n!}$ , where  $n \in \mathbb{Z}_{\geq 1}$ . Prove that for any  $k \in \mathbb{Z}$ ,  $f_n^{(k)}(0), f_n^{(k)}(1) \in \mathbb{Z}$ .**

Proof: If  $k \leq n-1$ , then  $f_n^{(k)}(0) = f_n^{(k)}(1) = 0$ . If  $k \geq n$ , then

$$\text{if } x^n(1-x)^n = \sum_{k=n}^{2n} c_k x^k, \text{ then } f_n^{(k)}(x) = \sum_{m=n}^{2n} c_k \binom{m}{k} x^{m-k} \in \mathbb{Z}[x].$$

Hence  $f_n^{(k)}(0), f_n^{(k)}(1) \in \mathbb{Z}$ .

**B11) Define the sequence**

$$F_n(x) = b^n(\pi^{2n}f_n(x) - \pi^{2n-2}f_n^{(2)}(x) + \cdots + (-1)^n f_n^{(2n)}(x)).$$

Prove that  $F_n(0), F_n(1) \in \mathbb{Z}$ .Proof: For  $0 \leq k \leq n$ ,  $b^n \pi^{2n-2k} f_n^{(2k)}(x) \in \mathbb{Z}$ , when  $x \in \{0, 1\}$ .**B12) For  $n \geq 1$ , define  $\{g_n\}_{n \geq 1}, \{A_n\}_{n \geq 1}$  as below:**

$$g_n(x) = F'_n(x) \sin(\pi x) - \pi F_n(x) \cos(\pi x), \quad A_n = \pi \int_0^1 a^n f_n(x) \sin(\pi x) dx.$$

Prove that  $A_n \in \mathbb{Z}$  and  $g'_n = \pi^2 a^n f_n(x) \sin(\pi x)$ .

Proof: Note that

$$\begin{aligned} g'_n(x) &= b^n \pi^{2n} \sum_{k=0}^n \left( f_n^{(2k)}(x) \sin(\pi x) - \pi f_n^{(2k+1)}(x) \cos(\pi x) \right)' (-\pi^2)^k \\ &= b^n \pi^{2n+2} f_n(x) \sin(\pi x). \end{aligned}$$

And

$$\begin{aligned} A_n &= \frac{1}{\pi} \int_0^1 dg_n(x) = \frac{1}{\pi} (g_n(1) - g_n(0)) \\ &= F_n(0) + F_n(1) \in \mathbb{Z}. \end{aligned}$$

**B13) Prove that there exists  $n \in \mathbb{Z}$  such that for all  $x \in [0, 1]$ ,  $a^n f_n(x) < 1/2$ .**

Proof:

$$f_n(x) = \frac{1}{n!} (x(1-x))^n \leq \frac{1}{n! 4^n} \rightarrow 0.$$

**B14) Prove that there exists  $n \in \mathbb{Z}$  such that  $A_n \in (0, 1)$ , leading to contradiction.**Proof:  $f_n, \sin(\pi x) \geq 0$ , when  $x \in [0, 1]$ , hence  $A_n > 0$ .Take  $n$  such that  $a^n f_n < 1/2$  then  $A_n < \frac{\pi}{2} \int_0^1 \sin(\pi x) dx = 1$ .Therefore  $A_n \in (0, 1)$ , contradicting with  $A_n \in \mathbb{Z}$ .

### 10.3 PSC: Calculation of Integrals

 $a \neq 0, b \neq 0$ 

(1)  $\int_0^\pi \sin^3 x dx$

$$\int_0^\pi \sin^3(x) dx = -2 \int_0^{\pi/2} \sin^2(x) d \cos(x) = 2 \int_0^1 (1-x^2) dx = \frac{4}{3}.$$

(2)  $\int_{-\pi}^\pi x^2 \cos x dx$

$$\int_{-\pi}^\pi x^2 \cos(x) dx = (x^2 - 2) \sin(x) + 2x \cos(x) \Big|_{-\pi}^\pi = -4\pi.$$

(3)  $\int_0^1 \frac{x}{1+\sqrt{1+x}} dx$

$$\int_0^1 \frac{x}{1+\sqrt{1+x}} dx = \int_0^1 \sqrt{1+x} - 1 dx = \frac{2}{3}(1+x)^{3/2} - x \Big|_0^1 = \frac{4\sqrt{2}-5}{3}.$$

(4)  $\int_0^{\sqrt{3}} x \arctan x dx$

$$\begin{aligned} \int_0^{\sqrt{3}} x \arctan x dx &= \frac{1}{2} \int_0^{\sqrt{3}} \arctan x dx^2 \\ &= \frac{1}{2} x^2 \arctan x \Big|_0^{\sqrt{3}} - \frac{1}{2} \int_0^{\sqrt{3}} \frac{x^2}{1+x^2} dx \\ &= \frac{3}{2} \frac{\pi}{3} - \frac{\sqrt{3}}{2} + \frac{1}{2} \arctan \sqrt{3} = \frac{2\pi}{3} - \frac{\sqrt{3}}{2}. \end{aligned}$$

(5)  $\int_{-1}^0 (2x+1)\sqrt{1-x-x^2} dx$

$$\int_{-1}^0 (2x+1)\sqrt{1-x-x^2} dx = \int_{-1}^1 \frac{y}{4} \sqrt{5-y^2} dy = 0$$

(6)  $\int_{\frac{1}{e}}^e |\log x| dx$

$$\begin{aligned} \int_{\frac{1}{e}}^e |\log x| dx &= \int_1^e \log x dx + \int_1^{1/e} \log x dx \\ &= (x \log x - x) \Big|_1^e + (x \log x - x) \Big|_1^{1/e} = 2 - \frac{2}{e} \end{aligned}$$

(7)  $\int_0^a x^2 \sqrt{a^2 - x^2} dx$

$$\int_0^a x^2 \sqrt{a^2 - x^2} dx = a^4 \int_0^{\pi/2} \sin^2 t \cos^2 t dt = \frac{a^4 \pi}{16}$$

(8)  $\int_0^{\log 2} \sqrt{e^x - 1} dx$

$$\begin{aligned} \int_0^{\log 2} \sqrt{e^x - 1} dx &= \int_1^2 \frac{\sqrt{y-1}}{y} dy = \int_0^1 \frac{\sqrt{x}}{1+x} dx \\ &= \int_0^{\pi/4} 2 \tan^2 \theta d\theta = 2 - \frac{\pi}{2}. \end{aligned}$$

(9)  $\int_1^2 x^{100} \log x \, dx$

$$\begin{aligned} \int_1^2 x^{100} \log x \, dx &= \int_1^2 \log x \, d\frac{x^{101}}{101} = \frac{2^{101} \log 2}{101} - \int_1^2 \frac{x^{100}}{101} \, dx \\ &= \frac{2^{101} \log 2}{101} - \frac{2^{101} - 1}{101^2}. \end{aligned}$$

(10)  $\int_0^a \log(x + \sqrt{x^2 + a^2}) \, dx$

$$\begin{aligned} \int_0^a \log(x + \sqrt{x^2 + a^2}) \, dx &= \\ \int_0^a \log(x + \sqrt{x^2 + a^2}) \, dx &= a \int_0^1 \log a + \log(t + \sqrt{t^2 + 1}) \, dt \\ &= a \log a + a \int_0^1 \log(t + \sqrt{t^2 + 1}) \, dt \\ &= a \log a + (\log(1 + \sqrt{2}) + \sqrt{2} - 1)a. \\ \int_0^1 \log(x + \sqrt{x^2 + 1}) \, dx &= x \log(x + \sqrt{x^2 + 1}) \Big|_0^1 - \int_0^1 \frac{x}{\sqrt{1 + x^2}} \, dx \\ (x = \tan \theta) &= \log(1 + \sqrt{2}) + \int_0^{\pi/4} \frac{1}{\cos^2 \theta} \, d \cos \theta \\ &= \log(1 + \sqrt{2}) + \sqrt{2} - 1. \end{aligned}$$

(11)  $\int_0^{\pi/2} \frac{\cos x \sin x}{a^2 \sin^2 x + b^2 \cos^2 x} \, dx$

$$\begin{aligned} \int_0^{\pi/2} \frac{\cos x \sin x}{a^2 \sin^2 x + b^2 \cos^2 x} \, dx &= \int_0^{\pi/2} \frac{\sin 2x}{a^2 + b^2 + (b^2 - a^2) \cos 2x} \, dx \\ &= \frac{1}{2} \int_{-1}^1 \frac{1}{(a^2 + b^2) + (b^2 - a^2)t} \, dt \\ &= \frac{1}{2(a^2 - b^2)} \log \left( \frac{a^2}{b^2} \right). \end{aligned}$$

(12)  $\int_0^{\pi/2} \frac{\cos^2 x}{\cos x + \sin x} \, dx$

$$\begin{aligned} \int_0^{\pi/2} \frac{\cos^2 x}{\cos x + \sin x} \, dx &= \int_0^{\pi/2} \frac{\sin^2 x}{\cos x + \sin x} \, dx \\ &= \frac{1}{2} \int_0^{\pi/2} \frac{1}{\cos x + \sin x} \, dx \\ &= \int_{\pi/4}^{\pi/2} \frac{1}{\sqrt{2} \sin x} \, dx \\ &= -\frac{\log \tan \left( \frac{\pi}{8} \right)}{\sqrt{2}}. \end{aligned}$$

$$(13) \int_0^{\pi/2} \frac{\sin^2 x}{\cos x + \sin x} dx$$

See (12)

$$(14) \int_0^{\pi/4} \log(1 + \tan x) dx$$

$$\begin{aligned} \int_0^{\pi/4} \log(1 + \tan x) dx &= \int_0^{\pi/4} \log \frac{\sin x + \cos x}{\cos x} dx \\ &= \int_0^{\pi/4} \log \frac{\sqrt{2} \sin(x + \pi/4)}{\cos x} dx \\ &= \frac{\pi}{8} \log 2. \end{aligned}$$

$$(15) \int_0^4 \frac{|x-1|}{|x-2|+|x-3|} dx$$

$$\begin{aligned} \int_0^1 \frac{|x-1|}{|x-2|+|x-3|} dx &= \int_0^1 \frac{1-x}{5-2x} dx = \frac{1}{2} - \frac{3}{4} \log \frac{5}{3}, \\ \int_1^2 \frac{|x-1|}{|x-2|+|x-3|} dx &= \int_0^1 \frac{x}{3-2x} dx = -\frac{1}{2} + \frac{3}{4} \log 3, \\ \int_2^3 \frac{|x-1|}{|x-2|+|x-3|} dx &= \int_0^1 (x+1) dx = \frac{3}{2}, \\ \int_3^4 \frac{|x-1|}{|x-2|+|x-3|} dx &= \int_0^1 \frac{x+2}{2x+1} dx = \frac{1}{2} + \frac{3}{4} \log 3, \\ \Rightarrow \int_0^4 \frac{|x-1|}{|x-2|+|x-3|} dx &= 2 + \frac{3}{4} \log \frac{27}{5} \end{aligned}$$

$$(16) \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx$$

$$\begin{aligned} \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx &= \int_0^{\pi} x d \arctan \cos x \\ &= -x \arctan \cos x \Big|_0^{\pi} + \int_0^{\pi} \arctan \cos x dx \\ &= \frac{\pi^2}{4} + 0 = \frac{\pi^2}{4}. \end{aligned}$$

$$(17) \int_0^{\pi/4} \frac{\sin x}{\sin x + \cos x} dx$$

$$\begin{aligned} \int_0^{\pi/4} \frac{\sin x}{\sin x + \cos x} dx &= \frac{1}{2} \int_{\pi/4}^{\pi/2} \frac{\sin t - \cos t}{\sin t} dt \\ &= \frac{\pi}{8} - \frac{\log 2}{4}. \end{aligned}$$

(18)  $\int_0^{\pi/2} \frac{\sin 2019x}{\sin x} dx$

$$\int_0^{\pi/2} \frac{\sin(2n+1)x}{\sin x} dx = \int_0^{\pi/2} 1 + \sum_{k=1}^n \cos(2kx) dx = \frac{\pi}{2}.$$

(19)  $\int_2^4 \frac{\log \sqrt{9-x}}{\log \sqrt{9-x} + \log \sqrt{x+3}} dx$

$$\int_2^4 \frac{\log \sqrt{9-x}}{\log \sqrt{9-x} + \log \sqrt{x+3}} dx = \int_{-1}^1 \frac{\log \sqrt{6+t}}{\log \sqrt{6+t} + \log \sqrt{6-t}} dt = 1.$$

(20)  $\int_0^1 \frac{1}{\sqrt{1+x^2} + \sqrt{1-x^2}} dx$

$$\begin{aligned} \int_0^1 \frac{dx}{\sqrt{1+x^2} + \sqrt{1-x^2}} &= \int_0^1 \frac{1}{2} (\sqrt{1+x^2} - \sqrt{1-x^2}) dx \\ &= -\frac{\pi}{8} + \frac{\sqrt{2}}{4} + \frac{1}{8} \log \frac{\sqrt{2}+1}{\sqrt{2}-1}. \end{aligned}$$

(21)  $\int_0^1 \sqrt{x + \sqrt{x+1}} dx$

$$\begin{aligned} \int_0^1 \sqrt{x + \sqrt{x+1}} dx &= \int_1^{1+\sqrt{2}} \sqrt{y} d \frac{2y+1-\sqrt{4y+5}}{2} \\ &= \int_1^{1+\sqrt{2}} \sqrt{y} - \frac{\sqrt{y}}{\sqrt{4y+5}} dy \\ \left( y = \frac{z^2-5}{4} \right) &= \frac{2}{3} y^{3/2} \Big|_1^{1+\sqrt{2}} - \int_3^{1+2\sqrt{2}} \frac{\sqrt{z^2-5}}{4} dz \\ &= \frac{2}{3} ((1+\sqrt{2})^{3/2} - 1) - \frac{3\sqrt{2}-1}{8} + \frac{5}{32} \log \frac{3+\sqrt{2}}{5}. \end{aligned}$$

(22)  $\int_{-1}^1 \frac{\sin \sin \sin x}{x^{800} + 1} dx$

$$\int_{-1}^1 \frac{\sin \sin \sin x}{x^{800} + 1} dx = 0. (\text{by symmetry})$$

## 11 Homework 11: Density of Sum of Squares

### 11.1 PSA: Riemann Integral

**A1)**  $f \in C([a, b])$ ,  $g \in \mathcal{R}([a, b])$ , where  $g$  is positive. Prove that there exists  $\xi \in (a, b)$ , such that

$$\int_a^b fg = f(\xi) \int_a^b g.$$

Proof: Since  $g$  is positive on  $[a, b]$ ,

$$\inf_{x \in [a, b]} f(x) \int_a^b g \leq \int_a^b fg \leq \sup_{x \in [a, b]} f(x) \int_a^b g.$$

By  $f \in C([a, b])$ , there exists such an  $\xi \in (a, b)$ .

**A2) Prove without using Lebesgue theorem: if  $f$  is monotonously increasing on  $[a, b]$ , then  $f \in \mathcal{R}([a, b])$ .**

Proof: For any  $\varepsilon > 0$  let  $n = [1/\varepsilon] + 1$ , and

$$\mathcal{C} = \left\{ x_k = a + (b - a) \frac{k}{n} : k = 0, 1, \dots, n \right\}.$$

Then

$$g(x) = \max_{x_k \leq x} \{f(x_k)\} \leq f, h(x) = \min_{x_k \geq x} \{f(x_k)\} \geq f.$$

and both are monotonous simple functions.

Therefore

$$\overline{\int_a^b f} - \underline{\int_a^b f} \leq \overline{S}(f; \mathcal{C}) - \underline{S}(f; \mathcal{C}) = \frac{1}{n}(f(b) - f(a)) \rightarrow 0.$$

Hence  $f$  is Riemann integrable.

**A3) Prove that  $1_{\mathbb{Q}}$  is not Riemann integrable on  $[0, 1]$ .**

Proof: Let  $\varepsilon = \frac{1}{2}$ . For any  $\mathcal{C} = \{0 = x_0 < \dots < x_n = 1\}$ ,  $\omega(x_{k-1}, x_k) = 1$ , hence

$$\sum_{k=1}^n \omega(x_{k-1}, x_k)(x_k - x_{k-1}) = 1 > \varepsilon.$$

Therefore  $1_{\mathbb{Q}}$  is not Riemann integrable.

**A4) Prove that if  $f \in \mathcal{R}([a, b])$ , then  $|f|^p \in \mathcal{R}([a, b])$ , where  $p \geq 0$ .**

Proof: Since  $x \mapsto |x|^p$  is continuous,  $|f|^p$  is continuous as  $x$  whenever  $f$  is continuous at  $x$ . Hence

$$f \in \mathcal{R}([a, b]) \implies |f|^p \in \mathcal{R}([a, b]).$$

**A5) Prove Hölder's Inequality: if  $f, g \in \mathcal{R}([a, b])$ ,  $p, q > 0$ ,  $1/p + 1/q = 1$ , then**

$$\left| \int_a^b fg \right| \leq \left( \int_a^b |f|^p \right)^{1/p} \left( \int_a^b |g|^q \right)^{1/q}.$$

Proof: By A4) the functions are integrable. We can assume that

$$\int_a^b |f|^p = \int_a^b |g|^q = 1.$$

Then by Young's inequality,

$$\left| \int_a^b fg \right| \leq \int_a^b |f| \cdot |g| \leq \int_a^b \frac{1}{p} |f|^p + \frac{1}{q} |g|^q = \frac{1}{p} + \frac{1}{q} = 1.$$

**A6) Prove Minkowski's inequality: if  $f, g \in \mathcal{R}([a, b])$ ,  $p \geq 1$ , then**

$$\left( \int_a^b |f + g|^p \right)^{1/p} \leq \left( \int_a^b |f|^p \right)^{1/p} + \left( \int_a^b |g|^p \right)^{1/p}.$$

Proof: Note that if  $1/p + 1/q = 1$ , then

$$\begin{aligned} \int_a^b |f + g|^p &= \int_a^b |f| \cdot |f + g|^{1-p} + \int_a^b |g| \cdot |f + g|^{1-p} \\ &\leq \left( \left( \int_a^b |f|^p \right)^{1/p} + \left( \int_a^b |g|^p \right)^{1/p} \right) \left( \int_a^b |f + g|^{(1-p)q} \right)^{1/q} \end{aligned}$$

Hence

$$\left( \int_a^b |f + g|^p \right)^{1/p} \leq \left( \int_a^b |f|^p \right)^{1/p} + \left( \int_a^b |g|^p \right)^{1/p}.$$

The equality holds, when  $|f|/|f + g|^{1-p}$ ,  $|g|/|f + g|^{1-p}$  are both constant, which is equivalent to  $|f|/|g|$  is constant.

## 11.2 PSB: Convex Functions

**B1) Assume  $f \in \mathcal{R}([a, b])$  and  $f$  is convex, prove that**

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) \, dx \leq \frac{f(a) + f(b)}{2}.$$

Proof: Note that  $f\left(\frac{a+b}{2}\right) \leq \frac{f(x) + f(a+b-x)}{2} \leq \frac{f(a) + f(b)}{2}$ , and

$$\int_a^b f(x) \, dx = \int_a^b \frac{f(x) + f(a+b-x)}{2} \, dx.$$

Hence

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) \, dx \leq \frac{f(a) + f(b)}{2}.$$

**B2) Assume  $f$  is twice differentiable on  $[a, b]$  and for any  $x$ ,  $f''(x) > 0$ ,  $f(x) \leq 0$ . Prove that for any  $x$ ,**

$$f(x) \geq \frac{2}{b-a} \int_a^b f(y) \, dy.$$

Proof: For any  $x \leq y \leq b$ ,

$$f(y) \leq \frac{b-y}{b-x} f(x) + \frac{y-x}{b-x} f(b) \leq \frac{b-y}{b-x} f(x),$$

hence

$$\int_x^b f(y) \, dy \leq f(x) \int_x^b \frac{b-y}{b-x} \, dy = \frac{b-x}{2} f(x).$$



Likewise,

$$\int_a^x f(y) \, dy \leq f(x) \int_a^x \frac{y-a}{x-a} \, dy = \frac{x-a}{2} f(x).$$

Therefore

$$f(x) \geq \frac{2}{b-a} \int_a^b f(y) \, dy.$$

**B3) Assume  $f$  is twice differentiable on  $\mathbb{R}$  and  $f''(x) \geq 0$ ,  $\varphi \in C([a, b])$ . Prove that**

$$\frac{1}{b-a} \int_a^b (f \circ \varphi)(t) \, dt \geq f \left( \frac{1}{b-a} \int_a^b \varphi(t) \, dt \right).$$

Proof: We prove the proposition for any convex function  $f$  and  $\varphi$  on the set  $X$ .

Let

$$\langle g \rangle = \frac{1}{\mu(X)} \int_X g \, d\mu.$$

Then since  $f$  is convex, there is a constant  $K$  such that  $f(y) - f(\langle \varphi \rangle) \geq K(y - \langle \varphi \rangle)$ . Hence

$$\begin{aligned} \langle f(\varphi) \rangle &= \frac{1}{\mu(X)} \int_X f(\varphi(t)) \, d\mu \\ &\geq \frac{1}{\mu(X)} \int_X f(\langle \varphi \rangle) \, d\mu + \frac{1}{\mu(X)} \int_X K(\varphi(t) - \langle \varphi \rangle) \, d\mu \\ &= f(\langle \varphi \rangle). \end{aligned}$$

**B4) Assume  $f \in C([a, b])$  and for any  $x$ ,  $f(x) > 0$ . Prove that**

$$\log \left( \frac{1}{b-a} \int_a^b f \right) \geq \frac{1}{b-a} \int_a^b \log f.$$

Proof: Since  $-\log x$  is convex, we can use B3).

**B5) Prove that if  $f$  is convex on  $\mathbb{R}$ ,  $\varphi \in C([0, 1])$ , then**

$$f \left( \int_0^1 \varphi \right) \leq \int_0^1 f \circ \varphi.$$

Proof: A special case of what we proved in B3).

### 11.3 PSC: Integrals and Derivatives

**C1) Assume  $f \in C^1([0, 2])$ ,  $|f'| \leq 1$ ,  $f(0) = f(2) = 1$ . Prove that**

$$1 \leq \int_0^2 f \leq 3.$$

Proof: Note that for  $0 \leq x \leq 1$ ,

$$|f(x) - 1| = x|f'(\xi)| \leq x.$$

and for  $1 \leq x \leq 2$ ,

$$|f(x) - 1| = (2-x)|f'(\xi)| \leq 2-x.$$

Hence

$$\int_0^2 |f(x) - 1| dx \leq \int_0^1 x dx + \int_1^2 (2 - x) dx = 1.$$

**C2) Assume that  $f \in C^2([0, 1])$ . Prove that  $\exists \xi \in [0, 1]$ , such that**

$$\int_0^1 f(x) dx = f\left(\frac{1}{2}\right) + \frac{1}{24}f''(\xi).$$

Proof: Let  $g(x) = f(x) + f(1 - x)$ , then

$$\begin{aligned} \int_0^1 f(x) dx - f\left(\frac{1}{2}\right) &= \int_0^{1/2} g(x) - 2f\left(\frac{1}{2}\right) dx \\ (\text{integration by parts}) &= - \int_0^{1/2} x g'(x) dx = -\frac{1}{2} \int_0^{1/2} g'(x) dx^2 \\ (\text{integration by parts}) &= \frac{1}{2} \int_0^{1/2} x^2 g''(x) dx. \end{aligned}$$

Note that  $g'' \in C([0, 1])$  hence by A1),  $\exists \eta \in (0, \frac{1}{2})$ ,

$$\int_0^1 f(x) dx - f\left(\frac{1}{2}\right) = g''(\eta) \frac{1}{2} \int_0^{1/2} x^2 dx = \frac{1}{48} g''(\eta).$$

Since  $f'' \in C([0, 1])$ , there exists  $\xi \in (\eta, 1 - \eta)$ , such that

$$f''(\xi) = \frac{f''(\eta) + f''(1 - \eta)}{2} = \frac{g''(\eta)}{2}.$$

Therefore

$$\int_0^1 f(x) dx = f\left(\frac{1}{2}\right) + \frac{1}{24}f''(\xi).$$

**C3) Assume  $f \in C^1([0, 1])$ . Prove that**

$$\max_{x \in [a, b]} |f(x)| \leq \frac{1}{b - a} \left| \int_a^b f(x) dx \right| + \int_a^b |f'(x)| dx.$$

Proof: For any  $t \in [a, b]$ ,

$$(b - a)|f(t)| \leq \left| \int_a^b f(x) dx \right| + \left| \int_a^b f(x) - f(t) dx \right|$$

where

$$\begin{aligned} \left| \int_a^b f(x) - f(t) dx \right| &= \left| \int_a^b \left( \int_t^x f'(u) du \right) dx \right| \\ &\leq \int_a^b \int_t^x |f'(u)| du dx \\ &\leq (b - a) \int_a^b |f'(u)| du. \end{aligned}$$

**C4) Suppose  $f \in C([0, 1])$  and for any  $g \in C([0, 1])$ ,  $g(0) = g(1) = 0$ , we have**

$$\int_0^1 f(x)g(x) \, dx = 0.$$

Prove that  $f(x) \equiv 0$ .

Proof: Otherwise assume  $f(t) > 0$  for some  $t \in (0, 1)$ , then there exists an  $\varepsilon > 0$  such that  $(t - \varepsilon, t + \varepsilon) \subset [0, 1]$  and  $\forall x \in (t - \varepsilon, t + \varepsilon), f(x) > f(t)/2$ .

Let

$$g(x) = \begin{cases} 0, & x \notin (t - \varepsilon, t + \varepsilon), \\ 1 - \frac{|x - t|}{\varepsilon}, & x \in (t - \varepsilon, t + \varepsilon). \end{cases}$$

Then

$$\int_0^1 f(x)g(x) \, dx > \int_{t-\varepsilon}^{t+\varepsilon} \frac{f(t)}{2} g(x) \, dx > 0,$$

leading to contradiction. Hence  $f(x) \equiv 0$ .

**C5) Suppose  $f \in C([0, 1])$  and for any  $n \in \mathbb{Z}_{\geq 0}$ ,**

$$\int_0^1 f(x)x^n \, dx = 0.$$

Prove that  $f(x) \equiv 0$ .

Proof: Otherwise,  $\int_0^1 f^2 > 0$ . By Stone-Weierstrass theorem, for any  $\varepsilon > 0$ , there is a polynomial  $P$  such that  $\sup_{x \in [0, 1]} |f(x) - P(x)| < \varepsilon$ . Hence

$$0 = \int_0^1 f(x)P(x) \, dx = \int_0^1 f^2 - \int_0^1 f(x)(f(x) - P(x)) \, dx \geq \int_0^1 f^2 - \sup_{x \in [0, 1]} |f(x)|\varepsilon > 0$$

when  $\varepsilon \rightarrow 0$ , leading to contradiction.

**C6) (Gronwall's Inequality) Suppose  $\varphi \in C([0, T])$  and for any  $t \in [0, T]$ ,  $|\varphi(t)| \leq M + k \int_0^t |\varphi(s)| \, ds$ , where  $M, k$  are positive real numbers. Prove that  $\forall t \in [0, T]$ ,  $|\varphi(t)| \leq Me^{kt}$ .**

Proof: Let

$$f : \left[0, \frac{T}{k}\right] \rightarrow \mathbb{R}, t \mapsto \frac{e^{-t}|\varphi(t/k)|}{M},$$

then for any  $t \in [0, T/k]$ ,

$$f(t) \leq e^{-t} + e^{-t} \int_0^t f(s)e^s \, ds.$$

Let  $f(t) = \sup_{s \in [0, T/k]} \{f(s)\}$  then

$$f(t) \leq e^{-t} + e^{-t} \int_0^t f(t)e^s \, dx = e^{-t} + f(t)(1 - e^{-t}).$$

Hence  $f(s) \leq f(t) \leq 1, \implies |\varphi(t)| \leq Me^{kt}$ .

**C7) Assume  $a, b > 0$ ,  $f \in C([-a, b])$ . If for any  $x \in (-a, b)$ ,  $f(x) > 0$  and  $\int_{-a}^b xf(x) \, dx = 0$ . Prove that**

$$\int_{-a}^b x^2 f(x) \, dx \leq ab \int_{-a}^b f(x) \, dx.$$

Proof: Note that

$$\int_{-a}^b (x+a)(x-b)f(x) \, dx \leq 0.$$

Combined with  $\int_{-a}^b xf(x) \, dx = 0$  we get

$$\int_{-a}^b x^2 f(x) \, dx \leq ab \int_{-a}^b f(x) \, dx.$$

**C8) Assume  $f \in C([-1, 1])$ . Prove that**

$$\lim_{\lambda \rightarrow 0^+} \int_{-1}^1 \frac{\lambda}{\lambda^2 + x^2} f(x) \, dx = \pi f(0).$$

Proof: Let  $M = \sup_{|x| \leq 1} |f(x)|$  and

$$g(x) = \frac{\lambda}{\lambda^2 + x^2},$$

then ( $g$  is sort of a good kernel)

$$\int_{-1}^1 g(x) \, dx = 2 \arctan \frac{1}{\lambda}.$$

Hence

$$\begin{aligned} & \left| \int_{-1}^1 f(x)g(x) \, dx - \pi f(0) \right| \\ & \leq \left| \pi - 2 \arctan \frac{1}{\lambda} \right| f(0) + \int_{-\varepsilon}^{\varepsilon} |f(x) - f(0)|g(x) \, dx + \int_{\varepsilon \leq |x| \leq 1} M g(x) \, dx \\ & \leq \left| \pi - 2 \arctan \frac{1}{\lambda} \right| f(0) + \sup_{|x| \leq \varepsilon} |f(x) - f(0)| \pi + 2M \left| \arctan \frac{1}{\lambda} - \arctan \frac{\varepsilon}{\lambda} \right| \\ & \rightarrow 0 \end{aligned}$$

since

$$\arctan \frac{1}{\lambda} - \arctan \frac{\varepsilon}{\lambda} = \arctan \frac{\lambda(1-\varepsilon)}{\lambda^2 + \varepsilon} \rightarrow 0, \text{ when } \lambda \rightarrow 0^+.$$

and  $\sup_{|x| \leq \varepsilon} |f(x) - f(0)| \rightarrow 0$  when  $\varepsilon \rightarrow 0$ .

**C9) Assume  $f$  is differentiable on  $[1, \infty)$  and both  $\int_1^\infty f(x) \, dx$  and  $\int_1^\infty f'(x) \, dx$  converges. Prove that**

$$\lim_{x \rightarrow \infty} f(x) = 0.$$

Proof: For any  $\varepsilon > 0$ , there exists  $N > 1$ , such that  $\forall u, v > N$ ,

$$\left| \int_u^v f'(x) \, dx \right| < \varepsilon, \text{ i.e. } |f(u) - f(v)| < \varepsilon$$

Hence for any  $u > N$ , if  $|f(u)| > \varepsilon$ ,

$$\left| \int_u^M f(x) \, dx \right| \geq (M - u)(|f(u) - \varepsilon|) \rightarrow \infty, \text{ as } M \rightarrow \infty,$$

which contradicts the fact that  $\int_1^\infty f(x) \, dx$  converges. Therefore  $|f(u)| < \varepsilon$  for any  $u > N$ , which implies  $\lim_{x \rightarrow \infty} f(x) = 0$ .

**C10) Prove that**

$$\int_0^\infty \frac{\sin^2 x}{x^2} dx = \int_0^\infty \frac{\sin x}{x} dx, \int_0^\infty \frac{\cos x}{1+x} dx = \int_0^\infty \frac{\sin x}{(1+x)^2} dx.$$

Proof:

$$\begin{aligned} \int_0^\infty \frac{\sin^2 x}{x^2} dx &= - \int_0^\infty \sin^2 x d\frac{1}{x} = \int_0^\infty \frac{\sin 2x}{x} dx = \int_0^\infty \frac{\sin x}{x} dx. \\ \int_0^\infty \frac{\cos x}{1+x} dx &= \int_0^\infty \frac{1}{1+x} d\sin x = \int_0^\infty \frac{\sin x}{(1+x)^2} dx. \end{aligned}$$

## 11.4 PSD: Calculation of improper integrals

**D1)**

$$\int_0^1 \log x dx = (x \log x - x) \Big|_0^1 = -1.$$

**D2)**

$$\int_{-\infty}^\infty \frac{1}{1+x^2} dx = \arctan x \Big|_{-\infty}^\infty = \pi.$$

**D3)**

Calculating residues, we get

$$\int_{-\infty}^\infty \frac{dx}{x^4+1} = 2\pi i \cdot (\operatorname{Res}(f; e^{i\pi/4}) + \operatorname{Res}(f; e^{3i\pi/4})) = \frac{\pi}{\sqrt{2}}.$$

Hence

$$\int_0^\infty \frac{dx}{x^4+1} = \frac{\pi}{2\sqrt{2}}.$$

**D4)**

Same as D3)

$$\int_{-\infty}^\infty \frac{1+x^2}{1+x^4} dx = \sqrt{2}\pi.$$

Hence

$$\int_0^\infty \frac{1+x^2}{1+x^4} dx = \frac{\pi}{\sqrt{2}}.$$

**D5)**

$$\int_{-\infty}^0 x e^x dx = \int_{-\infty}^0 x d e^x = - \int_{-\infty}^0 e^x dx = -1.$$

D6)

$$\int_0^\infty e^{-\sqrt{x}} dx = 2 \int_0^\infty y e^{-y} dy = 2 \int_0^\infty e^{-y} dy = 2.$$

D7)

$$\int_0^\infty \frac{dx}{(a^2 + x^2)^{3/2}} = \frac{1}{a^2} \int_0^\infty \frac{dx}{(1 + x^2)^{3/2}} = \frac{1}{2a^2} B\left(\frac{1}{2}, 1\right) = \frac{1}{a^2}.$$

(We can also substitute  $x = a \tan \theta$ ).

D8)

$$\int_2^\infty \frac{dx}{x^2 + x - 2} = \frac{1}{3} \log \frac{x-1}{x+2} \Big|_2^\infty = \frac{\log 3}{3}.$$

D9)

$$\begin{aligned} \int_{-\infty}^\infty \frac{dx}{(x^2 + x + 1)^2} &= \frac{8}{3\sqrt{3}} \int_{-\infty}^\infty \frac{du}{(1 + u^2)^2} \\ (u = \tan \theta) &= \frac{8}{3\sqrt{3}} \int_{-\pi/2}^{\pi/2} \cos^2 \theta d\theta = \frac{4\sqrt{3}\pi}{9}. \end{aligned}$$

D10)

$$\int_{-1}^1 \frac{dx}{\sqrt{1-x^2}} = \int_{-\pi/2}^{\pi/2} 1 d\theta = \pi.$$

D11)

$$\int_{-1}^1 \frac{\arcsin x}{\sqrt{1-x^2}} dx = \int_{-\pi/2}^{\pi/2} \theta d\theta = 0.$$

D12)

Let  $\gamma$  be the unit circle, then

$$\begin{aligned} \int_{-1}^1 \frac{dx}{(2-x)^2 \sqrt{1-x^2}} &= \int_{-\pi/2}^{\pi/2} \frac{d\theta}{(2-\sin \theta)^2} = \frac{1}{2} \int_{-\pi}^{\pi} \frac{d\theta}{(2-\sin \theta)^2} \\ &= \frac{1}{2} \int_{\gamma} -\frac{4}{i} \frac{z dz}{(z^2 - 4iz - 1)^2} \\ &= -4\pi \operatorname{Res} \left( \frac{z}{(z^2 - 4iz - 1)^2}; (2 - \sqrt{3})i \right) \\ &= \frac{2\pi}{3\sqrt{3}}. \end{aligned}$$

D13)

$$\int_0^1 \frac{\arcsin \sqrt{x}}{x(1-x)} dx > \int_{1/4}^1 \frac{\pi}{6} \frac{1}{1-x} dx \text{ which diverges.}$$

D14)

$$\int_0^1 (1-x)^n x^{1/2-1} dx = B\left(n+1, \frac{1}{2}\right) = \frac{\Gamma(n+1)\Gamma(\frac{1}{2})}{\Gamma(n+\frac{3}{2})} = \frac{n!2^{n+1}}{(2n+1)!!}.$$

D15)

$$\int_0^1 \frac{x^n}{\sqrt{1-x^2}} dx = \int_0^{\pi/2} \sin^n x dx = \begin{cases} \frac{(n-1)!!}{n!!}, n \text{ is odd,} \\ \frac{(n-1)!!}{n!!} \cdot \frac{\pi}{2}, n \text{ is even.} \end{cases}$$

D16)

Using integration by parts, and substitute  $x = e^{-y}$ ,

$$\begin{aligned} \int_0^1 x^m (\log x)^n dx &= (-1)^n \int_0^\infty e^{-(m+1)y} y^n dy \\ &= (-1)^n \frac{n!}{(m+1)^n} \int_0^\infty e^{-(m+1)y} dy = \frac{(-1)^n n!}{(m+1)^{n+1}}. \end{aligned}$$

D17)

$$\int_2^\infty \frac{dx}{x(\log x)^p} = \int_{\log 2}^\infty \frac{dy}{y^p} = \frac{(\log 2)^{1-p}}{p-1}.$$

D18)

Substitute  $x = ay$ , then

$$\int_0^\infty \frac{\log x}{x^2 + a^2} dx = \frac{\pi \log a}{2a} + \frac{1}{a} \int_0^\infty \frac{\log y}{1 + y^2} dy = \frac{\pi \log a}{2a}.$$

since by substituting  $y = 1/z$ ,

$$\int_0^\infty \frac{\log y}{1 + y^2} dy = - \int_0^\infty \frac{\log z}{1 + z^2} dz = 0.$$

D19)

$$\int_0^\infty x^n e^{-x} dx = \Gamma(n) = (n-1)!.$$

**D20)**

$$\int_{-\infty}^{\infty} \frac{dx}{(ax^2 + 2bx + c)^n} = \frac{1}{d^n} \sqrt{\frac{d}{a}} \int_{-\infty}^{\infty} \frac{du}{(1+u^2)^n} = \frac{1}{d^n} \sqrt{\frac{d}{a}} \pi \frac{(2n-3)!!}{(2n-2)!!}.$$

where  $d = \frac{ac-b^2}{a}$ **D21)**

$$\int_0^{\infty} x^{2n-1} e^{-x^2} dx = \frac{1}{2} \int_0^{\infty} y^{n-1} e^{-y} dy = \frac{(n-1)!}{2}.$$

**D22)**

The Poisson kernel

$$\begin{aligned} \frac{1-r^2}{1-2r\cos x+r^2} &= \frac{1-r^2}{(1-re^{ix})(1-re^{-ix})} \\ &= (1-r^2) \sum_{n=0}^{\infty} r^n e^{inx} \sum_{m=0}^{\infty} r^m e^{-imx} \\ &= \sum_{k=-\infty}^{\infty} r^{|k|} e^{ikx}. \end{aligned}$$

Hence

$$\int_{-\pi}^{\pi} \frac{1-r^2}{1-2r\cos x+r^2} dx = 2\pi.$$

**D23)**

$$\begin{aligned} \int_0^{\infty} e^{-ax} \cos bx dx &= \frac{1}{b} \int_0^{\infty} e^{-ax} d \sin bx = \frac{a}{b} \int_0^{\infty} e^{-ax} \sin bx dx \\ &= -\frac{a}{b^2} \int_0^{\infty} e^{-ax} d \cos bx = \frac{a}{b^2} - \frac{a^2}{b^2} \int_0^{\infty} e^{-ax} \cos bx dx \\ &= \frac{a}{a^2+b^2}. \end{aligned}$$

**D24)**

Same as (23),

$$\int_0^{\infty} e^{-ax} \sin bx dx = \frac{b}{a^2+b^2}.$$



D25)

$$\begin{aligned}
 \int_1^\infty \frac{dx}{x(x+1)\cdots(x+n)} &= \lim_{N \rightarrow \infty} \int_0^N \sum_{k=0}^n \frac{(-1)^k}{x+k} \binom{n}{k} dx \\
 &= \lim_{N \rightarrow \infty} \sum_{k=0}^n (-1)^k \binom{n}{k} \log \left( \frac{N+k}{k+1} \right) \\
 &= - \sum_{k=0}^n (-1)^k \binom{n}{k} \log(k+1) + \lim_{N \rightarrow \infty} \sum_{k=0}^n (-1)^k \binom{n}{k} \log \left( 1 + \frac{k}{N} \right) \\
 &= - \sum_{k=0}^n (-1)^k \binom{n}{k} \log(k+1).
 \end{aligned}$$

D26)

$$\begin{aligned}
 \int_0^\pi \log \sin x \, dx &= 2 \int_0^{\pi/2} \log \sin x \, dx = 2 \int_0^{\pi/2} \log \cos x \, dx \\
 &= \int_0^{\pi/2} \log \sin 2x - \log 2 \, dx = \frac{1}{2} \int_0^\pi \log \sin x \, dx - \frac{\pi}{2} \log 2 \\
 &= -\pi \log 2.
 \end{aligned}$$

D27)

$$\int_0^\infty e^{-x^2} \, dx = \frac{\sqrt{\pi}}{2}.$$

Note that

$$\max\{0, 1 - x^2\} < e^{-x^2} < \frac{1}{1 + x^2}.$$

Hence

$$\frac{(2n)!!}{(2n+1)!!} < \int_0^\infty e^{-nx^2} \, dx < \frac{(2n-3)!!}{(2n-2)!!} \cdot \frac{\pi}{2}.$$

Therefore

$$\sqrt{n} \frac{(2n)!!}{(2n+1)!!} < \int_0^\infty e^{-x^2} \, dx < \sqrt{n} \frac{(2n-3)!!}{(2n-2)!!} \cdot \frac{\pi}{2}.$$

By Wallis's formula,

$$\int_0^\infty e^{-x^2} \, dx = \frac{\sqrt{\pi}}{2}.$$

## 11.5 PSE: Density of sum of squares

Let  $I = (0, \infty)$ .

**Part 1****E1) Prove that  $e^{-u}/\sqrt{u}$  is integrable on  $I$ , and let  $K = \int_0^\infty e^{-u}/\sqrt{u} du$ .**

Proof:

$$\begin{aligned}\int_1^\infty e^{-u}/\sqrt{u} du &< \int_1^\infty e^{-u} du = \frac{1}{e}. \\ \int_0^1 e^{-u}/\sqrt{u} du &< \int_0^1 u^{-1/2} du = \frac{1}{2}.\end{aligned}$$

**E2) Prove that for any  $x \in I$ ,**

$$F(x) = \int_0^\infty \frac{e^{-u}}{\sqrt{u(u+x)}} du$$

is well-defined.

Proof:

$$F(x) < \int_0^\infty \frac{e^{-u}}{x\sqrt{u}} du \text{ converges.}$$

**E3) Prove that  $F \in C^1(I)$  and calculate  $F'(x)$ .**

Solution: Let  $f(x, u) = \frac{e^{-u}}{\sqrt{u(u+x)}}$ , then  $f$  is uniformly continuous on any closed subinterval of  $I$ , and so is

$$\frac{d}{dx} f(x, u) = -\frac{e^{-u}}{\sqrt{u}(u+x)^2}.$$

Also,

$$\int_0^\infty \frac{d}{dx} f(x, u) du$$

converges uniformly.

Hence  $F$  is continuously differentiable and

$$F'(x) = -\int_0^\infty \frac{e^{-u}}{\sqrt{u}(u+x)^2} du.$$

**E4) Prove that for any  $x \in I$ ,**

$$xF'(x) - \left(x - \frac{1}{2}\right) F(x) = -K.$$

Proof: We show that

$$x \int_0^\infty \frac{e^{-u}}{\sqrt{u}(u+x)^2} du + \left(x - \frac{1}{2}\right) \int_0^\infty \frac{e^{-u}}{\sqrt{u}(u+x)} du = \int_0^\infty \frac{e^{-u}}{\sqrt{u}} du.$$

Note that, by substituting  $u \rightarrow ux$ ,

$$\begin{aligned}x \int_0^\infty \frac{e^{-u}}{\sqrt{u}(u+x)^2} du &= \frac{1}{\sqrt{x}} \int_0^\infty \frac{e^{-ux}}{\sqrt{u}(1+u)^2} du, \\ \left(x - \frac{1}{2}\right) \int_0^\infty \frac{e^{-u}}{\sqrt{u}(u+x)} du &= \left(\sqrt{x} - \frac{1}{2\sqrt{x}}\right) \int_0^\infty \frac{e^{-ux}}{\sqrt{u}(1+u)} du, \\ \int_0^\infty \frac{e^{-u}}{\sqrt{u}} du &= \sqrt{x} \int_0^\infty \frac{e^{-ux}}{\sqrt{u}} du.\end{aligned}$$

Hence it is equivalent to

$$x \int_0^\infty \frac{e^{-ux}}{\sqrt{u}} du = \int_0^\infty \frac{e^{-ux}}{\sqrt{u}(1+u)^2} du + \left(x - \frac{1}{2}\right) \int_0^\infty \frac{e^{-ux}}{\sqrt{u}(1+u)} du.$$

Note that  $d e^{-ux} \sqrt{u} = -e^{-ux} \left(x\sqrt{u} - \frac{1}{2\sqrt{u}}\right) du$ , hence

$$\begin{aligned} & x \int_0^\infty \frac{e^{-ux}}{\sqrt{u}} du - \left(x + \frac{1}{2}\right) \int_0^\infty \frac{e^{-ux}}{\sqrt{u}(1+u)} du \\ &= \int_0^\infty e^{-ux} \left(x\sqrt{u} - \frac{1}{2\sqrt{u}}\right) \frac{du}{1+u} \\ &= - \int_0^\infty \frac{d e^{-ux} \sqrt{u}}{1+u} = - \int_0^\infty e^{-ux} \sqrt{u} \frac{du}{(1+u)^2} \\ &= \int_0^\infty \frac{e^{-ux}}{\sqrt{u}} \frac{du}{(1+u)^2} - \int_0^\infty \frac{e^{-ux}}{\sqrt{u}} \frac{du}{1+u}. \end{aligned}$$

$$(\sqrt{u} = \frac{1}{\sqrt{u}}((1+u) - 1))$$

**E5) Define  $G : I \rightarrow \mathbb{R}, x \mapsto \sqrt{x}e^{-x}F(x)$ . Prove that  $\exists C \in \mathbb{R}$  such that**

$$G(x) = C - K \int_0^x \frac{e^{-t}}{\sqrt{t}} dt.$$

Proof: By B4)

$$G'(x) = \sqrt{x}e^{-x}F'(x) + \left(\frac{1}{2\sqrt{x}} - \sqrt{x}\right)e^{-x}F(x) = -K \frac{e^{-x}}{\sqrt{x}}.$$

Hence let  $C = G(0)$ , then

$$G(x) = C + \int_0^x G'(x) dx = C - K \int_0^x \frac{e^{-t}}{\sqrt{t}} dt.$$

**E6) Calculate the value of  $K$ .**

Solution: Note that when  $x \rightarrow \infty$ ,  $F(x) \rightarrow 0$  hence  $G(x) \rightarrow 0$ . Therefore

$$0 = \lim_{x \rightarrow \infty} G(x) = G(0) - K \int_0^\infty \frac{e^{-t}}{t} dt = G(0) - K^2.$$

Where

$$\begin{aligned} G(0) &= \lim_{x \rightarrow 0^+} \frac{\sqrt{x}}{e^x} \int_0^\infty \frac{e^{-u}}{\sqrt{u}(x+u)} du = \lim_{x \rightarrow 0} \int_0^\infty \frac{e^{-ux}}{\sqrt{u}(1+u)} du \\ &= \int_0^\infty \frac{1}{\sqrt{u}(1+u)} du = \int_0^\infty \frac{2dt}{1+t^2} = \pi. \end{aligned}$$

Hence  $K = \sqrt{\pi}$ .

## Part 2

Define

$$f(x) = \sum_{n=1}^\infty \frac{e^{-nx}}{\sqrt{n}}, g(x) = \sum_{n=0}^\infty \sqrt{n}e^{-nx}.$$

**E7) Prove that  $f, g$  are well-defined on  $I$  and are both continuous on  $I$ .**

Proof: Let  $C = \sup_{x \geq 0} x^3 e^{-x}$ , then

$$\sum_{n=1}^{\infty} \frac{e^{-nx}}{\sqrt{n}} < \sum_{n=0}^{\infty} \sqrt{n} e^{-nx} \leq \sum_{n=1}^{\infty} \frac{C}{(nx)^2 \sqrt{x}} \text{ converges.}$$

On any closed sub-interval of  $I$ , the two series both converge uniformly, and  $e^{-nx}$  is continuous, hence  $f, g$  are both continuous on  $I$ .

**E8) Prove that  $\forall x \in I$ ,**

$$\int_1^{\infty} \frac{e^{-ux}}{\sqrt{u}} du \leq f(x) \leq \int_0^{\infty} \frac{e^{-ux}}{\sqrt{u}} du.$$

Proof: The function  $e^{-ux}/\sqrt{u}$  is monotonously decreasing by  $u$ , hence

$$\begin{aligned} \int_1^N \frac{e^{-ux}}{\sqrt{u}} du &\leq \sum_{n=1}^{N-1} \frac{e^{-nx}}{\sqrt{n}} \leq f(x). \\ \sum_{n=1}^N \frac{e^{-nx}}{\sqrt{n}} &\leq \int_0^N \frac{e^{-ux}}{\sqrt{u}} du \leq \int_0^{\infty} \frac{e^{-ux}}{\sqrt{u}} du. \end{aligned}$$

Therefore

$$\int_1^{\infty} \frac{e^{-ux}}{\sqrt{u}} du \leq f(x) \leq \int_0^{\infty} \frac{e^{-ux}}{\sqrt{u}} du.$$

**E9) Prove that  $\exists C_0$  such that**

$$\lim_{x \rightarrow 0^+} \sqrt{x} f(x) = C_0.$$

Proof: By E8)

$$\begin{aligned} \sqrt{x} f(x) &\leq \int_0^{\infty} \frac{e^{-ux}}{\sqrt{ux}} du = \int_0^{\infty} \frac{e^{-t}}{\sqrt{t}} dt = \sqrt{\pi}. \\ \sqrt{x} f(x) &\geq \int_1^{\infty} \frac{e^{-ux}}{\sqrt{ux}} du = \int_x^{\infty} \frac{e^{-t}}{\sqrt{t}} dt \rightarrow \sqrt{\pi}. \end{aligned}$$

Hence

$$\lim_{x \rightarrow 0^+} \sqrt{x} f(x) = \sqrt{\pi}.$$

**E10) Define the sequence  $\{a_n\}_{n \geq 1}$  as follows:**

$$a_n = \left( \sum_{k=1}^n \frac{1}{\sqrt{k}} \right) - 2\sqrt{n}.$$

Prove that  $\{a_n\}$  converges.

Proof: By Euler-Maclaurin formula, for  $f(x) = 1/\sqrt{x}$ ,

$$\begin{aligned}\sum_{k=1}^n \frac{1}{\sqrt{k}} &= \frac{f(1) + f(n)}{2} + \int_1^n \frac{1}{\sqrt{x}} dx + \int_1^n \tilde{B}_1(x) f'(x) dx \\ &= 2\sqrt{n} - \frac{3}{2} + \frac{1}{2\sqrt{n}} + \int_1^n \tilde{B}_1(x) f'(x) dx\end{aligned}$$

Hence

$$\lim_{n \rightarrow \infty} a_n = - \int_1^\infty \frac{\tilde{B}_1(x)}{2x^{3/2}} dx - \frac{3}{2}.$$

**E11) Prove that for any  $x \in I$ , the function**

$$h(x) = \sum_{n \geq 1} \left( \sum_{k=1}^n \frac{1}{\sqrt{k}} \right) e^{-nx}$$

is well-defined.

Proof: By E10),  $|a_n|$  is bounded, hence

$$h(x) = \sum_{n \geq 1} 2\sqrt{n}e^{-nx} + a_n e^{-nx} = 2g(x) + \sum_{n \geq 1} a_n e^{-nx} \leq 2g(x) + \sup_n |a_n| \cdot \frac{1}{e^x - 1}.$$

**E12) Express  $h(x)$  using  $f(x)$  and find a constant  $C_1$  such that**

$$\lim_{x \rightarrow 0^+} x^{\frac{3}{2}} h(x) = C_1.$$

Proof: Since  $e^{-nx}/k > 0$ , we can interchange the sums

$$h(x) = \sum_{k=1}^{\infty} \frac{1}{\sqrt{k}} \sum_{n=k}^{\infty} e^{-nx} = \sum_{k=1}^{\infty} \frac{e^{-kx}}{\sqrt{k}} \frac{1}{1 - e^{-x}} = \frac{1}{1 - e^{-x}} f(x).$$

Therefore

$$\lim_{x \rightarrow 0^+} x^{3/2} h(x) = \lim_{x \rightarrow 0^+} \sqrt{x} f(x) = \sqrt{\pi}.$$

**E13) Prove that**

$$\lim_{x \rightarrow 0^+} x^{\frac{3}{2}} g(x) = \frac{\sqrt{\pi}}{2}.$$

Proof:

$$\lim_{x \rightarrow 0^+} x^{3/2} |h(x) - 2g(x)| \leq \lim_{x \rightarrow 0^+} \sup_n |a_n| \cdot \frac{x^{3/2}}{e^x - 1} = 0.$$

Hence

$$\lim_{x \rightarrow 0^+} x^{3/2} g(x) = \frac{1}{2} \lim_{x \rightarrow 0^+} x^{3/2} h(x) = \frac{\sqrt{\pi}}{2}.$$

**Part 3**

Given  $A \subset \mathbb{Z}_{\geq 0}$ , we can define a sequence  $\{a_n\}_{n \geq 0}$ :

$$a_n = \begin{cases} 1, & \text{if } n \in A; \\ 0, & \text{if } n \notin A. \end{cases}$$

Define the set  $I_A \subset \mathbb{R}_{\geq 0}$  as follows:

$$I_A = \left\{ x \geq 0 : \text{the series } \sum_{n \geq 0} a_n e^{-nx} \text{ converges} \right\}.$$

Define the function  $f_A : I_A \rightarrow \mathbb{R}$  as follows:

$$f_A(x) = \sum_{n \geq 0} a_n e^{-nx}.$$

Let  $\Phi(A) = \lim_{x \rightarrow 0} x f_A(x)$  (if the limit exists) and let

$$\mathcal{S} = \{A \subset \mathbb{Z}_{\geq 0} : \lim_{x \rightarrow 0^+} x f_A(x) \text{ exists}\}.$$

For example, let

$$A_1 = \{n^2 : n \in \mathbb{Z}_{\geq 1}\}, A_2 = \{p^2 + q^2 : p, q \in \mathbb{Z}_{\geq 1}\}.$$

**E14) Determine the set  $I_A$ .**

Solution: If  $A$  is finite, then  $I_A = \mathbb{R}_{\geq 0}$ . Otherwise  $I_A = \mathbb{R}_{> 0} = I$ .

**E15) Given  $A \subset \mathbb{Z}_{\geq 0}$ , for any  $n \geq 0$ , define the set  $A_{\leq n}$ :**

$$A_{\leq n} = \{ \in A : \leq n \}.$$

Prove that for any  $x > 0$ , the series

$$\sum_{n=0}^{\infty} |A_{\leq n}| \cdot e^{-nx}$$

converges, and satisfy

$$\sum_{n=0}^{\infty} |A_{\leq n}| \cdot e^{-nx} = \frac{f_A(x)}{1 - e^{-x}}.$$

Proof:  $|A_{\leq n}| \leq n + 1$ , hence

$$\sum_{n=0}^{\infty} |A_{\leq n}| \cdot e^{-nx} \text{ converges.}$$

Therefore

$$\sum_{n=0}^{\infty} |A_{\leq n}| \cdot e^{-nx} = \sum_{n=0}^{\infty} \sum_{k=0}^n a_k \cdot e^{-nx} = \sum_{k=0}^{\infty} a_k \cdot \frac{e^{-kx}}{1 - e^{-x}} = \frac{f_A(x)}{1 - e^{-x}}.$$

**E16) Prove that for any  $x > 0$**

$$\frac{f_{A_1}(x)}{1 - e^{-x}} = \sum_{n=0}^{\infty} \lfloor \sqrt{n} \rfloor e^{-nx}.$$

Proof: By E15),

$$|A_1 \leq n| = \sum_{k=0}^n [\sqrt{k} \in \mathbb{Z}_{\geq 1}] = \lfloor \sqrt{n} \rfloor.$$

**E17) Prove that**

$$\lim_{x \rightarrow 0^+} \sqrt{x} f_{A_1}(x)$$

exists and calculate the value of  $\Phi(A_1)$ .

Proof:

$$\lim_{x \rightarrow 0^+} \sqrt{x} f_{A_1}(x) = \lim_{x \rightarrow 0^+} \sqrt{x} (1 - e^{-x}) \left( g(x) - \sum_{n=0}^{\infty} \{\sqrt{n}\} e^{-nx} \right).$$

Since  $1 - e^{-x} \sim x$ ,  $g(x) \sim \frac{\sqrt{\pi}}{2} x^{-3/2}$ , and

$$\left| \sum_{n=0}^{\infty} \{\sqrt{n}\} e^{-nx} \right| \leq \frac{1}{1 - e^{-x}}.$$

Hence

$$\lim_{x \rightarrow 0^+} \sqrt{x} f_{A_1}(x) = \frac{\sqrt{\pi}}{2}.$$

and

$$\Phi(A_1) = \lim_{x \rightarrow 0^+} x f_{A_1}(x) = 0.$$

**E18) Let  $v(n) = \#\{(p, q) \in \mathbb{Z}_{\geq 1}^2 : p^2 + q^2 = n\}$ . Prove that for any  $x > 0$ , the series**

$$\sum_{n \geq 1} v(n) e^{-nx}$$

converges and

$$\sum_{n \geq 1} v(n) e^{-nx} = (f_{A_1}(x))^2.$$

Proof: Since  $v(n) \leq n$ ,  $\sum_{n \geq 1} v(n) e^{-nx}$  converges.

$$\sum_{n \geq 1} v(n) e^{-nx} = \sum_{n \geq 1} \sum_{k=0}^n a_k a_{n-k} e^{-nx} = \sum_{n \geq 1} \sum_{k=0}^n a_k e^{-kx} \cdot a_{n-k} e^{-(n-k)x} = (f_{A_1}(x))^2.$$

**E19) Prove that for any  $x > 0$** 

$$f_{A_2}(x) \leq (f_{A_1}(x))^2$$

and give an upper-bound of  $\Phi(A_2)$  (assuming it exists).

Proof:

$$f_{A_2}(x) = \sum_{n \geq 1} [v(n) \geq 1] \cdot e^{-nx} \leq \sum_{n \geq 1} v(n) e^{-nx} = (f_{A_1}(x))^2.$$

Hence

$$\Phi(A_2) = \lim_{x \rightarrow 0^+} x f_{A_2}(x) \leq \lim_{x \rightarrow 0^+} (\sqrt{x} f_{A_1}(x))^2 = \frac{\pi}{4}.$$

**Part 4**Assume  $\{a_n\}_{n \geq 0}$  is a sequence of non-negative numbers, such that for any  $x > 0$  the series

$$S(x) = \sum_{n \geq 0} a_n e^{-nx}$$

converges. Moreover, assume that the limit below exists:

$$\lim_{x \rightarrow 0^+} x S(x) = \lim_{x \rightarrow 0^+} x \sum_{n \geq 0} a_n e^{-nx} = \in [0, +\infty).$$

Let  $F = \{f : [0, 1] \rightarrow \mathbb{R}\}$ ,  $E_0 = C([0, 1])$ . Let  $E$  be the space of piecewise continuous functions, and define the norm on  $E$ :

$$\|\psi\|_\infty = \sup_{x \in [0, 1]} |\psi(x)|.$$

**E20) Define  $L : E \rightarrow F$  as follows:**

$$(L(\psi))(x) = \sum_{n=0}^{\infty} a_n e^{-nx} \psi(e^{-nx}), \psi \in E.$$

Prove that  $L$  is well-defined and is linear. Moreover, if for any  $x \in [0, 1]$ ,  $\psi_1(x) \leq \psi_2(x)$ , then for any  $x \in [0, 1]$ ,

$$(L(\psi_1))(x) \leq (L(\psi_2))(x).$$

Proof: Since  $\psi \in E$ ,  $\psi$  is bounded, hence  $L$  is well-defined and is clearly linear. The inequality holds since  $a_n$  are non-negative.**E21) Define the subspace of  $E$** 

$$E_1 = \{\psi \in E : \lim_{x \rightarrow 0^+} x(L(\psi))(x) \text{ exists}\}.$$

Define the linear map  $\Delta : E_1 \rightarrow \mathbb{R}$  as follows:

$$\Delta(\psi) = \lim_{x \rightarrow 0^+} x(L(\psi))(x), \psi \in E_1.$$

Prove that  $E_1$  is a subspace of  $E$  and there is a constant  $M > 0$  such that for any  $\psi \in E_1$ ,

$$|\Delta(\psi)| \leq M \|\psi\|_\infty.$$

Proof: Since  $L$  is linear, so is  $\Delta$ , thus  $E_1$  is clearly a subspace of  $E$ .

$$|\Delta(\psi)| = \left| \lim_{x \rightarrow 0^+} x \sum_{n=0}^{\infty} a_n e^{-nx} \psi(e^{-nx}) \right| \leq \|\psi\|_\infty \cdot \left| \lim_{x \rightarrow 0^+} x S(x) \right| = \|\psi\|_\infty.$$



**E22) For the polynomial  $P_n(x) = x^n$ , prove that  $P_n \in E_1$  and calculate  $\Delta(P_n)$ .**

Proof:

$$\Delta(P_n) = \lim_{x \rightarrow 0^+} x \sum_{k=0}^{\infty} a_k e^{-kx} e^{-nkx} = \frac{1}{n+1}.$$

**E23) Prove that  $E_0 \subset E_1$  and for every  $\psi \in E_0$  calculate  $\Delta(\psi)$ .**

Proof: Since  $\Delta$  is linear, by E22) we know that for any polynomial  $P$ ,

$$\Delta(P) = \int_0^1 P(x) dx.$$

By Stone-Weierstraß theorem, any continuous function on  $[0, 1]$  can be uniformly approximated with polynomials, hence (same as E24)

$$\Delta(\psi) = \int_0^1 \psi(x) dx, \forall \psi \in E_0.$$

**E24) For  $a \in (0, 1)$ ,  $\varepsilon \in (0, \min(a, 1 - a))$ , define the functions**

$$g_{-}(x) = \begin{cases} 1, & x \in [0, a - \varepsilon]; \\ \frac{a-x}{\varepsilon}, & x \in (a - \varepsilon, a); \\ 0, & x \in [a, 1] \end{cases}, g_{+}(x) = \begin{cases} 1, & x \in [0, a]; \\ \frac{a+\varepsilon-x}{\varepsilon}, & x \in (a, a + \varepsilon); \\ 0, & x \in [a + \varepsilon, 1] \end{cases}.$$

Prove that  $g_{\pm} \in E_0$  and calculate  $\Delta(g_{\pm})$ . Further prove that  $\mathbf{1}_{[0,a]} \in E_1$  and calculate  $\Delta(\mathbf{1}_{[0,a]})$ .

Proof:  $g_{\pm} \in E_0$  is trivial, and  $\Delta(g_{\pm}) = \int_0^1 g_{\pm} = (a \pm \varepsilon/2)$ . Since  $g_{-} \leq \mathbf{1}_{[0,a]} \leq g_{+}$ ,

$$x(L(g_{-}))(x) \leq x(L(\mathbf{1}_{[0,a]}))(x) \leq x(L(g_{+}))(x)$$

For any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that for any  $0 < x < \delta$ ,

$$|x(L(g_{-}))(x) - \Delta(g_{-})|, |x(L(g_{+}))(x) - \Delta(g_{+})| < \frac{\varepsilon}{2}.$$

Hence for any  $0 < x < \delta$ ,

$$\begin{aligned} x(L(\mathbf{1}_{[0,a]}))(x) &\leq x(L(g_{+}))(x) \leq \Delta(g_{+}) + \frac{\varepsilon}{2} = a + \varepsilon. \\ x(L(\mathbf{1}_{[0,a]}))(x) &\geq x(L(g_{-}))(x) \geq \Delta(g_{-}) - \frac{\varepsilon}{2} = a - \varepsilon. \end{aligned}$$

Therefore

$$\Delta(\mathbf{1}_{[0,a]}) = \lim_{x \rightarrow 0^+} x(L(\mathbf{1}_{[0,a]}))(x) = a.$$

**E25) Prove that  $E_1 = E$  and for  $\psi \in E$  determine the formula of  $\Delta(\psi)$ .**

Proof: Use the same method as E24) applied to Darboux's sum. Hence

$$E_1 = E, \text{ and } \Delta(\psi) = \int_0^1 \psi(x) dx.$$

**E26) Define the function**

$$\psi(x) = \begin{cases} 0, & x \in [0, e^{-1}); \\ \frac{1}{x}, & x \in [e^{-1}, 1]. \end{cases}$$

Prove the following equation by calculating  $L(\psi)\left(\frac{1}{N}\right)$ :

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=0}^N a_k = .$$

Proof:

$$(L(\psi))\left(\frac{1}{N}\right) = \sum_{n=0}^{\infty} a_n e^{-n/N} \psi(e^{-n/N}) = \sum_{n=0}^N a_n.$$

Hence by E25),

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^N a_n = \Delta(\psi) = \int_0^1 \psi(x) dx = .$$

**E27) Consider  $A \in \mathcal{S}$ , and calculate**

$$\lim_{n \rightarrow \infty} \frac{|A_{\leq n}|}{n}.$$

which is called the asymptomatic density of  $A$  on  $\mathbb{Z}_{\geq 0}$ .

Solution:

$$\lim_{n \rightarrow \infty} \frac{|A_{\leq n}|}{n} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^N a_n = \lim_{x \rightarrow 0^+} x \sum_{n=0}^{\infty} a_n e^{-nx} = \Phi(A).$$

**E28) Calculate**

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n v(k)}{n},$$

and give an upper-bound of the asymptomatic density of  $A_2$ .

Solution:

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n v(k)}{n} = \lim_{x \rightarrow 0^+} x \sum_{n=0}^{\infty} v(n) e^{-nx} = \lim_{x \rightarrow 0^+} x (f_{A_1}(x))^2 = \frac{\pi}{4}.$$

From E19)  $\Phi(A_2) \leq \frac{\pi}{4}$ .

Quote:

God does not care about our mathematical difficulties. He integrates empirically.

—Albert Einstein

## 12 Homework 12: Oscillatory Integral

### 12.1 PSA: Stieltjes Integral

Let  $\mu$  be a monotonic function on  $I = [a, b]$ .

**A1) For any pair of partitions  $\sigma, \sigma' \in \mathcal{S}(I)$ ,**

$$\underline{S}_\mu(f; \sigma) \leq \overline{S}_\mu(f; \sigma').$$

Proof: Suppose  $\mathcal{C} = \sigma \cup \sigma'$ , then

$$\underline{S}_\mu(f; \sigma) \leq \underline{S}_\mu(f; \mathcal{C}) \leq \overline{S}_\mu(f; \mathcal{C}) \leq \overline{S}_\mu(f; \sigma').$$

**A2) For any  $\rho \in C([a, b])$ ,  $\rho \geq 0$ ,  $\mu(x) = \int_a^x \rho(t) dt$ . Prove that for any  $f \in \mathcal{R}([a, b])$ ,  $f \in \mathcal{R}([a, b]; \mu)$  and**

$$\int_a^b f d\mu = \int_a^b f(x) \rho(x) dx.$$

Proof: Consider any  $\mathcal{C} = \{x_0, x_1, \dots, x_n\}$ , then if we denote  $m_i = \inf_{x \in [x_{i-1}, x_i]} f(x)$ ,  $u_i = \inf_{x \in [x_{i-1}, x_i]} \rho(x)$ ,  $v_i = \inf_{x \in [x_{i-1}, x_i]} f(x) \rho(x)$ ,  $M = \sup_{x \in [a, b]} f(x)$ ,  $v_i - m_i u_i \leq f(t) \rho(t) - f(t) u_i \leq M \omega_\rho(x_{i-1}, x_i)$ . Hence for any  $\varepsilon > 0$  there exists a  $\delta > 0$ , for any  $\max\{x_i - x_{i-1}\} < \delta$ ,  $\sup_{x, y \in [x_{i-1}, x_i]} |\rho(x) - \rho(y)| < \varepsilon$ . Then

$$\begin{aligned} \underline{S}(f\rho; \mathcal{C}) &= \sum_{k=1}^n v_k(x_k - x_{k-1}) \leq \sum_{k=1}^n u_k m_k(x_k - x_{k-1}) + M\varepsilon(b-a) \\ &\leq M\varepsilon(b-a) + \sum_{k=1}^n m_k \int_{x_{k-1}}^{x_k} \rho(t) dt = M\varepsilon(b-a) + \underline{S}_\mu(f; \mathcal{C}). \end{aligned}$$

The other side is similar, hence  $\sup\{\underline{S}_\mu(f; \mathcal{C})\} = \inf\{\overline{S}_\mu(f; \mathcal{C})\}$  so  $f \in \mathcal{R}([a, b]; \mu)$  and

$$\int_a^b f d\mu = \int_a^b f(x) \rho(x) dx.$$

**A3) Prove that  $\mathcal{R}(I; \mu)$  is a linear space on  $\mathbb{R}$  and the integration operator**

$$\int_a^b \cdot d\mu : \mathcal{R}(I; \mu) \rightarrow \mathbb{R}, f \mapsto \int_a^b f d\mu.$$

is linear.

Proof: Since  $\underline{S}_\mu(\cdot; \mathcal{C})$  and  $\overline{S}_\mu(\cdot; \mathcal{C})$  is linear for any  $\mathcal{C}$ ,  $\mathcal{R}(I; \mu)$  is clearly a linear space on  $\mathbb{R}$ , and  $\int_a^b \cdot d\mu$  is a linear operator.

**A4) Suppose  $f_1, f_2 \in \mathcal{R}(I; \mu)$ . If the any  $x \in I$ ,  $f_1(x) \leq f_2(x)$ , then**

$$\int_a^b f_1 d\mu \leq \int_a^b f_2 d\mu.$$

Proof: By A3), we can assume  $f_1 = 0$ . Then for any  $\mathcal{C}$ ,  $\underline{S}_\mu(f; \mathcal{C}) \geq 0$  since  $f \geq 0$ , hence  $\int_a^b f d\mu = \sup\{\underline{S}_\mu(f; \mathcal{C})\} \geq 0$ .

**A5) If  $f \in \mathcal{R}([a, b]; \mu)$ , then for any  $c \in [a, b]$ ,  $f|_{[a, c]}$  and  $f|_{[c, b]}$  are both Stieltjes integrable and**

$$\int_a^b f \, d\mu = \int_a^c f \, d\mu + \int_c^b f \, d\mu.$$

Proof: For any partition  $\sigma$ , let  $\sigma' = \sigma \cup \{c\}$ , then  $\sigma'$  can be split into two partitions of the intervals  $[a, c]$  and  $[c, b]$ :  $\sigma' = \sigma_1 \cup \sigma_2$ , such that  $\underline{S}_\mu(f; \sigma') = \underline{S}_\mu(f; \sigma_1) + \underline{S}_\mu(f; \sigma_2)$  and  $\overline{S}_\mu(f; \sigma') = \overline{S}_\mu(f; \sigma_1) + \overline{S}_\mu(f; \sigma_2)$ . Hence

$$\inf \underline{S}_\mu(f; \sigma_1) + \inf \underline{S}_\mu(f; \sigma_2) \leq \inf \underline{S}_\mu(f; \sigma') \leq \sup \overline{S}_\mu(f; \sigma') \leq \sup \overline{S}_\mu(f; \sigma_1) + \sup \overline{S}_\mu(f; \sigma_2).$$

Therefore

$$\int_a^b f \, d\mu = \int_a^c f \, d\mu + \int_c^b f \, d\mu.$$

**A6) If  $f, g \in \mathcal{R}([a, b]; \mu)$ , then  $f \cdot g \in \mathcal{R}([a, b]; \mu)$ .**

Proof: Same as in the case of the Riemann integral.

**A7) Define Stieltjes integral on the interval  $[0, \infty)$ : Suppose  $f \in C([0, \infty))$  is continuous and bounded, define**

$$\int_0^\infty f \, d\mu = \lim_{M \rightarrow \infty} \int_0^M f \, d\mu.$$

Suppose  $\{\alpha_n\}_{n \geq 1}$  is a sequence of positive real numbers and  $\sum_{n=1}^\infty \alpha_n$  converges, define the monotonic function  $\mu = \sum_{n=1}^\infty \alpha_n \mathbf{1}_{\geq n}$ , then

$$\int_1^\infty f \, d\mu = \sum_{n=1}^\infty \alpha_n f(n).$$

Proof: Note that

$$\mu(x+0) - \mu(x-0) = \begin{cases} 0, & x \notin \mathbb{Z}, \\ \alpha_x, & x \in \mathbb{Z}. \end{cases}$$

Hence

$$\int_0^N f \, d\mu = \sum_{n=1}^{N-1} f(n) \alpha_n.$$

By definition,

$$\int_0^\infty f \, d\mu = \sum_{n=1}^\infty \alpha_n f(n).$$

**A8)  $f, g \in \mathcal{R}([a, b]; \mu)$  are real-valued Riemann integrable functions. Suppose for any  $x \in [a, b]$ ,  $g(x) \geq 0$ . Let**

$$m = \inf_{x \in I} f(x), \quad M = \sup_{x \in I} f(x).$$

Then there exists  $\epsilon \in [m, M]$  such that

$$\int_a^b fg \, d\mu = \int_a^b g \, d\mu.$$

Proof: Note that  $mg \leq fg \leq Mg$ , and use A4).

**A9) Construct a Stieltjes integral to show that Abel summation method is a special case of integration by parts.**

Proof:

The Abel summation formula states that

$$\sum_{i=1}^n T_i(S_i - S_{i-1}) = T_n S_n - T_1 S_0 - \sum_{i=1}^{n-1} S_i(T_{i+1} - T_i).$$

Consider the monotonically increasing function  $\mu : [0, n] \rightarrow \mathbb{R}, x \mapsto T_{\lceil x \rceil}, \mu(0) = T_1$ , and  $f$  be a polynomial such that  $f(k) = S_k$  for  $k = 0, 1, \dots, n$ . Then

$$\int_0^n f' \mu \, dx = \sum_{k=1}^n \int_{k-1}^k f' \mu = \sum_{k=1}^n \int_{k-1}^k f'(x) T_k \, dx = \sum_{k=1}^n T_k (S_k - S_{k-1}).$$

While

$$\int_0^n f \, d\mu = \sum_{k=1}^{n-1} f(k)(\mu(k+0) - \mu(k)) = \sum_{k=1}^{n-1} S_k (T_{k+1} - T_k).$$

and

$$f\mu \Big|_0^n = T_n S_n - T_1 S_0.$$

Hence the formula is a special case of integration by parts.

## 12.2 PSB: Convergence of Improper Integrals

$b$  can be  $\infty$ .

**B1) Assume  $f : [a, b) \rightarrow \mathbb{R}$ , and for any  $b^- < b$ ,  $f$  is integrable on  $[a, b^-]$ . Prove that the integral  $\int_a^b f(x) \, dx$  exists iff: for any  $\varepsilon > 0$ ,  $\exists b(\varepsilon) \in (a, b)$  such that for any  $b', b'' > b(\varepsilon)$ ,  $\left| \int_{b'}^{b''} f(x) \, dx \right| < \varepsilon$ .**

Proof: Let

$$F(t) = \int_a^t f(x) \, dx, \forall t \in [a, b).$$

Then  $\int_a^b f(x) \, dx$  exists iff  $\lim_{t \rightarrow b^-} F(t)$  exists, which is equivalent to

$$\forall \varepsilon > 0, \exists b(\varepsilon) \in (a, b), \forall b', b'' > b(\varepsilon), \left| \int_{b'}^{b''} f(x) \, dx \right| = |F(b'') - F(b')| < \varepsilon.$$

**B2) If  $|f(x)| \leq F(x), x \in [a, b)$  and  $\int_a^b F(x) \, dx$  converges, then  $\int_a^b f(x) \, dx$  converges.**

Proof: Use B1) and

$$\left| \int_u^v f(x) \, dx \right| \leq \int_u^v F(x) \, dx.$$

**B3) Prove the Dirichlet test for convergence: if  $f, g : [a, \infty) \rightarrow \mathbb{R}$  satisfy**

- $f$  is continuous and there exists  $A > 0$ , such that for any  $M \geq a$ ,

$$\left| \int_a^M f(x) dx \right| \leq A.$$

- $g$  is monotonic and  $\lim_{x \rightarrow \infty} g(x) = 0$ .  
Then  $\int_a^\infty f(x)g(x) dx$  converges.

**Lemma: The Second Integral Mean Value Theorem**

If  $f$  is integrable and  $g$  is monotonic and non-negative(or non-positive) on  $[a, b]$ , then there exists  $c \in (a, b)$  such that

$$\int_a^b f(x)g(x) dx = g(a) \int_a^c f(x) dx + g(b) \int_c^b f(x) dx.$$

Proof: Assume that  $g$  is non-negative and monotonically decreasing. It is easy to see that there exists  $\xi \in (a, b)$  such that

$$\int_a^b f(x)g(x) dx = g(a) \int_a^\xi f(x) dx.$$

Apply the above formula to  $f(x)$  and  $g(x) - g(b)$  and we get

$$\int_a^b f(x)g(x) dx = g(a) \int_a^\xi f(x) dx + g(b) \int_\xi^b f(x) dx.$$

Proof of B3): Since  $|\int_u^v f(x) dx| \leq 2A$ , by lemma

$$\left| \int_u^v f(x)g(x) dx \right| \leq 2A(|g(u)| + |g(v)|).$$

By B1), the integral converges.

**B4) Prove the Abel test of convergence:**

If  $f, g : [a, \infty) \rightarrow \mathbb{R}$  satisfy:

- $\int_a^\infty f(x) dx$  exists.
- $g$  is monotonic and  $g$  is bounded.  
Then  $\int_a^\infty f(x)g(x) dx$  converges.

Proof: Suppose  $g$  is monotonically increasing, then

$$\left| \int_u^v f(x)(g(x) - g(a)) dx \right| \leq 2M \left( \left| \int_u^\xi f(x) dx \right| + \left| \int_\xi^v f(x) dx \right| \right) \rightarrow 0$$

since  $\int_a^\infty f(x) dx$  converges. Therefore both  $\int_a^\infty f(x)(g(x) - g(a)) dx$  and  $\int_a^\infty f(x)g(a) dx$  converges, hence  $\int_a^\infty f(x)g(x) dx$  converges.

**B5) Determine whether the following integrals converges:**

(1)

$$\int_0^\infty \frac{\log(1+x)}{x^p} dx$$

(absolutely) convergent when  $1 < p < 2$ , diverges when  $p \leq 1$  or  $p \geq 2$ .

(2)

$$\int_1^\infty \frac{\sin x}{x^p} dx$$

Absolutely convergent when  $p > 1$ , conditionally convergent when  $0 < p \leq 1$ , diverges when  $p \geq 0$ .

(3)

$$\int_1^\infty \sin x^2 dx = \frac{1}{2} \int_1^\infty \frac{\sin y}{y^{1/2}} dy$$

is conditionally convergent.

(4)

$$\int_0^\infty \frac{\sin^2 x}{x} dx$$

diverges

(5)  $p, q > 0$ ,

$$\int_0^{2\pi} \sin^{-p} x \cos^{-q} x dx$$

Absolutely convergent when  $p, q < 1$ , diverges when  $p \geq 1$  or  $q \geq 1$ .

(6)

$$\int_0^\infty x^p \sin(x^q) dx$$

If  $q = 0$  the integral diverges. Assume  $q \neq 0$  below.

$$\int_0^\infty x^p \sin(x^q) dx = \frac{1}{q} \int_0^\infty y^{(p+1)/q-1} \sin y dy.$$

Let  $\alpha = \frac{p+1}{q} - 1$ , then the integral

- diverges if  $\alpha \leq -2$  or  $\alpha \geq 0$ ,
  - converges absolutely if  $-2 < \alpha < -1$ .
  - converges conditionally if  $-1 \leq \alpha < 0$ .
- (7)  $q \geq 0$ ,

$$\int_0^\infty \frac{x^p \sin x}{1+x^q} dx$$

If  $p \leq -2$ , then the integral diverges near 0, since  $x^p \sin x \sim x^{p+1}$ . The integral converges (absolutely) near 0 otherwise. Assume  $p > -2$  below.

If  $p - q < -1$  then the integral converges absolutely when it tends to infinity, since  $\frac{x^p}{1+x^q} \sim x^{p-q}$ .

If  $-1 \leq p - q < 0$  then the integral converges conditionally, since the integral of  $(x^{p-q})'$  converges.

(8)

$$\begin{aligned} \int_0^{\pi/2} \frac{\log \sin x}{\sqrt{x}} dx &= 2 \int_0^{\pi/2} \log \sin x d\sqrt{x} \\ &= 2\sqrt{x} \log \sin x \Big|_0^{\pi/2} - 2 \int_0^{\pi/2} \sqrt{x} \cot x dx \\ &= -2 \int_0^{\pi/2} \frac{\sqrt{x}}{\sin x} \cos x dx \end{aligned}$$

converges, since  $\int_0^1 \frac{1}{\sqrt{x}} dx$  converges.

(9)

$$\int_e^\infty \frac{\log \log x}{\log x} \sin x dx = \int_1^\infty \frac{\log y}{y} e^y \sin e^y dy.$$

It is easy to see the integral does not converge absolutely.

Meanwhile

$$f'(x) = \left( \frac{\log \log x}{\log x} \right)' = \frac{1 - \log \log x}{(\log x)^2 x},$$

and

$$\int_e^\infty \frac{\log \log x - 1}{(\log x)^2 x} dx = \int_1^\infty \frac{\log y - 1}{y^2} dy = \int_0^\infty \frac{t - 1}{e^t} dt.$$

converges.

By Lagrange mean value theorem,

$$\begin{aligned} \int_{2\pi}^\infty \frac{\log \log x}{\log x} \sin x dx &= \sum_{n=1}^\infty \int_{2n\pi}^{(2n+1)\pi} (f(x) - f(x + \pi)) \sin x dx \\ &\leq \sum_{n=1}^\infty -2\pi f'(2n\pi) \leq 2\pi \int_e^\infty -f'(x) dx \end{aligned}$$

converges.

### 12.3 PSC: Oscillatory Integral

$F(t), G(t) : [1, \infty) \rightarrow \mathbb{R}$ ,  $\lim_{t \rightarrow \infty} G(t) = 0$ . Assume that for any  $t \geq 1$ ,  $G(t) \neq 0$ . If

$$\lim_{t \rightarrow \infty} \frac{F(t)}{G(t)} = 1.$$

Then we say  $F, G$  have the same order, and  $F \sim G$ .

#### Part 1

**C1)**  $d > 0$  is a given real number. Assume  $g \in C^1([0, d])$ . Prove that there is a constant  $C$ , such that

$$\left| \int_0^d e^{-tx} g(x) dx \right| \leq \frac{C}{t}.$$

Proof: Let  $C = \sup_{x \in [0, d]} |g(x)|$ , then

$$\left| \int_0^d e^{-tx} g(x) dx \right| \leq C \int_0^d e^{-tx} dx = \frac{C}{t} (1 - e^{-td}) \leq \frac{C}{t}.$$



**C2)** Assume  $d > 0$ ,  $g \in C([0, d])$  and  $g(0) \neq 0$ . Prove that

$$\int_0^d e^{-tx} g(x) dx \sim \frac{g(0)}{t}.$$

Proof: Let  $M = \sup_{x \in [0, d]} |g(x)|$ , then

$$\begin{aligned} \left| \int_0^d e^{-tx} t \frac{g(x)}{g(0)} dx - 1 \right| &= \left| \int_0^{td} e^{-u} \frac{g(u/t)}{g(0)} du - \int_0^\infty e^{-u} du \right| \\ &\leq \int_{td}^\infty e^{-u} du + \int_0^{td} e^{-u} \left| \frac{g(u/t)}{g(0)} - 1 \right| du + \int_N^{td} e^{-u} \left| \frac{g(u/t)}{g(0)} - 1 \right| du \\ &\leq e^{-td} + \sup_{0 \leq x \leq N/t} \left| \frac{g(x)}{g(0)} - 1 \right| + \left( \frac{M}{|g(0)|} + 1 \right) \int_N^{td} e^{-u} du \rightarrow 0. \end{aligned}$$

(let  $t \rightarrow \infty$  then let  $N \rightarrow \infty$ ).

**C3)**  $d > 0, g \in C([0, d]), g(0) \neq 0$ . Prove that

$$\int_0^d e^{-tx^2} g(x) dx \sim \frac{\sqrt{\pi} \cdot g(0)}{2\sqrt{t}}.$$

Proof: Same as C2), let  $M = \sup_{x \in [0, d]} |g(x)/g(0)|$ , then

$$\begin{aligned} \left| \int_0^d e^{-tx^2} \sqrt{t} \frac{g(x)}{g(0)} dx - \frac{\sqrt{\pi}}{2} \right| &= \left| \int_0^{d\sqrt{t}} e^{-u^2} \frac{g(u/\sqrt{t})}{g(0)} du - \int_0^\infty e^{-u^2} du \right| \\ &\leq \int_{d\sqrt{t}}^\infty e^{-u^2} du + \int_0^{d\sqrt{t}} e^{-u^2} \left| \frac{g(u/\sqrt{t})}{g(0)} - 1 \right| du + \int_N^{d\sqrt{t}} e^{-u^2} (M+1) du. \end{aligned}$$

which tends to 0, same as C2).

---

For  $t \geq 1$ ,  $f, \varphi \in C([a, b])$ , define the function

$$F(t) = \int_a^b e^{-t\varphi(x)} f(x) dx.$$

Our goal is to study  $F(t)$  when  $t \rightarrow \infty$ .

**C4)** Assume  $\varphi \in C^1([a, b])$ , and for any  $x \in [a, b]$ ,  $\varphi'(x) \neq 0$ . Further assume that  $\varphi'(x) > 0$ . Let  $d = \varphi(b) - \varphi(a)$ . Prove that

$$\Psi : [a, b] \rightarrow [0, d], x \mapsto \varphi(x) - \varphi(a),$$

is a continuously differentiable bijection on  $[a, b]$ .

Proof:  $\varphi$  is monotonic by  $\varphi'(x) > 0$ , hence  $\Psi$  is a bijection and  $\Psi' = \psi'$ .

**C5) Assume**  $\varphi \in C^1([a, b])$ , **and for any**  $x \in [a, b]$ ,  $\varphi'(x) \neq 0$ . **Prove that if**  $f(a) \neq 0$ , **then when**  $t \rightarrow \infty$ ,

$$F(t) \sim \frac{f(a)}{\varphi'(a)} \frac{e^{-t\varphi(a)}}{t}.$$

Proof: Let  $g(x) = f(x)/\Psi'(x)$ , and  $h = (t\Psi)^{-1}$  then

$$\begin{aligned} \left| F(t) \frac{t}{e^{-t\varphi(a)}} - \frac{f(a)}{\varphi'(a)} \right| &= \left| \int_a^b e^{-t\Psi(x)} t f(x) dx - \frac{f(a)}{\Psi'(a)} \right| = \left| \int_a^b e^{-t\Psi(x)} g(x) dt\Psi(x) - g(a) \right| \\ &= \left| \int_0^{t\Psi(b)} e^{-u} g(h(u)) du - g(h(0)) \int_0^\infty e^{-u} du \right| \\ &= |g(h(0))| \int_{t\Psi(b)}^\infty e^{-u} du + \int_0^{N\Psi(b)} e^{-u} |g(h(u)) - g(h(0))| du \\ &\quad + \int_{N\Psi(b)}^{t\Psi(b)} e^{-u} |g(h(u)) - g(h(0))| du \\ &\leq |g(a)| e^{-t\Psi(b)} + \sup_{x \in [a, \Psi^{-1}(N\Psi(b)/t)]} |g(x) - g(a)| + \int_{N\Psi(b)}^{t\Psi(b)} e^{-u} 2M du. \end{aligned}$$

which tends to 0 since  $g$  is continuous. ( $M = \sup_{x \in [a, b]} |g(x)|$ ).

**C6) Assume that**  $\varphi \in C^2([a, b])$ ,  $\varphi'(a) = 0$ ,  $\varphi''(x) > 0$  **and for any**  $x \in (a, b]$ ,  $\varphi'(x) > 0$ . **Let**  $d = \sqrt{\varphi(b) - \varphi(a)}$ . **Prove that**

$$\Psi : [a, b] \rightarrow [0, d], x \mapsto \sqrt{\varphi(x) - \varphi(a)}.$$

is a continuously differentiable bijection on  $[a, b]$ , and calculate  $\Psi'(a)$ .

Proof: Trivial.  $\Psi' = \frac{\varphi'}{2\Psi}$ , hence

$$\Psi'(a) = \lim_{x \rightarrow a^+} \frac{\varphi'(x)}{2\sqrt{\varphi(x) - \varphi(a)}} = \lim_{x \rightarrow a^+} \frac{\varphi''(x)}{\varphi'(x)/\sqrt{\varphi(x) - \varphi(a)}} = \sqrt{\frac{\varphi''(a)}{2}}.$$

**C7) Assume**  $\varphi \in C^2([a, b])$ ,  $\varphi'(a) = 0$ ,  $\varphi''(a) > 0$ . **Prove that if**  $f(a) \neq 0$ , **when**  $t \rightarrow \infty$ ,

$$F(t) \sim \frac{\sqrt{\pi} f(a)}{\sqrt{2\varphi''(a)}} \frac{e^{-t\varphi(a)}}{\sqrt{t}}.$$

Proof: Let  $g = f/\Psi'$ ,  $h = (\sqrt{t}\Psi)^{-1}$ , then

$$F(t) \frac{\sqrt{t}}{e^{-t\varphi(a)}} = \int_a^b e^{-t\Psi^2(x)} f(x) \sqrt{t} dx = \int_a^b e^{-t\Psi^2(x)} g(x) d\sqrt{t}\Psi(x) = \int_0^{\sqrt{t}\Psi(b)} e^{-u^2} g(h(u)) du.$$

Hence

$$\begin{aligned}
 \left| F(t) \frac{\sqrt{t}}{e^{-t\varphi(a)}} - \frac{\sqrt{\pi}}{2} g(a) \right| &= \left| \int_0^{\sqrt{t}\Psi(b)} e^{-u^2} g(h(u)) \, du - \int_0^\infty e^{-u^2} g(h(0)) \, du \right| \\
 &\leq g(a) \int_{\sqrt{t}\Psi(b)}^\infty e^{-u^2} \, du + \int_0^{N\Psi(b)} e^{-u^2} |g(h(u)) - g(h(0))| \, du \\
 &\quad + \int_{N\Psi(b)}^{\sqrt{t}\Psi(b)} e^{-u^2} 2M \, du \\
 &\leq g(a) e^{-\sqrt{t}\Psi(b)} + \sqrt{\pi} \sup_{x \in [a, \Psi^{-1}(N\Psi(b)/\sqrt{t})]} |g(x) - g(a)| + 2M e^{-N\Psi(b)}.
 \end{aligned}$$

which tends to 0 as  $t \rightarrow \infty$  and  $N \rightarrow \infty$ , since  $g$  is continuous.  
 (A much simpler solution can be given using the Laplace method)

**C8) Given  $f \in C((0, \infty))$ ,  $\varphi \in C^2((0, \infty))$ . Assume that**

- exists a unique  $a \in (0, \infty)$  such that  $\varphi'(a) = 0$ ;
- $\varphi''(a) > 0$ ,  $f(a) \neq 0$ ;
- $\int_0^\infty e^{-\varphi(x)} |f(x)| \, dx$  converges.

Prove that when  $t \rightarrow \infty$ , the function  $G(t) = \int_0^\infty e^{-t\varphi(x)} f(x) \, dx$  satisfy

$$G(t) \sim \frac{\sqrt{2\pi} f(a)}{\sqrt{\varphi''(a)}} \frac{e^{-t\varphi(a)}}{\sqrt{t}}.$$

Proof: (Simple application of the Laplace method)

Apply C7) to the intervals  $[a/2, a]$  and  $[a, 2a]$ , then

$$\int_{a/2}^{2a} e^{-t\varphi(x)} f(x) \, dx \sim \frac{\sqrt{2\pi} f(a)}{\sqrt{\varphi''(a)}} \cdot \frac{e^{-t\varphi(a)}}{\sqrt{t}}.$$

While the integral on the intervals  $(0, a/2)$ ,  $(2a, \infty)$  converges rapidly. Hence

$$G(t) \sim \frac{\sqrt{2\pi} f(a)}{\sqrt{\varphi''(a)}} \frac{e^{-t\varphi(a)}}{\sqrt{t}}.$$

**C9)  $\Gamma(n) = (n-1)!$ .**

Proof:

$$\Gamma(n+1) = \int_0^\infty t^n e^{-t} \, dt = - \int_0^\infty t^n \, de^{-t} = n \int_0^\infty t^{n-1} e^{-t} \, dt = n\Gamma(n).$$

**C10) Prove Stirling's approximation**

$$n! \sim \sqrt{2\pi} \frac{n^{n+1/2}}{e^n}.$$

Proof: Note that, by substituting  $t = ns$ ,

$$n! = \Gamma(n+1) = \int_0^\infty e^{-t} t^n \, dt = n^{n+1} \int_0^\infty e^{-n(s-\log s)} \, ds.$$

Hence

$$\frac{\Gamma(t+1)}{t^{t+1}} \sim \sqrt{2\pi} \frac{e^{-t}}{\sqrt{t}}.$$

## Part 2

For  $\lambda \geq 1$ ,  $f, \varphi \in C^\infty([a, b])$ , define the function

$$I(\lambda) = \int_a^b e^{i\lambda\varphi(x)} f(x) dx.$$

Our goal is to study  $I(\lambda)$  when  $\lambda \rightarrow \infty$ .

**C11) Assume that for any  $x \in [a, b]$ ,  $\varphi'(x) \neq 0$ . Define the maps**

$$\begin{aligned} L : C^\infty([a, b]) &\rightarrow C^\infty([a, b]), h \mapsto \frac{1}{i\lambda\varphi'(x)} h'(x), \\ M : C^\infty([a, b]) &\rightarrow C^\infty([a, b]), h \mapsto -\left(\frac{h}{i\varphi'}\right)'(x). \end{aligned}$$

Assume that  $f, g \in C^\infty([a, b])$ . Prove that if there exists  $c > 0$  such that for any  $x \in [a, a+c] \cup [b-c, b]$ ,  $h(x) = 0$ , then  $M/\lambda$  is the adjoint of  $L$ :

$$\int_a^b h \cdot Lg = \frac{1}{\lambda} \int_a^b g \cdot Mh.$$

Proof: By integration of parts,

$$\int_a^b h \cdot Lg = \int_a^b \frac{h}{i\lambda\varphi'} dg = - \int_a^b g d\left(\frac{h}{i\lambda\varphi'}\right) = \frac{1}{\lambda} \int_a^b g \cdot Mh.$$

**C12) Assume that for any  $x \in [a, b]$ ,  $\varphi'(x) \neq 0$  and  $f$  vanishes near  $a$  and  $b$ . prove that for any  $n \in \mathbb{Z}_{\geq 1}$ , there is a constant  $c_n$  independent of  $\lambda$  such that**

$$|I(\lambda)| \leq \frac{c_n}{\lambda^n}.$$

Proof: Let  $g = e^{i\lambda}\varphi$ , then  $Lg = g$ .  $f \in C^\infty([a, b])$  vanishes near  $a, b$  hence  $M^n f$  vanishes near  $a, b$  for any  $n \in \mathbb{Z}_{\geq 0}$ . Therefore

$$|I(\lambda)| = \left| \int_a^b fg \right| = \frac{1}{\lambda} \left| \int_a^b g \cdot Mf \right| = \cdots = \frac{1}{\lambda^n} \left| \int_a^b g \cdot M^n f \right|.$$

so  $c_n = \left| \int_a^b g \cdot M^n f \right|$  is valid.

**C13)** If there exists  $\delta > 0$ , such that for any  $x \in [a, b]$ ,  $|\varphi'(x)| \geq \delta$  and  $\varphi'(x)$  is monotonic on  $[a, b]$ . Prove that there is a constant  $C_1$  independent of  $\lambda, \varphi, a, b$  such that

$$\left| \int_a^b e^{i\lambda\varphi(x)} dx \right| \leq \frac{C_1}{\lambda\delta}.$$

Proof: Let  $C_1 = 4$  then (since  $\varphi'$  maintains the same sign)

$$\begin{aligned} \left| \int_a^b e^{i\lambda\varphi(x)} dx \right| &= \left| \int_a^b \frac{de^{i\lambda\varphi}}{\lambda\varphi'} \right| \leq \left| \frac{e^{i\lambda\varphi}}{\lambda\varphi'} \right|_a^b + \frac{1}{\lambda} \left| \int_a^b e^{i\lambda\varphi} \frac{\varphi''}{(\varphi')^2} dx \right| \\ &\leq \frac{2}{\lambda\delta} + \frac{1}{\lambda} \int_a^b \left| \frac{\varphi''}{(\varphi')^2} \right| \\ &= \frac{2}{\lambda\delta} + \frac{1}{\lambda} \int_a^b d \frac{1}{\varphi'} \leq \frac{4}{\lambda\delta}. \end{aligned}$$

**C14)** Suppose for any  $x \in [a, b]$ ,  $|\varphi''(x)| \geq 1$ . Prove that there is a unique  $c \in [a, b]$  such that

$$|\varphi'(c)| = \inf_{x \in [a, b]} |\varphi'(x)|.$$

Further prove that for any  $x \in [a, b]$ ,

$$|\varphi'(x)| \geq |x - c|.$$

Proof: Since  $\varphi \in C^\infty([a, b])$  and  $|\varphi''| \geq 1$ ,  $\varphi''$  maintains the same sign. Assume that  $\forall x \in [a, b]$ ,  $\varphi''(x) \geq 1$ , then  $\varphi'$  is monotonically increasing. Therefore, if  $\varphi' \neq 0$ , then  $c \in \{a, b\}$ , otherwise,  $c$  is the unique null point of  $\varphi'$ .

Either  $\varphi'(c) = 0$  or  $c = a$ , when  $\varphi'$  maintains the same sign, so we always have  $|\varphi'(x)| \geq |\varphi'(x) - \varphi'(c)|$ , and

$$\forall x \in [a, b], \exists \xi \in [x, c], |\varphi'(x) - \varphi'(c)| \geq |x - c| \cdot \varphi'(\xi) \geq |x - c|.$$

Therefore  $|\varphi'(x)| \geq |x - c|$ .

**!C15)** Assume that for any  $x \in [a, b]$ ,  $|\varphi''(x)| \geq 1$ . Prove that there is a constant  $C_2$  independent of  $\lambda, \varphi, a, b$ , such that

$$\left| \int_a^b e^{i\lambda\varphi(x)} dx \right| \leq \frac{C_2}{\sqrt{\lambda}}.$$

Proof: Since  $\varphi \in C^\infty([a, b])$ , we can assume  $\varphi''(x) \geq 1$ . For an arbitrary  $\delta > 0$ , divide the interval  $[a, b]$  into two parts:

$E_1 = \{x : |\varphi'(x)| \leq \delta\}$  and  $E_2 = \{x : |\varphi'(x)| > \delta\}$ .

Note that  $\forall x, y \in E_1$ ,  $|\varphi'(x) - \varphi'(y)| \leq 2\delta$ , but  $|\varphi'(x) - \varphi'(y)| \geq \left| \int_x^y \varphi''(t) dt \right| \geq |x - y|$ . Therefore  $E_1$  is an interval of length at most  $2\delta$ , so

$$\left| \int_{E_1} e^{i\lambda\varphi(x)} dx \right| \leq 2\delta.$$

Now consider the integral on  $E_2$ , which is the union of one or two intervals. By C13),

$$\left| \int_{E_2} e^{i\lambda\varphi(x)} dx \right| \leq 2 \cdot \frac{4}{\lambda\delta}.$$

Therefore

$$\left| \int_a^b e^{i\lambda\varphi(x)} dx \right| \leq 2\delta + \frac{8}{\lambda\delta} = \frac{8}{\sqrt{\lambda}}.$$

(if we let  $\delta = \sqrt{4/\lambda}$ .)

**C16) Assume that for any  $x \in [a, b]$ ,  $|\varphi''(x)| \geq 1$ . Prove that there is a constant  $C_2$  independent of  $\lambda, \varphi, f, a, b$  such that**

$$\left| \int_a^b e^{i\lambda\varphi(x)} f(x) dx \right| \leq \frac{C_2}{\sqrt{\lambda}} \left( |f(a)| + \int_a^b |f'(x)| dx \right).$$

Proof: By C15),

$$\begin{aligned} \left| \int_a^b e^{i\lambda\varphi(x)} f(x) dx \right| &\leq \left| \int_a^b e^{i\lambda\varphi(x)} f(a) dx \right| + \left| \int_a^b e^{i\lambda\varphi(x)} \int_a^x f'(t) dt dx \right| \\ &\leq |f(a)| \frac{C_2}{\sqrt{\lambda}} + \left| \int_a^b f'(t) \int_t^b e^{i\lambda\varphi(x)} dx dt \right| \\ &\leq \frac{C_2}{\sqrt{\lambda}} \left( |f(a)| + \int_a^b |f'(x)| dx \right). \end{aligned}$$