


Guideline Title: Hypoglycemia

Guideline Number: GUID-3203-M021

Issue date: 09/09/2014	Replaces Dept. Policy:
Revision dates: 4/30/24	Developed by: Air Care Team
Approved by: Dr. Christopher Hunter, MD Air Care Team Medical Director	Approved by:
Signature: 	Signature:
Department Numbers: 3203	

- I. **PURPOSE:** Recognition of the patient who hypoglycemic and treat according to condition with goal to return blood glucose to a minimum of 60 mg/dL without complications.
- II. **DEFINITIONS:**
- III. **DEPARTMENT GUIDELINE:**
 - a. Establish care as defined by *GUID-G002 – General Patient Care*.
 - b. Assess for - Subjective signs: altered mental status, dizziness, fatigue, blurred vision, headache. Physical assessment: seizure, drowsiness, unconsciousness, personality changes, sweating, staggering.
 - c. Check a blood glucose level per *GUID-PR003 – Blood Glucose Monitoring*. If patient is hypothermic, hypotensive, on multiple vasopressors, or is in shock, obtain blood via IV line.
 - d. Review all diabetes medication and last dose. Review nutritional status.
 - e. Establish a peripheral intravenous catheter.
 - f. If patient is symptomatic and
 - i. Adult
 1. If glucose < 60 mg/dL:
 - a. **Dextrose 50% 25 grams (1 amp) IV/IO -OR-**
 - b. **Dextrose 10% 10 grams (100 ml) IV/IO** every 5 minutes to a maximum dose of 250ml
 - c. If no IV/IO access, **Glucagon 1 mg SQ/IM -OR-**
 - d. If the patient is a known diabetic with hypoglycemia, is allowed and able to drink, and is not being transferred for an interventional/surgical procedure or is an acute trauma or stroke patient give 15 grams of an **oral glucose product: Liquid Dex-4**.
 - ii. Pediatric with blood glucose level is < 60 mg/dL
 1. **Dextrose 10% 0.2-1 g/kg IV/IO (2-10 ml/kg) -OR-**
 2. **Dextrose 25% 2 ml/kg IV/IO -OR-**
 3. **Glucagon < 20 kg: 0.5 mg SC/IM or if > 20 kg: 1 mg SC/IM**

Guideline Title: **Hypoglycemia**

Guideline Number: **GUID-3203-M021**

iii. Infant/neonate <3mo

1. **Dextrose 5% 0.5-1 g/kg IV** (10-20 ml/kg)
2. **D10 2-4 ml/kg IV**
3. **If < 5 kg: Glucagon 0.25 mg IM**

g. Repeat blood glucose level 15 minutes after initial intervention and every 15 minutes thereafter until level is > 60 mg/dL.

IV. **DOCUMENTATION:**

- a. EMS Charts

V. **REFERENCES:**

- a. Orlando Health Patient Care Policy and Procedure #5115: Hypoglycemia Treatment: Adult and Pediatric Inpatients
- b. American Diabetes Association. (2019). Standard of medical care in diabetes- 2019. Diabetes Care, 42(1).