

Guideline Title: Heat Related Illness/Hyperthermia

Guideline Number: GUID-3203-M009

Issue date: 09/09/2014	Replaces Dept. Policy:
Revision dates: 4/30/24	Developed by: Air Care Team
Approved by: Dr. Christopher Hunter, MD Air Care Team Medical Director	Approved by:
Signature: 	Signature:
Department Numbers: 3203	

- I. **PURPOSE:** Therapeutic goal: To recognize the patient with heat related illness and to provide and maintain adequate oxygenation/ventilation and tissue perfusion. To provide adequate temperature control and expeditious transport to the nearest appropriate medical facility.
- II. **DEFINITIONS:**
 - a. BP – Blood Pressure
 - b. MAP – Mean Arterial Pressure
- III. **DEPARTMENT GUIDELINE:**
 - a. Establish care as defined by GUID3203-G002 – *General Patient Care*.
 - b. Remove the patient from the current environment and place in an area that is cool.
 - c. Provide supplemental oxygen.
 - d. Keep patient supine.
 - e. Establish 2 large bore IV access
 - f. If the patient is hypotensive
 - i. Adults: 0.9% NS IV/IO fluid boluses per GUID3203-M011 - *Hypotension*.
 - ii. Pediatric: 0.9% NS 10-20 ml/kg IV/IO fluid boluses per GUID3203-PED009 – *Pediatric Hypotension*.
 - g. Actively cool the patient: remove excess clothing, apply cool water to skin, increase movement of air over the patient, and/or apply cool packs to groin and axilla if available.
 - h. Perform a bedside glucose check and if glucose is <60 mg/dL and the patient is symptomatic, treat per GUID3203-M021 – *Hypoglycemia*.
 - i. If cocaine/sympathomimetic toxicity strongly suspected with signs of hypertension, tachycardia, or agitation consider **Midazolam 0.05-0.1 mg/kg IV/IO**, may repeat every 3-5 minutes PRN, maximum dose 5 mg
 - j. If seizures occur, treat per GUID3203-M015 – *Seizures*.
- IV. **DOCUMENTATION:**
 - a. EMS Charts
- V. **REFERENCES:**