


Guideline Title: Management of Previously Initiated & Continuous IV Medication

Guideline Number: GUID-3203-G003

Issue date: 09/09/2014	Replaces Dept. Policy:
Revision dates: 4/30/24	Developed by: Air Care Team
Approved by: Dr. Christopher Hunter, MD Air Care Team Medical Director	Approved by:
Signature: 	Signature:
Department Numbers: 3203	

I. **PURPOSE:** To maximize the quality of care delivered to patients during transport.

II. **DEPARTMENT GUIDELINE:**

- A. Determine the recent, current, or anticipated use of IV medications. The team should evaluate the medications the patient is receiving and determine the reasons the referring physician is utilizing the medication. Identify any parameters that the physician has been utilizing in the use of these medications including pain scores, blood pressure, heart rates or sedation levels.
- B. Consider continuing the medication as per the referring physician's guidelines unless contraindicated by changes in the patient condition, discovery of adverse reactions, or other contraindications. The transport team should evaluate whether the medication is being given at the correct dose and route. Verify concentration, drip rates, and dosage of all previously mixed continuous medications prior to transport.
- C. Consider discontinuing the medication if they identify potential or actual problems as identified in the above section.
- D. Contact Medical Control for consultation at any time to review the medication regimes in question.

III. **DOCUMENTATION:**

- A. EMS Charts

IV. **REFERENCES:** None