

**Guideline Title: Tranexamic Acid for Severe Hemorrhage
in Trauma Patients**

Guideline Number: GUID-3203-T016

Issue date: 06/06/2016	Replaces Dept. Policy: GUID-3203-5016
Revision dates: 07/01/2026	Developed by: Air Care Team
Approved by: Dr. Danielle DiCesare, MD Air Care Team Medical Director	Approved by:
Signature: 	Signature:
Department Numbers: 3203	

- I. **PURPOSE:** This department guideline outlines the administration of Tranexamic Acid (TXA) by the Air Care Team for trauma patients, with indications of severe hemorrhage from blunt or penetrating trauma. TXA is an antifibrinolytic agent that blocks plasminogen from converting to plasmin, an enzyme that dissolves blood clots. It works to stabilize clots that have formed at the site of injury.
- II. **DEFINITIONS:**
 - a. TXA - Tranexamic Acid
 - b. SBP – Systolic blood pressure
- III. **DEPARTMENT GUIDELINE:**
 - a. Indications: TXA is given to trauma patients with suspected hemorrhagic shock as a result of a penetrating or blunt injury as evidence by:
 - i. Patients ≥ 16 years old
 1. Sustained SBP < 90 mmHg
 2. Sustained heart rate of greater than 110 bpm
 3. Initiation of *Massive Transfusion Protocol* at receiving facility
 - ii. Pediatric patients ≤ 15 years old
 1. Sustained SBP of < 70 mmHg + (age in years x 2) for children < 10 years old and SBP < 90 mmHg for children > 10 years old
 2. Sustained tachycardia appropriate for clinical context and age
 - a. Neonates (less than 28 days) < 100 bpm or > 180 bpm
 - b. Infant (1 month – 1 year) < 80 bpm or > 160 bpm
 - c. Child (1 – 10 years) < 60 bpm or > 130 bpm
 - d. Older Child (> 10 years) < 60 bpm or > 110 bpm
 3. Obvious signs of sustained hemorrhagic shock and hypoperfusion (pallor, peripheral vasoconstriction including cool, pale skin and delayed capillary refill > 2 seconds) after 20ml/kg of IV crystalloid infusion.
 - b. TXA must be started within 3 hours from time of injury with best results occurring if started within 1 hour from time of injury.

**Guideline Title: Tranexamic Acid for Severe Hemorrhage
in Trauma Patients**

Guideline Number: GUID-3203-T016

- c. Contraindications:
 - i. Known hypersensitivity
 - ii. History of known thromboembolic disease such as a deep vein thrombosis (DVT) or pulmonary embolism (PE).
 - iii. Pregnancy with fetus of viable gestational age
 - iv. Patients who have contraindication to anti-fibrinolytic therapy agents
- d. Establish care as defined by *GUID3203-G002 – General Patient Care* or *GUID3203-PED001 – Pediatric General Management*.
 - i. Patients ≥ 16 years old
 - 1. **TXA 2 grams IV/IO** slow IV push.
 - 2. Continue any maintenance infusion from a sending facility or consider a maintenance infusion if bolus was previously received: Typically, TXA 1 gram in 250ml or 500ml 0.9% NS over 8 hours.
 - ii. Pregnant patients with severe post-partum maternal hemorrhage
 - 1. **TXA 1 gram mixed in 100ml of 0.9% NS IV/IO** over 10 minutes or 600ml/hour.
 - iii. Pediatric patients ≤ 15 years old
 - 1. **TXA 15 mg/kg mixed in 10mg/ml of 0.9% NS IV/IO** (maximum 1 gram/100ml) over 10 minutes. Do not give IV push (more rapidly than 10 mL/min) due to potential for hypotension.
- e. Administration and patient care
 - i. If blood products are infusing, use an alternative IV site for TXA administration.
 - ii. Monitor for side effects: seizure, vision changes, renal impairment, and hypotension with rapid injection
 - iii. Do not delay transport; TXA may be given en route to trauma center/receiving facility.
 - iv. Provide a radio and verbal report to emergency department staff and trauma physician. Notify receiving facility of time the bolus was given and need to start infusion or time the bolus was given and the time the maintenance infusion was started.

IV. DOCUMENTATION:

- a. EMS Charts patient care record

V. REFERENCES:

- a. CRASH-2 Trial Collaborators, Roberts et al., (2010) Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in traumatic patients with significant hemorrhage (CRASH-2): a randomized placebo-controlled trial. *Lancet* 376:23.
- b. Joint Theater Trauma System Clinical Practice Guideline for Damage-Control Resuscitation http://www.usaasr.amedd.army.mil/assets/cpgs/Damage_Control_Resuscitation_10_Aug_2011%20pdf
- c. Levy, J. H., Koster, A., Quinones, Q. J., Milling, T. J., & Key, N. S. (2018). Antifibrinolytic Therapy and Perioperative Considerations. *Anesthesiology*, 128(3), 657–670. <https://doi.org/10.1097/ALN.0000000000001997>
- d. Napolitano et al., (2013). Tranexamic acid in trauma: How should we use it? *J Trauma Acute Care Surg*, 74(6), 1575-1586.