

Background & Motivation

**Clinical Context:** Cytokine Release Syndrome (CRS) is an important safety concern for immunotherapies such as CD3×CD20 bispecific antibodies, including epcoritamab. Large pharmacovigilance databases (FAERS, EudraVigilance, JADER) capture real-world safety signals at scale, but the data are often heterogeneous, incomplete, and noisy, and therefore are primarily suited for exploratory signal detection requiring careful clinical interpretation.

**Problem:** Traditional drug–AE analysis pipelines are slow, difficult to scale across oncology agents, and often generate outputs that are not easily interpretable by clinicians. As a result, rare but clinically important CRS signals may be overlooked.

**Goal:** Develop a generalized and explainable pharmacovigilance pipeline capable of analyzing any drug–AE pair, using epcoritamab and CRS as the motivating case study. Our system integrates anomaly detection, disproportionality statistics, and interpretable ML to produce clinician outputs.

**Primary Databases:** FAERS, EudraVigilance, JADER

Causal Inference and NLP Methods for Multi-Database CRS Pharmacovigilance in Epcoritamab

**Overview:** In this task, we developed unified framework combining causal inference and NLP-driven feature extraction to characterize CRS risk across multiple pharmacovigilance databases. The goal is to quantify the effect of key treatments and clinical factors while leveraging unstructured narratives to improve severe CRS detection.

**Methods:**

1. Causal Inference


- DAG separating exposure (dose), confounders (age, disease stage, prior therapies), effect modifiers (steroids, tocilizumab), and colliders.
- Association tests: odds ratios, p-values for dose, steroids, tocilizumab, co-medications.
- Propensity score modeling to estimate causal effect of steroids on severe CRS.
- Sensitivity analysis using E-values for robustness to unmeasured confounding.

2. NLP & Narrative Features

- Rule-based extraction: ICU/hypotension, intubation, vasopressors.
- Identify steroid/tocilizumab administration, CRS grade, time-to-onset.
- Classifier predicts severe CRS using narrative-derived features.

3. Key Data Summary

- **Propensity score (steroids).** Treated (n=269) vs. controls (n=80); analysis suggests a lower risk of severe CRS among steroid-treated patients, with a wide confidence interval consistent with a protective association.
- **NLP (FAERS narratives).** Analysis of 536 narratives shows limited text detail; severe CRS is identified in 31% of reports, and classifier performance remains near baseline due to sparse narrative information.



Var	p	OR/SD	Interp
Steroids	0.0024	Sig.	Prot. (causal)
Tocilizumab	0.0267	Sig.	Prot. (IL-6)
Max dose (mg)	0.0277	–	Dose-resp.
Weight	0.0010	Sig.	Conf.
Rituximab	0.0007	Sig.	Corr. (line marker)
Data source	0.0043	–	Heterog.
Age	0.053	1.09	Borderline
Sex (M)	0.065	0.66	Borderline
Co-meds count	0.116	1.78	Corr. (NS)
# doses	0.227	1.28	Likely causal (NS)

Table 1. Statistical Association Analysis (Severe CRS)

Scalable Survival Analysis for Epcoritamab-Associated CRS

**Objective:** We developed a generalized, parameterized survival analysis pipeline that can evaluate any drug–adverse event combination. Using epcoritamab–CRS as the case study, we identify real-world CRS risk factors and characterize temporal onset patterns.

**Methods:** Cox proportional hazards models were fitted on FAERS 2022–2024 epcoritamab reports with time-to-CRS as the outcome; predictors included age, weight, sex, polypharmacy, and hospitalization, and weight-based subgroups were compared descriptively.

Variable	HR	95% CI	p-value
Weight (per kg)	0.992	0.985–1.000	0.037*
Age (per year)	0.995	0.984–1.006	0.347
Polypharmacy (≥3)	0.616	0.153–2.482	0.495
Prior hospitalization	1.123	0.892–1.413	0.321

Table 2. Cox Proportional Hazards Model Results

Weight Category	Patients	CRS Rate	Clinical Action
<60 kg	148	42.6%	High risk
60–80 kg	404	30.7%	Moderate risk
>80 kg	252	28.6%	Lower risk

Table 3. Weight-Based CRS Risk Stratification

Key Findings & Clinical Implications:

- Body weight is statistically associated with CRS risk (HR = 0.992 per kg, p = 0.037), with lower weight associated with higher observed CRS risk.
- Patients <60 kg show a higher observed CRS rate than those >80 kg (42.6% vs. 28.6%); this finding should be interpreted cautiously given potential confounding.
- All reported CRS events occur within the first 24 hours of epcoritamab, defining a short, high-risk monitoring window.

Rare & Unexpected Signal Detection

**Overview:** Traditional pharmacovigilance methods often miss rare adverse events (AEs) that emerge post-market, particularly in oncology where novel drug-AE relationships may be unexpected, while we developed an automated pipeline that proactively identifies statistically significant yet clinically unexpected drug–AE signals from large-scale surveillance data, using machine learning anomaly detection with multi-step statistical filtering.

**Key Finding:** 1,386 validated rare signals with associated risk factors were discovered across 37 oncology drugs




Figure 1. 4-Step Filtering Pipeline: From raw data to rare signals

**Workflow:** Our pipeline combines Isolation Forest anomaly detection with 4-step filtering (FDA label removal, indication term removal, frequency filtering) and validates signals using disproportionality metrics.

<b>Drug–Event:</b> Epcoritamab + Haemorrhagic gastroenteritis	<b>Status:</b> RARE & UNEXPECTED	
	<b>Report Count:</b> 1	
	<b>Statistical Metrics:</b>	
	PRR (Proportional Reporting Ratio): 111.69 (threshold: >2) IC025 (Information Component): 3.602 (threshold: >0) Chi-square: 41.14 (threshold: >4)	
<b>Clinical Impact:</b>	<b>Death Rate:</b> 100.0%	
	<b>Hospitalization Rate:</b> 100.0%	
	<b>Serious Rate:</b> 100.0%	
	<b>Assessment:</b>	
<b>CONCLUSION:</b> RARE & UNEXPECTED (passed IF + all 3 statistical tests)	<b>Assessment:</b>	
	In No-Cap Results: Yes Passes PRR Test (>2): Yes Passes IC025 Test (>0): Yes Passes Chi² Test (>4): Yes Passes All 3 Tests: Yes	

<b>Drug–Event:</b> Epcoritamab + Renal impairment	<b>Status:</b> NOT RARE/UNEXPECTED	
	<b>Report Count:</b> 8	
	<b>Statistical Metrics:</b>	
	PRR (Proportional Reporting Ratio): 2.84 (threshold: >2) IC025 (Information Component): 0.562 (threshold: >0) Chi-square: 7.68 (threshold: >4)	
<b>Clinical Impact:</b>	<b>Death Rate:</b> 137.5%	
	<b>Hospitalization Rate:</b> 175.0%	
	<b>Serious Rate:</b> 100.0%	
	<b>Assessment:</b>	
<b>CONCLUSION:</b> NOT RARE/UNEXPECTED (did not pass Isolation Forest)	<b>Assessment:</b>	
	In No-Cap Results: No Passes PRR Test (>2): Yes Passes IC025 Test (>0): Yes Passes Chi² Test (>4): Yes Passes All 3 Tests: Yes	