

# On Almost Lyapunov Functions for Systems with Inputs

Shenyu Liu, Daniel Liberzon

Coordinated Science Laboratory,  
University of Illinois at Urbana-Champaign, IL, USA

*{sliu113, liberzon}@illinois.edu*

December 11th, 2019

# Overview

- 1 Introduction
- 2 Autonomous systems
- 3 Systems with inputs
- 4 Main result
- 5 Example
- 6 Conclusion

# Introduction

- Lyapunov's direct method tells that a time invariant, autonomous system

$$\dot{x} = f(x) \quad (1)$$

has a globally asymptotically stable origin if there exists a **positive definite**, radially unbounded function  $V(x)$  such that  $\dot{V}(x) := \langle \frac{d}{dx} V(x), f(x) \rangle$  is **negative definite** [Khalil, 2002].

- Finding such  $V$  satisfying the opposite sign definite constraints is difficult.
- **Non-monotonic** Lyapunov functions are widely studied [Aeyels and Peuteman, 1998, Ahmadi and Parrilo, 2008, Karafyllis, 2011, Meigoli and Nikraves, 2012]
- The analysis complicates with the presence of inputs;  $\dot{x} = f(x, u)$ .

- Lyapunov's direct method tells that a time invariant, autonomous system

$$\dot{x} = f(x) \quad (1)$$

has a globally asymptotically stable origin if there exists a **positive definite**, radially unbounded function  $V(x)$  such that  $\dot{V}(x) := \langle \frac{d}{dx} V(x), f(x) \rangle$  is **negative definite** [Khalil, 2002].

- Finding such  $V$  satisfying the opposite sign definite constraints is difficult.
- **Non-monotonic** Lyapunov functions are widely studied [Aeyels and Peuteman, 1998, Ahmadi and Parrilo, 2008, Karafyllis, 2011, Meigoli and Nikraves, 2012]
- The analysis complicates with the presence of inputs;  $\dot{x} = f(x, u)$ .

- Lyapunov's direct method tells that a time invariant, autonomous system

$$\dot{x} = f(x) \quad (1)$$

has a globally asymptotically stable origin if there exists a **positive definite**, radially unbounded function  $V(x)$  such that  $\dot{V}(x) := \langle \frac{d}{dx} V(x), f(x) \rangle$  is **negative definite** [Khalil, 2002].

- Finding such  $V$  satisfying the opposite sign definite constraints is difficult.
- **Non-monotonic** Lyapunov functions are widely studied [Aeyels and Peuteman, 1998, Ahmadi and Parrilo, 2008, Karafyllis, 2011, Meigoli and Nikraves, 2012]
- The analysis complicates with the presence of inputs;  $\dot{x} = f(x, u)$ .

- Lyapunov's direct method tells that a time invariant, autonomous system

$$\dot{x} = f(x) \quad (1)$$

has a globally asymptotically stable origin if there exists a **positive definite**, radially unbounded function  $V(x)$  such that  $\dot{V}(x) := \langle \frac{d}{dx} V(x), f(x) \rangle$  is **negative definite** [Khalil, 2002].

- Finding such  $V$  satisfying the opposite sign definite constraints is difficult.
- **Non-monotonic** Lyapunov functions are widely studied [Aeyels and Peuteman, 1998, Ahmadi and Parrilo, 2008, Karafyllis, 2011, Meigoli and Nikraves, 2012]
- The analysis complicates with the presence of inputs;  $\dot{x} = f(x, u)$ .

“ $V(x(t))$  decreases more than it increases over any long enough time”  
which can be guaranteed by

- 1 Considering the finite step difference of Lyapunov function, i.e., existence of  $T > 0$  such that  $V(x(T)) - V(x(0)) < 0$  for all  $x(0) \neq 0$  [Aeyels and Peuteman, 1998]; or
- 2 Imposing some conditions on the higher order derivatives of  $V$  [Meigoli and Nikraves, 2009]; or
- 3 The study of almost Lyapunov functions such that  $\Omega := \{x \in \mathbb{R}^n : \dot{V}(x) > -aV(x)\}$  is small enough [Liu et al., 2016], [Liu et al., 2019].



“ $V(x(t))$  decreases more than it increases over any long enough time”  
which can be guaranteed by

- ① Considering the finite step difference of Lyapunov function, i.e., existence of  $T > 0$  such that  $V(x(T)) - V(x(0)) < 0$  for all  $x(0) \neq 0$  [Aeyels and Peuteman, 1998]; or
- ② Imposing some conditions on the higher order derivatives of  $V$  [Meigoli and Nikraves, 2009]; or
- ③ The study of almost Lyapunov functions such that  $\Omega := \{x \in \mathbb{R}^n : \dot{V}(x) > -aV(x)\}$  is small enough [Liu et al., 2016], [Liu et al., 2019].

“ $V(x(t))$  decreases more than it increases over any long enough time”  
which can be guaranteed by

- 1 Considering the finite step difference of Lyapunov function, i.e., existence of  $T > 0$  such that  $V(x(T)) - V(x(0)) < 0$  for all  $x(0) \neq 0$  [Aeyels and Peuteman, 1998]; or
- 2 Imposing some conditions on the higher order derivatives of  $V$  [Meigoli and Nikraves, 2009]; or
- 3 The study of almost Lyapunov functions such that  $\Omega := \{x \in \mathbb{R}^n : \dot{V}(x) > -aV(x)\}$  is small enough [Liu et al., 2016], [Liu et al., 2019].

“ $V(x(t))$  decreases more than it increases over any long enough time”  
which can be guaranteed by

- 1 Considering the finite step difference of Lyapunov function, i.e., existence of  $T > 0$  such that  $V(x(T)) - V(x(0)) < 0$  for all  $x(0) \neq 0$  [Aeyels and Peuteman, 1998]; or
- 2 Imposing some conditions on the higher order derivatives of  $V$  [Meigoli and Nikraves, 2009]; or
- 3 The study of almost Lyapunov functions such that  $\Omega := \{x \in \mathbb{R}^n : \dot{V}(x) > -aV(x)\}$  is small enough [Liu et al., 2016], [Liu et al., 2019].

“ $V(x(t))$  decreases more than it increases over any long enough time”  
which can be guaranteed by

- ① Considering the finite step difference of Lyapunov function, i.e., existence of  $T > 0$  such that  $V(x(T)) - V(x(0)) < 0$  for all  $x(0) \neq 0$  [Aeyels and Peuteman, 1998]; or
- ② Imposing some conditions on the higher order derivatives of  $V$  [Meigoli and Nikraves, 2009]; or
- ③ The study of almost Lyapunov functions such that  $\Omega := \{x \in \mathbb{R}^n : \dot{V}(x) > -aV(x)\}$  is small enough [Liu et al., 2016], [Liu et al., 2019].

# Autonomous systems

# Global asymptotic stability

$$\dot{x} = f(x) \quad (1)$$

- The system (1) is **globally asymptotically stable** (GAS) if there exists  $\beta \in \mathcal{KL}$  such that

$$|x(t)| \leq \beta(|x_0|, t) \quad \forall x_0 \in \mathbb{R}^n, t \geq 0. \quad (2)$$

- GAS can be shown via a Lyapunov function  $V(x) : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$  such that for some  $\alpha_1, \alpha_2 \in \mathcal{K}_\infty, a > 0$  such that

$$\alpha_1(|x|) \leq V(x) \leq \alpha_2(|x|) \quad \forall x \in \mathbb{R}^n, \quad (3)$$

$$\dot{V}(x) = \left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad \forall x \in \mathbb{R}^n. \quad (4)$$

# Global asymptotic stability

$$\dot{x} = f(x) \quad (1)$$

- The system (1) is **globally asymptotically stable** (GAS) if there exists  $\beta \in \mathcal{KL}$  such that

$$|x(t)| \leq \beta(|x_0|, t) \quad \forall x_0 \in \mathbb{R}^n, t \geq 0. \quad (2)$$

- GAS can be shown via a Lyapunov function  $V(x) : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$  such that for some  $\alpha_1, \alpha_2 \in \mathcal{K}_\infty, a > 0$  such that

$$\alpha_1(|x|) \leq V(x) \leq \alpha_2(|x|) \quad \forall x \in \mathbb{R}^n, \quad (3)$$

$$\dot{V}(x) = \left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad \forall x \in \mathbb{R}^n. \quad (4)$$

# Global asymptotic stability

$$\dot{x} = f(x) \quad (1)$$

- The system (1) is **globally asymptotically stable** (GAS) if there exists  $\beta \in \mathcal{KL}$  such that

$$|x(t)| \leq \beta(|x_0|, t) \quad \forall x_0 \in \mathbb{R}^n, t \geq 0. \quad (2)$$

- GAS can be shown via a Lyapunov function  $V(x) : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$  such that for some  $\alpha_1, \alpha_2 \in \mathcal{K}_\infty, a > 0$  such that

$$\alpha_1(|x|) \leq V(x) \leq \alpha_2(|x|) \quad \forall x \in \mathbb{R}^n, \quad (3)$$

$$\dot{V}(x) = \left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad \forall x \in \mathbb{R}^n. \quad (4)$$



# Almost Lyapunov functions for autonomous systems

$$\left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad (4)$$

- The inequality (4) may not hold for all  $x \in \mathbb{R}^n$ , because of:
  - 1 Poor choice of  $V$ ;
  - 2 Numerical computation;
  - 3 System perturbation;
  - 4 ...
- The inequality (4) may not be satisfied for  $x \in \Omega \subset \mathbb{R}^n$ .
- Our previous work [Liu et al., 2019] shows that as long as the volume of the connected components of  $\Omega$  are small enough, the system (1) is GAS.

# Almost Lyapunov functions for autonomous systems


$$\left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad (4)$$

- The inequality (4) may not hold for all  $x \in \mathbb{R}^n$ , because of:
  - 1 Poor choice of  $V$ ;
  - 2 Numerical computation;
  - 3 System perturbation;
  - 4 ...
- The inequality (4) may not be satisfied for  $x \in \Omega \subset \mathbb{R}^n$ .
- Our previous work [Liu et al., 2019] shows that as long as the volume of the connected components of  $\Omega$  are small enough, the system (1) is GAS.

# Almost Lyapunov functions for autonomous systems

$$\left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad (4)$$


- The inequality (4) may not hold for all  $x \in \mathbb{R}^n$ , because of:
  - 1 Poor choice of  $V$ ;
  - 2 Numerical computation;
  - 3 System perturbation;
  - 4 ...
- The inequality (4) may not be satisfied for  $x \in \Omega \subset \mathbb{R}^n$ .
- Our previous work [Liu et al., 2019] shows that as long as the volume of the connected components of  $\Omega$  are small enough, the system (1) is GAS.



# Almost Lyapunov functions for autonomous systems

$$\left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad (4)$$


- The inequality (4) may not hold for all  $x \in \mathbb{R}^n$ , because of:
  - 1 Poor choice of  $V$ ;
  - 2 Numerical computation;
  - 3 System perturbation;
  - 4 ...
- The inequality (4) may not be satisfied for  $x \in \Omega \subset \mathbb{R}^n$ .
- Our previous work [Liu et al., 2019] shows that as long as the volume of the connected components of  $\Omega$  are small enough, the system (1) is GAS.



# Almost Lyapunov functions for autonomous systems

$$\left\langle \frac{d}{dx} V(x), f(x) \right\rangle \leq -aV(x) \quad (4)$$

- The inequality (4) may not hold for all  $x \in \mathbb{R}^n$ , because of:
  - 1 Poor choice of  $V$ ;
  - 2 Numerical computation;
  - 3 System perturbation;
  - 4 ...
- The inequality (4) may not be satisfied for  $x \in \Omega \subset \mathbb{R}^n$ .
- Our previous work [Liu et al., 2019] shows that as long as the volume of the connected components of  $\Omega$  are small enough, the system (1) is GAS.



# Systems with inputs

# Input-to-state Stability

$$\dot{x} = f(x, u) \quad (5)$$

- The system (5) is **input-to-state stable** (ISS) [Sontag, 1989] if there exists  $\beta \in \mathcal{KL}, \gamma \in \mathcal{K}_\infty$  such that

$$|x(t; x_0, u)| \leq \beta(|x_0|, t) + \gamma(\text{ess sup}_{s \in [0, t]} |u(s)|) \quad (6)$$

for all  $x_0 \in \mathbb{R}^n, t \geq 0$  and  $u \in \mathcal{U} \subseteq \mathbb{R}^m$ .

- ISS can be shown via a Lyapunov function  $V(x) : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$  such that for some  $\alpha_1, \alpha_2, \rho \in \mathcal{K}_\infty, a > 0$ , (3) holds and

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

for all  $x \in \mathbb{R}^n, |u| \leq \rho(|x|)$  [Sontag and Wang, 1995].

# Input-to-state Stability

$$\dot{x} = f(x, u) \quad (5)$$

- The system (5) is **input-to-state stable** (ISS) [Sontag, 1989] if there exists  $\beta \in \mathcal{KL}, \gamma \in \mathcal{K}_\infty$  such that

$$|x(t; x_0, u)| \leq \beta(|x_0|, t) + \gamma(\text{ess sup}_{s \in [0, t]} |u(s)|) \quad (6)$$

for all  $x_0 \in \mathbb{R}^n, t \geq 0$  and  $u \in \mathcal{U} \subseteq \mathbb{R}^m$ .

- ISS can be shown via a Lyapunov function  $V(x) : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$  such that for some  $\alpha_1, \alpha_2, \rho \in \mathcal{K}_\infty, a > 0$ , (3) holds and

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

for all  $x \in \mathbb{R}^n, |u| \leq \rho(|x|)$  [Sontag and Wang, 1995].



$$\dot{x} = f(x, u) \quad (5)$$

- The system (5) is **input-to-state stable** (ISS) [Sontag, 1989] if there exists  $\beta \in \mathcal{KL}, \gamma \in \mathcal{K}_\infty$  such that

$$|x(t; x_0, u)| \leq \beta(|x_0|, t) + \gamma(\operatorname{ess\,sup}_{s \in [0, t]} |u(s)|) \quad (6)$$

for all  $x_0 \in \mathbb{R}^n, t \geq 0$  and  $u \in \mathcal{U} \subseteq \mathbb{R}^m$ .

- ISS can be shown via a Lyapunov function  $V(x) : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$  such that for some  $\alpha_1, \alpha_2, \rho \in \mathcal{K}_\infty, a > 0$ , (3) holds and

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

for all  $x \in \mathbb{R}^n, |u| \leq \rho(|x|)$  [Sontag and Wang, 1995].

# Problem set up

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

- Denote the solution of the **auxiliary system**

$$\dot{x} = f_\rho(x, d) := f(x, \rho(|x|)d) \quad (8)$$

by  $x_\rho(t; x_0, d)$ , where  $\|d\| \leq 1$ .

- (7) holds for all  $|u| \leq \rho(|x|)$  is equivalent to

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle \leq -aV(x). \quad (9)$$

- Appealing to our almost Lyapunov function framework, we allow (9) to be violated for  $x \in \Omega \subset \mathbb{R}^n$ .
- Can we still show ISS if we only have (9) for all  $x \in \mathbb{R}^n \setminus \Omega$ , while  $\Omega$  is “small” enough?

# Problem set up

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

- Denote the solution of the **auxiliary system**

$$\dot{x} = f_\rho(x, d) := f(x, \rho(|x|)d) \quad (8)$$

by  $x_\rho(t; x_0, d)$ , where  $\|d\| \leq 1$ .

- (7) holds for all  $|u| \leq \rho(|x|)$  is equivalent to

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle \leq -aV(x). \quad (9)$$

- Appealing to our almost Lyapunov function framework, we allow (9) to be violated for  $x \in \Omega \subset \mathbb{R}^n$ .
- Can we still show ISS if we only have (9) for all  $x \in \mathbb{R}^n \setminus \Omega$ , while  $\Omega$  is “small” enough?

# Problem set up

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

- Denote the solution of the **auxiliary system**

$$\dot{x} = f_\rho(x, d) := f(x, \rho(|x|)d) \quad (8)$$

by  $x_\rho(t; x_0, d)$ , where  $\|d\| \leq 1$ .

- (7) holds for all  $|u| \leq \rho(|x|)$  is equivalent to

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle \leq -aV(x). \quad (9)$$

- Appealing to our almost Lyapunov function framework, we allow (9) to be violated for  $x \in \Omega \subset \mathbb{R}^n$ .
- Can we still show ISS if we only have (9) for all  $x \in \mathbb{R}^n \setminus \Omega$ , while  $\Omega$  is “small” enough?

# Problem set up

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

- Denote the solution of the **auxiliary system**

$$\dot{x} = f_\rho(x, d) := f(x, \rho(|x|)d) \quad (8)$$

by  $x_\rho(t; x_0, d)$ , where  $\|d\| \leq 1$ .

- (7) holds for all  $|u| \leq \rho(|x|)$  is equivalent to

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle \leq -aV(x). \quad (9)$$

- Appealing to our almost Lyapunov function framework, we allow (9) to be violated for  $x \in \Omega \subset \mathbb{R}^n$ .
- Can we still show ISS if we only have (9) for all  $x \in \mathbb{R}^n \setminus \Omega$ , while  $\Omega$  is “small” enough?

# Problem set up

$$\left\langle \frac{d}{dx} V(x), f(x, u) \right\rangle \leq -aV(x) \quad (7)$$

- Denote the solution of the **auxiliary system**

$$\dot{x} = f_\rho(x, d) := f(x, \rho(|x|)d) \quad (8)$$

by  $x_\rho(t; x_0, d)$ , where  $\|d\| \leq 1$ .

- (7) holds for all  $|u| \leq \rho(|x|)$  is equivalent to

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle \leq -aV(x). \quad (9)$$

- Appealing to our almost Lyapunov function framework, we allow (9) to be violated for  $x \in \Omega \subset \mathbb{R}^n$ .
- Can we still show ISS if we only have (9) for all  $x \in \mathbb{R}^n \setminus \Omega$ , while  $\Omega$  is “small” enough?

# How to quantify the size of $\Omega$ ?

- Although volume of  $\Omega$  in Euclidean space is used in our previous work, it is conservative.
- Small  $\Omega$  does not necessarily imply  $x_\rho(t; x_0, d)$  will stay inside  $\Omega$  for finite time.
- Instead, we directly impose assumptions on the  $\Omega$  dwell time:

$$T := \sup_{x_0 \in \Omega, \|d\| \leq 1} \inf_{t \geq 0} \{t : x_\rho(t; x_0, d) \notin \Omega\} \quad (10)$$

- Only upper bound of  $T$  is needed; depending on the size and shape of  $\Omega$  and the vector field  $f(x, u)$ , it can be estimated without computing the solutions of the system.

# How to quantify the size of $\Omega$ ?

- Although volume of  $\Omega$  in Euclidean space is used in our previous work, it is conservative.
- Small  $\Omega$  does not necessarily imply  $x_\rho(t; x_0, d)$  will stay inside  $\Omega$  for finite time.
- Instead, we directly impose assumptions on the  $\Omega$  dwell time:

$$T := \sup_{x_0 \in \Omega, \|d\| \leq 1} \inf_{t \geq 0} \{t : x_\rho(t; x_0, d) \notin \Omega\} \quad (10)$$

- Only upper bound of  $T$  is needed; depending on the size and shape of  $\Omega$  and the vector field  $f(x, u)$ , it can be estimated without computing the solutions of the system.



# How to quantify the size of $\Omega$ ?

- Although volume of  $\Omega$  in Euclidean space is used in our previous work, it is conservative.
- Small  $\Omega$  does not necessarily imply  $x_\rho(t; x_0, d)$  will stay inside  $\Omega$  for finite time.
- Instead, we directly impose assumptions on the  **$\Omega$  dwell time**:

$$T := \sup_{x_0 \in \Omega, \|d\| \leq 1} \inf_{t \geq 0} \{t : x_\rho(t; x_0, d) \notin \Omega\} \quad (10)$$

- Only upper bound of  $T$  is needed; depending on the size and shape of  $\Omega$  and the vector field  $f(x, u)$ , it can be estimated without computing the solutions of the system.

# How to quantify the size of $\Omega$ ?

- Although volume of  $\Omega$  in Euclidean space is used in our previous work, it is conservative.
- Small  $\Omega$  does not necessarily imply  $x_\rho(t; x_0, d)$  will stay inside  $\Omega$  for finite time.
- Instead, we directly impose assumptions on the  $\Omega$  dwell time:

$$T := \sup_{x_0 \in \Omega, \|d\| \leq 1} \inf_{t \geq 0} \{t : x_\rho(t; x_0, d) \notin \Omega\} \quad (10)$$

- Only upper bound of  $T$  is needed; depending on the size and shape of  $\Omega$  and the vector field  $f(x, u)$ , it can be estimated without computing the solutions of the system.

# The upper bound of time derivative of $V$

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle, \quad (11)$$

## Lemma

Let  $V \in C^1(\mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0})$  be a positive definite function and assume the system (5) has an equilibrium at 0. Then  $V'$  defined via (11) exists for all  $x \in \mathbb{R}^n$  and is Lipschitz when both  $f_\rho(x, d)$ ,  $\frac{d}{dx} V(x)$  are Lipschitz in  $x$ .

## Lemma

Assume  $\Omega$  is bounded and all the assumptions in the above lemma hold. Then  $\frac{d}{dx} V'$  exists almost everywhere in  $\Omega$ . In addition, there exists  $c > 0$  such that for all  $x \in \Omega$  where  $\frac{d}{dx} V'(x)$  exists,

$$\left| \left\langle \frac{d}{dx} V'(x), f_\rho(x, d) \right\rangle \right| \leq cV(x) \quad \forall |d| \leq 1 \quad (12)$$

# The upper bound of time derivative of $V$

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle, \quad (11)$$

## Lemma

Let  $V \in C^1(\mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0})$  be a positive definite function and assume the system (5) has an equilibrium at 0. Then  $V'$  defined via (11) exists for all  $x \in \mathbb{R}^n$  and is Lipschitz when both  $f_\rho(x, d)$ ,  $\frac{d}{dx} V(x)$  are Lipschitz in  $x$ .

## Lemma

Assume  $\Omega$  is bounded and all the assumptions in the above lemma hold. Then  $\frac{d}{dx} V'$  exists almost everywhere in  $\Omega$ . In addition, there exists  $c > 0$  such that for all  $x \in \Omega$  where  $\frac{d}{dx} V'(x)$  exists,

$$\left| \left\langle \frac{d}{dx} V'(x), f_\rho(x, d) \right\rangle \right| \leq cV(x) \quad \forall |d| \leq 1 \quad (12)$$

# The upper bound of time derivative of $V$

$$V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle, \quad (11)$$

## Lemma

Let  $V \in C^1(\mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0})$  be a positive definite function and assume the system (5) has an equilibrium at 0. Then  $V'$  defined via (11) exists for all  $x \in \mathbb{R}^n$  and is Lipschitz when both  $f_\rho(x, d)$ ,  $\frac{d}{dx} V(x)$  are Lipschitz in  $x$ .

## Lemma

Assume  $\Omega$  is bounded and all the assumptions in the above lemma hold. Then  $\frac{d}{dx} V'$  exists almost everywhere in  $\Omega$ . In addition, there exists  $c > 0$  such that for all  $x \in \Omega$  where  $\frac{d}{dx} V'(x)$  exists,

$$\left| \left\langle \frac{d}{dx} V'(x), f_\rho(x, d) \right\rangle \right| \leq cV(x) \quad \forall |d| \leq 1 \quad (12)$$

## Main result

- $f_\rho(x, d) := f(x, \rho(|x|)d)$  is Lipschitz in  $x$ ;
- $V \in C^1(\mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0})$  has Lipschitz gradient;
- $\alpha_1(|x|) \leq V(x) \leq \alpha_2(|x|) \quad \forall x \in \mathbb{R}^n$
- $V'(x) := \sup_{|d| \leq 1} \left\langle \frac{d}{dx} V(x), f_\rho(x, d) \right\rangle$ ;
- $V'(x) \leq -a V(x) \quad \forall x \in \mathbb{R}^n \setminus \Omega$ ;
- $\left| \left\langle \frac{d}{dx} V'(x), f_\rho(x, d) \right\rangle \right| \leq c V(x) \quad \forall |d| \leq 1, a.a. x \in \Omega$ ;
- $T := \sup_{x_0 \in \Omega, \|d\| \leq 1} \inf_{t \geq 0} \{t : x_\rho(t; x_0, d) \notin \Omega\}$ .

## Theorem

*Consider a system with input (5) with the assumptions listed in the previous slide satisfied with some  $a, c > 0$ . There exists an increasing function  $\alpha : [0, 1) \rightarrow [0, \infty)$  with  $\alpha(0) = 0, \lim_{t \rightarrow 1^-} \alpha(t) = \infty$  such that as long as the  $\Omega$  dwell time  $T$  satisfies*

$$T < \frac{1}{\sqrt{c}} \min \left\{ \frac{\pi}{2}, \alpha \left( \frac{a}{\sqrt{c}} \right) \right\}, \quad (13)$$

*the system (5) is ISS.*

$$\alpha(t) = \ln \left( \frac{1+t}{1-t} \right) + 2 \arccos \left( \frac{1}{\sqrt{t^2 + 1}} \right) \quad (14)$$



## Theorem

*Consider a system with input (5) with the assumptions listed in the previous slide satisfied with some  $a, c > 0$ . There exists an increasing function  $\alpha : [0, 1) \rightarrow [0, \infty)$  with  $\alpha(0) = 0, \lim_{t \rightarrow 1^-} \alpha(t) = \infty$  such that as long as the  $\Omega$  dwell time  $T$  satisfies*

$$T < \frac{1}{\sqrt{c}} \min \left\{ \frac{\pi}{2}, \alpha \left( \frac{a}{\sqrt{c}} \right) \right\}, \quad (13)$$

*the system (5) is ISS.*


$$\alpha(t) = \ln \left( \frac{1+t}{1-t} \right) + 2 \arccos \left( \frac{1}{\sqrt{t^2 + 1}} \right) \quad (14)$$

# Example

# Example

$$\dot{x} = \begin{pmatrix} -\lambda(x) & -\mu \\ \mu & -\lambda(x) \end{pmatrix} x + u, \quad (15)$$

where  $\lambda(x) = \left(\frac{a+b}{2}\right) \min\left\{\frac{|x-x_c|}{r}, 1\right\} - \frac{b}{2} + k$ .




- Take  $V = \frac{1}{2}|x|^2, \rho(s) = ks$ .  
 $V'(x) = 2(-\lambda(x) + k)V(x)$ .
- $\Omega = \{x \in \mathbb{R}^2 : |x - x_c| < r\}$ .
- For numerical values  $a = 1, b = 0.5, k = 0.1, r = 0.1, \mu = 2$  and  $x_c = (0.8, 0)^\top$ , it is computed  $c \approx 30.54$  and  $T < 0.125$ .
- (13),(14) give an upper bound of  $\Omega$  dwell time of 0.131. Hence the system (15) is ISS.

# Example

$$\dot{x} = \begin{pmatrix} -\lambda(x) & -\mu \\ \mu & -\lambda(x) \end{pmatrix} x + u, \quad (15)$$

where  $\lambda(x) = \left(\frac{a+b}{2}\right) \min\left\{\frac{|x-x_c|}{r}, 1\right\} - \frac{b}{2} + k$ .




- Take  $V = \frac{1}{2}|x|^2$ ,  $\rho(s) = ks$ .  
 $V'(x) = 2(-\lambda(x) + k)V(x)$ .
- $\Omega = \{x \in \mathbb{R}^2 : |x - x_c| < r\}$ .
- For numerical values  $a = 1$ ,  $b = 0.5$ ,  $k = 0.1$ ,  $r = 0.1$ ,  $\mu = 2$  and  $x_c = (0.8, 0)^\top$ , it is computed  $c \approx 30.54$  and  $T < 0.125$ .
- (13), (14) give an upper bound of  $\Omega$  dwell time of 0.131. Hence the system (15) is ISS.

# Example

$$\dot{x} = \begin{pmatrix} -\lambda(x) & -\mu \\ \mu & -\lambda(x) \end{pmatrix} x + u, \quad (15)$$

where  $\lambda(x) = \left(\frac{a+b}{2}\right) \min\left\{\frac{|x-x_c|}{r}, 1\right\} - \frac{b}{2} + k$ .




- Take  $V = \frac{1}{2}|x|^2$ ,  $\rho(s) = ks$ .  
 $V'(x) = 2(-\lambda(x) + k)V(x)$ .
- $\Omega = \{x \in \mathbb{R}^2 : |x - x_c| < r\}$ .
- For numerical values  $a = 1$ ,  $b = 0.5$ ,  $k = 0.1$ ,  $r = 0.1$ ,  $\mu = 2$  and  $x_c = (0.8, 0)^\top$ , it is computed  $c \approx 30.54$  and  $T < 0.125$ .
- (13), (14) give an upper bound of  $\Omega$  dwell time of 0.131. Hence the system (15) is ISS.

# Example

$$\dot{x} = \begin{pmatrix} -\lambda(x) & -\mu \\ \mu & -\lambda(x) \end{pmatrix} x + u, \quad (15)$$

where  $\lambda(x) = \left(\frac{a+b}{2}\right) \min \left\{ \frac{|x-x_c|}{r}, 1 \right\} - \frac{b}{2} + k$ .




- Take  $V = \frac{1}{2}|x|^2$ ,  $\rho(s) = ks$ .  
 $V'(x) = 2(-\lambda(x) + k)V(x)$ .
- $\Omega = \{x \in \mathbb{R}^2 : |x - x_c| < r\}$ .
- For numerical values  $a = 1$ ,  $b = 0.5$ ,  $k = 0.1$ ,  $r = 0.1$ ,  $\mu = 2$  and  $x_c = (0.8, 0)^\top$ , it is computed  $c \approx 30.54$  and  $T < 0.125$ .
- (13),(14) give an upper bound of  $\Omega$  dwell time of 0.131. Hence the system (15) is ISS.

# Example

$$\dot{x} = \begin{pmatrix} -\lambda(x) & -\mu \\ \mu & -\lambda(x) \end{pmatrix} x + u, \quad (15)$$

where  $\lambda(x) = \left(\frac{a+b}{2}\right) \min \left\{ \frac{|x-x_c|}{r}, 1 \right\} - \frac{b}{2} + k$ .



- Take  $V = \frac{1}{2}|x|^2$ ,  $\rho(s) = ks$ .  
 $V'(x) = 2(-\lambda(x) + k)V(x)$ .
- $\Omega = \{x \in \mathbb{R}^2 : |x - x_c| < r\}$ .
- For numerical values  $a = 1$ ,  $b = 0.5$ ,  $k = 0.1$ ,  $r = 0.1$ ,  $\mu = 2$  and  $x_c = (0.8, 0)^\top$ , it is computed  $c \approx 30.54$  and  $T < 0.125$ .
- (13),(14) give an upper bound of  $\Omega$  dwell time of 0.131. Hence the system (15) is ISS.

- Our analysis via almost Lyapunov function also applies when there are multiple “bad” regions  $\Omega$ .
- Because different assumption on the size of  $\Omega$  is used in this work compared with our previous work [Liu et al., 2019], The system is shown to be ISS with even much larger  $\Omega$ .
- The trade-off is that we need some prior knowledge about the  $\Omega$  dwell time. Converting spatial bounds on  $\Omega$  to temporal bounds on  $\Omega$  is non-trivial and can be further developed.



- Our analysis via almost Lyapunov function also applies when there are multiple “bad” regions  $\Omega$ .
- Because different assumption on the size of  $\Omega$  is used in this work compared with our previous work [Liu et al., 2019], The system is shown to be ISS with even much larger  $\Omega$ .
- The trade-off is that we need some prior knowledge about the  $\Omega$  dwell time. Converting spatial bounds on  $\Omega$  to temporal bounds on  $\Omega$  is non-trivial and can be further developed.

- Our analysis via almost Lyapunov function also applies when there are multiple “bad” regions  $\Omega$ .
- Because different assumption on the size of  $\Omega$  is used in this work compared with our previous work [Liu et al., 2019], The system is shown to be ISS with even much larger  $\Omega$ .
- The trade-off is that we need some prior knowledge about the  $\Omega$  dwell time. Converting spatial bounds on  $\Omega$  to temporal bounds on  $\Omega$  is non-trivial and can be further developed.

# Sketch of proof

- Let  $t_1, t_2$  be the time the solution trajectory  $x_\rho(t; x_0, d)$  enters and leaves  $\Omega$ . We want to ultimately show  $V(x_\rho(t_2; x_0, d)) \leq \eta V(x_\rho(t_1; x_0, d))$  for some  $\eta < 1$ .
- From the definition of  $V'$  in (11) and the assumption (12) on  $V'$ , we have

$$\frac{d}{dt} V(x_\rho(t; x_0, d)) \leq V'(x_\rho(t; x_0, d)),$$

$$\left| \frac{d}{dt} V'(x_\rho(t; x_0, d)) \right| \leq c V(x_\rho(t; x_0, d)) \quad a.e.$$

- Analysing differential inequality of the vector  $\begin{bmatrix} V \\ V' \end{bmatrix}$  with boundary conditions

$$V'(x_\rho(t_i; x_0, d)) \leq -a V(x_\rho(t_i; x_0, d)), i = 1, 2. \quad (16)$$

# Sketch of proof

- Let  $t_1, t_2$  be the time the solution trajectory  $x_\rho(t; x_0, d)$  enters and leaves  $\Omega$ . We want to ultimately show  $V(x_\rho(t_2; x_0, d)) \leq \eta V(x_\rho(t_1; x_0, d))$  for some  $\eta < 1$ .
- From the definition of  $V'$  in (11) and the assumption (12) on  $V'$ , we have

$$\frac{d}{dt} V(x_\rho(t; x_0, d)) \leq V'(x_\rho(t; x_0, d)),$$

$$\left| \frac{d}{dt} V'(x_\rho(t; x_0, d)) \right| \leq c V(x_\rho(t; x_0, d)) \quad a.e.$$

- Analysing differential inequality of the vector  $\begin{bmatrix} V \\ V' \end{bmatrix}$  with boundary conditions

$$V'(x_\rho(t_i; x_0, d)) \leq -a V(x_\rho(t_i; x_0, d)), i = 1, 2. \quad (16)$$

# Sketch of proof

- Let  $t_1, t_2$  be the time the solution trajectory  $x_\rho(t; x_0, d)$  enters and leaves  $\Omega$ . We want to ultimately show  $V(x_\rho(t_2; x_0, d)) \leq \eta V(x_\rho(t_1; x_0, d))$  for some  $\eta < 1$ .
- From the definition of  $V'$  in (11) and the assumption (12) on  $V'$ , we have

$$\frac{d}{dt} V(x_\rho(t; x_0, d)) \leq V'(x_\rho(t; x_0, d)),$$

$$\left| \frac{d}{dt} V'(x_\rho(t; x_0, d)) \right| \leq c V(x_\rho(t; x_0, d)) \quad a.e.$$

- Analysing differential inequality of the vector  $\begin{bmatrix} V \\ V' \end{bmatrix}$  with boundary conditions

$$V'(x_\rho(t_i; x_0, d)) \leq -a V(x_\rho(t_i; x_0, d)), i = 1, 2. \quad (16)$$

# Sketch of proof

- For any  $t \in [t_1, t_2]$ , It is proven that


$$\frac{V(x_\rho(t; x_0, d))}{V(x_\rho(t_1; x_0, d))} \leq R_1(t - t_1),$$

$$\frac{V(x_\rho(t; x_0, d))}{V(x_\rho(t_2; x_0, d))} \geq R_2(t - t_2)$$

where

$$R_1(s) := \left( \cosh \sqrt{c}s - \frac{a}{\sqrt{c}} \sinh \sqrt{c}s \right),$$

$$R_2(s) := \left( \cos \sqrt{c}s - \frac{a}{\sqrt{c}} \sin \sqrt{c}s \right).$$



- It is then shown that when (13) holds, there always exists  $t^* \in [t_1, t_2]$  such that  $R_1(t^* - t_1) \leq \eta_1 < 1$ ,  $R_2(t^* - t_2) \geq \eta_2 > 1$  and thus

$$\frac{V(x_\rho(t_2; x_0, d))}{V(x_\rho(t_1; x_0, d))} \leq \frac{R_1(t^* - t_1)}{R_2(t^* - t_2)} \leq \frac{\eta_1}{\eta_2} < 1.$$

# Sketch of proof

- For any  $t \in [t_1, t_2]$ , It is proven that


$$\frac{V(x_\rho(t; x_0, d))}{V(x_\rho(t_1; x_0, d))} \leq R_1(t - t_1),$$

$$\frac{V(x_\rho(t; x_0, d))}{V(x_\rho(t_2; x_0, d))} \geq R_2(t - t_2)$$

where

$$R_1(s) := \left( \cosh \sqrt{c}s - \frac{a}{\sqrt{c}} \sinh \sqrt{c}s \right),$$

$$R_2(s) := \left( \cos \sqrt{c}s - \frac{a}{\sqrt{c}} \sin \sqrt{c}s \right).$$



- It is then shown that when (13) holds, there always exists  $t^* \in [t_1, t_2]$  such that  $R_1(t^* - t_1) \leq \eta_1 < 1$ ,  $R_2(t^* - t_2) \geq \eta_2 > 1$  and thus

$$\frac{V(x_\rho(t_2; x_0, d))}{V(x_\rho(t_1; x_0, d))} \leq \frac{R_1(t^* - t_1)}{R_2(t^* - t_2)} \leq \frac{\eta_1}{\eta_2} < 1.$$

# Conclusion



- The study of almost Lyapunov functions from our previous work was generalized to systems with inputs.
- When there are “bad” regions  $\Omega$  in the state space where  $V$  does not decrease fast enough, an upper bound of the  $\Omega$  dwell time was found to guarantee that the system with inputs is still ISS.
- Example showed that our method of showing stability via almost Lyapunov function is able to handle non-trivial perturbations and it is believed to be applicable to a broader class of systems.



Aeyels, D. and Peuteman, J. (1998).

A new asymptotic stability criterion for nonlinear time-variant differential equations.

*IEEE Transactions on Automatic Control*, 43(7):968–971.



Ahmadi, A. A. and Parrilo, P. A. (2008).

Non-monotonic Lyapunov functions for stability of discrete time nonlinear and switched systems.

In *2008 47th IEEE Conference on Decision and Control*, pages 614–621.



Karafyllis, I. (2011).

Can we prove stability by using a positive definite function with non sign-definite derivative?

*IMA Journal of Mathematical Control and Information*, 29(2):147–170.



Khalil, H. (2002).

*Nonlinear Systems*, 3rd ed.

Prentice Hall.



Liu, S., Liberzon, D., and Zharnitsky, V. (2016).

On almost Lyapunov functions for non-vanishing vector fields.

In *2016 IEEE 55th Conference on Decision and Control (CDC)*, pages 5557–5562.



Meigoli, V. and Nikraves, S. K. Y. (2009).

A new theorem on higher order derivatives of Lyapunov functions.

*ISA Transactions*, 48(2):173 – 179.



Meigoli, V. and Nikraves, S. K. Y. (2012).

Stability analysis of nonlinear systems using higher order derivatives of Lyapunov function candidates.

*Systems & Control Letters*, 61(10):973 – 979.



Sontag, E. D. (1989).

Smooth stabilization implies coprime factorization.

*IEEE Transactions on Automatic Control*, 34(4):435–443.



Sontag, E. D. and Wang, Y. (1995).

On characterizations of the input-to-state stability property.

*Systems & Control Letters*, 24(5):351 – 359.