The continuous variable is:	Rain fall on different days in a city.	Number of customers entering a store on different days.	Number of flights landing on an airport on different days.	None		
If the observations of the daily temperature on a certain a place are: -1, -4, -7, -8 and -11 then the range is	12	- 11	10	-5		
Extreme values in a data set have a strong effect on	the median	the mode	the mean	none		
The set containing the members that are common to two sets is said to be the:	Intersection	Union	Sample	Disperse		
The empty set is also known as:	Null Set	Half Set	Ultra Set	Nicon Set		
The intersection of A and B, written and defined by	A n B	A u B	A + B	A - B		
The value calculated from population is called	Sample	Parameter	Statistics	Inference		
A collection of observation is called	Data	Sets	Primary	Secondary		
Which of the following is a branch of statistics?	Descriptive statistics	Inferential statistics	Industry statistics	Both A and B		
Find the mode of the call received on 7 consecutive day 11,13,13,17,19,23,25	11	13	17	23		
Find the median of the call received on 7 consecutive days 11,13,13,17,19,23,25	17	19	13	25		
The number of hours for which a light bulb being functional	Discrete data	Continuous data	Constant data	None of them		
The middle value of an ordered array of numbers is the	Mode	Median	Mean	Mid-Point		
The no of Bottles in Lab	Discrete data	Continuous data	Constant data	None of them		

Amount of Money spent on books	Qualitative Variable	Quantitative Variable	Theoretical Variable	Constant Variable		
The complement of A, is defined by	A	$A^{/}$	S	G		
Number of flowers on Plant	Continuous data	Constant data	None of them	Discrete data		
Histogram is a graph of	Qualitative data	Time sereis	Frequency distribution	Ogive		
The data about the weights of plants	Discrete data	Continuous data	Constant data	None of them		
The average of population is	Parameter	Statistics	Nicon	Median		
Natural numbers are:	{+1, -1}	{1,2,3,4.....}	{0,1,2,3,8,9}	None of them		
The science of collecting, organizing, presenting, analyzing and interpreting data to assist in making more effective decisions is called:	Parameter	Statistics	Population	Inference		
Methods of organizing, summarizing, and presenting data in an informative way are called:	Descriptive statistics	Inferential statistics	Theoretical statistics	Applied statistics		
A part of the population selected for study is called a:	Variable	Data	Sample	Parameter		
A set of all units of interest in a study is called:	Sample	Population	Parameter	Statistic		
A specific characteristic of a sample is called:	Statistic	Parameter	Constant	Variable		
10 is less than 15, write in inequalities sign	10 < 15	10 > 15	Neither	Both		
When the characteristic being studied is nonnumeric, it is called a	Discrete variable	Continuous variable	Quantitative Variable	Qualitative Variable		
The methods used to determine something about a population on the basis	Inferential	Descriptive	Applied	Theoretical		

of a sample is called:	statistics	statistics	statistics	statistics		
Listings of the data in the form in which these are collected are known as	Secondary data	Raw data	Arrayed data	Qualitative data		
A variable that assumes only some selected values in a range is called:	Qualitative variable	Discrete variable	Quantitative variable	Continuous variable		
A value which does not change from one individual to another individual is called	Array	Constant	Statistic	Variable		
In the plural sense, statistics means:	Numerical data	Methods	Population data	Sample data		
The median of the data 2, 3, 5, 9, 11 (odd number of values)	7	9	11	2		
Merits of Median	easy to calculate	Difficult to calculate	Moderate in difficulty	Always constant		
Demerits of Median	not based on all observation	Based on all observation	Based on mean	Based on average		
Modal class will mandatory to find	Arithmetic Mean	Median	Mode	Average		
Measures of Central Tendency	used to describe the “centre” of the data	used to describe the “end” of the data	used to describe the “start” of the data	None of them		
Importance Of Central Tendency	To find representative value	To make more concise data	To make comparisons	All of them		
4, 6, 8, 10 A.M. of this data	7	9	5	6		
The figure 22.26 rounded to one decimal place is	22.2	22.3	22.1	22		
Students divided into different groups according to their intelligence and gender will generate:	Qualitative data	Quantitative data	Continuous data	Constant		
The number of classes will depend upon	Class marks	frequency	Class interval	Class boundary		
When the classes are 40----44, 45----49, 50----54, ... the class interval is:	4	8	5	10		
When the classes are 40----44, 45----49, boundaries are	39.5 – 44.5, 44.5 – 49.5	39 – 44, 44 – 49	39.5 – 44, 44 – 49.5	39.5 – 45, 45 – 49.5		
Frequency polygon can be drawn with the help of:	Historigram	Histogram	Pie chart	Circle		

When successive mid-points in a histogram are connected by straight lines, the graph is called a:	Historigram	Histogram	Pie chart	Frequency Polygon		
The graph of the cumulative frequency distribution is called:	Histogram	Pie chart	Ogive	Bar chart		
The word ogive is also used for:	Frequency polygon	Cumulative frequency polygon	Frequency curve	Histogram		
Cumulative frequency polygon can be used for the calculation of:	A.M.	Mode	Median	Circle		
The Percentage of Hydrogen content of gases collected from sample	**Discrete variable**	**Continuous variable**	**Constant**	**None of them**		
If there are no gaps between consecutive classes, the limits are called:	Class marks	frequency	Class interval	Class boundaries		
The average of lower and upper class limits is called:	Class limit	Class mark	Class frequency	Class boundary		
The lower and upper class limits are 20 and 30, the midpoints of the	20	25	30	50		
A frequency distribution that contains a class with limits of "10 and under 20" would have a midpoint:	10	14.5	15	20.5		
If maximum value of data is 45 and minimum is 25, the range will be	20	10	15	30		
If the range will be 30, and the maximum value is 60 find minimum value?	30	15	29	28		
6 is greater than 5, use sign of inequalities	6 - 5	6 > 5	6 < 5	6 = 5		
The part of the population from which information is collected is	**Population**	**Sample**	**Parameter**	**Statistics**		
The amount of milk produced by a cow is	Discrete variable	Continuous variable	Constant	None of them		
the construction of graphs, charts, and tables, and the calculation of various measures known as	Descriptive Statistics	Inferential Statistics	Random variable	Parameter		
a+(-a) = (-a) +a =?	0	1	2	3		
The whole population average of data is	**Sample**	**Parameter**	**Statistics**	**Inference**		
The part of information data average will be	**Mode**	**Parameter**	**Statistics**	**Inference**		
Which have more values population or sample	**Population**	**Parameter**	**Statistics**	**Inference**		
The data is 2, 4, 6, 8, 10, 8 Mode will be ?	8	4	6	10		
Find the range of data whereas, Maximum 34, Minimum 33	1	2	3	4		
A value which does not change from one individual to another individual is	Array	Constant	Statistic	Variable		

In the plural sense, statistics means	Numerical data	Methods	Population data	Sample data		
The median of the data 2,3,5,9,11	7	9	11	2		
a+0 = 0+a = ?	a	B	C	D		
Demerits of Median is	not based on all observation	Based on all observation	Based on mean	Based on average		
Modal class will mandatory to find what ?	Arithmetic Mean	Median	Mode	Average		
Measures of Central Tendency is	used to describe the "centre" of the data	used to describe the "end" of the data	used to describe the "start" of the data	None of them		
Importance Of Central Tendency will be	To find representative value	To make more concise data	To make comparisons	All of them		
4, 6, 8, 10, A.M. of this data set	7	9	5	6		
The figure 22.26 rounded to one decimal place	22.2	22.3	22.1	22		
Students divided into different groups according to their intelligence and gender will generate	**Qualitative data**	**Quantitative data**	**Continuous data**	**Constant**		
The number of classes will depend upon the	Class marks	frequency	Class interval	Class boundary		
When the classes are 40----44, 45----49, 50----54, ... the class interval is	4	8	5	10		
When the classes are 40----44, 45----49, Class boundaries are	39.5 – 44.5, 44.5 – 49.5	39 – 44, 44 – 49	39.5 – 44, 44 – 49.5	39.5 – 45, 45 – 49.5		
Frequency polygon can be drawn with the help of	Historigram	Histogram	Pie chart	Circle		
When successive mid-points in a histogram are connected by straight lines, the graph is called	Historigram	Histogram	Pie chart	Frequency Polygon		
The graph of the cumulative frequency distribution is called	Histogram	Pie chart	Ogive	Bar chart		
The word ogive is also used for	Frequency polygon	Cumulative frequency polygon	Frequency curve	Histogram		
Cumulative frequency polygon can be used for the calculation of	A.M.	Mode	Median	Circle		
The Percentage of Hydrogen content of gases collected from sample	**Discrete variable**	**Continuous variable**	**Constant**	**None of them**		
If there are no gaps between consecutive classes, the limits are called	Class marks	frequency	Class interval	Class boundaries		
The average of lower and upper class limits is called	Class limit	Class mark	Class	Class		

			frequency	boundary		
The lower and upper class limits are 20 and 30, the midpoints of the	20	25	30	50		
The frequency of the mode class is also called	Modal class	Average	Moderate	Selected		
Extreme values in a data set have a strong effect on	the median	the mode	the mean	none		
The class width of the following class limits: (-5.45) — (1.95) is	3.5	7.40	- 3.5	7.41		
The class boundaries of the following class limits: (-2.5) - (4.4) are	(-2.5) - (4.4)	(-2.45) - (4.45)	(-2) - (4)	(-2.55) - (4.45)		
The class mark (mid-point) of the following class limits: (-3.5) — (-1.5) is:	1	2.5	- 2.5	-1		
The value calculated from population is called	**Sample**	**Parameter**	**Statistics**	**Inference**		
A collection of observation is called	Data	Sets	Primary	Secondary		
Statistics is a branch of	Science	Maths	Physics	Chemistry		
A + B = B + A, regarding real numbers	True	False	Both	None of them		
A survey asks participants to indicate their level of agreement with a statement using a 5-point scale (1=strongly disagree, 2=disagree, 3=neutral, 4=agree, 5=strongly agree). What scale of measurement is being used?	Ratio	Interval	Ordinal	Nominal		
https://www.qualitygurus.com/quizzes/multiple-choice-questions-on-scales-of-measurement/						