

I have my notochord but where did I leave my tail?

Elijah Lowe^{*1}, Kanchan Pavangadkar², Likit Preeyanon², C. Titus Brown^{1, 2}, Billie J. Swalla³
¹Computer Science and Engineering and ²Microbiology and Molecular Genetics
Michigan State University, East Lansing, MI 48824
³Department of Biology
University of Washington, Seattle, WA 98195-1800


Abstract

In the molgulid clade of ascidians several species have individually evolved tail loss during their larval stage of development. The notochord, one of the key features of chordates, is found in the center of the tail in most ascidian larvae. In almost all solitary ascidians, including *M. oculata*, 40 notochord cells have been found extended in the tail. However, in a closely related species, *M. occulta* (tailless), only 20 notochord cells were found, and these express *brachyury*. These two species are only 3% divergent and are able to cross hybridize. Some of the hybrid also have 20 notochord cells, however the notochord cells extend and converge in a shortened version of *M. oculata* tail. Through the use of high throughput sequencing technologies and subsequent allelotyping analysis, we are able to quantitatively investigate the express of genes associated with notochord development in the parent species and in the hybrid. Our transcriptome assemblies are freely available for download; contact ctb@msu.edu. We thank the BEACON Center and USDA NIFA for funding.


Expression profiles

Gene	Locus	M. oculata (gast.)	Hybrid (gast.)	M. occulta (neur.)	M. oculata (neur.)	Hybrid (neur.)	M. oculata (tailbud)	Hybrid (tailbud)
ATP sulfurylase/APS kinase [Ciona intestinalis]	occulata	0	0	7	0	0	0	0
ATP sulfurylase/APS kinase [Ciona intestinalis]	oculata	17	0	0	13	0	9	0
ATP sulfurylase/APS kinase [Ciona intestinalis]	oculata	117	0	0	103	0	225	104
leprecan [Ciona intestinalis]	oculata	55	0	0	75	0	73	64
leprecan [Ciona intestinalis]	oculata	33	0	0	40	0	67	29
leprecan [Ciona intestinalis]	occulata	0	23	23	0	28	0	36
transcription factor protein foxa-a [Ciona intestinalis]	oculata	37	34	0	43	17	199	141
transcription factor protein foxa-a [Ciona intestinalis]	occulata	0	3	110	0	63	0	0
merlin [Molgula tectiformis]	occulata	0	5	16	0	20	0	10
merlin [Molgula tectiformis]	occulata	0	23	29	0	6	0	18
merlin [Molgula tectiformis]	oculata	9	0	0	7	0	13	0
merlin [Molgula tectiformis]	oculata	19	0	0	86	0	41	0
fibroblast growth factor 8/17/18 [Ciona intestinalis]	oculata	4	0	0	1	0	20	0
fibroblast growth factor 9/16/20 [Ciona intestinalis]	oculata	12	0	0	12	0	49	0
noto17 [Molgula tectiformis]	oculata	24	0	0	34	0	127	0
noto17 [Molgula tectiformis]	oculata	5	0	0	8	0	39	0
noto6 [Molgula tectiformis]	occulata	0	11	11	0	30	0	29
noto6 [Molgula tectiformis]	oculata	10	0	0	18	0	3	0
noto6 [Molgula tectiformis]	oculata	8	0	0	15	0	12	0
prickle 1 [Ciona intestinalis]	occulata	0	193	88	0	90	0	57
prickle 1 [Ciona intestinalis]	oculata	56	0	0	50	0	126	0
prickle 1 [Ciona intestinalis]	oculata	19	0	0	23	0	17	0

Molgula species studied




BLAST hits



Discussion

The notochord is important for tail formation. We examined 38 genes associated with notochord development. Of 38 selected, all but 9 were found in either *M. oculata* or *M. occulta*. 25 of the remaining 29 were found in both species, while 4 were found only in *M. oculata*. In the future we plan to confirm gene presence by in situ, increase our query gene set, and examine more time points in *M. occulta*.

Method



Gene list

- 706e(kl-3)
- ATPcitrate-lyase
- ATP sulfurylase. APS kinase
- Notch
- Mt-merlin
- Mt-noto17
- Not1
- Not2
- Not3
- Not4
- Not5
- Not6
- Not7
- Notch
- beta-catenin
- beta4galactosyltransferase
- Brachyury
- col1
- netrin
- cdc45
- ezrin.radixin.moesin(ERM)-like protein
- fibrinogen-like protein
- Fibroblastgrowthfactor3/7/10/22
- Fibroblastgrowthfactor8/17/18
- Fibroblastgrowthfactor9/16/20
- leprecan
- Mt-noto6
- noto8
- noto9
- pellino
- prickle1
- rel1
- snail
- Suppressor of hairless homolog
- foxa-a
- foxd
- tropomyosin-like protein
- tyrosinephosphatase

Key work cited

- Kugler et al. 2011. Evolutionary changes in the notochord genetic toolkit: a comparative analysis of notochord genes in the ascidian Ciona and the larvacean Oikopleura. BMC Evolutionary Biology 11:21
- Imai et al. 2004. Gene expression profiles of transcription factors and signaling molecules in the ascidian embryo: towards a comprehensive understanding of gene networks. Development 131, 4047-4058
- Hotta et al. 2008. Brachyury-downstream gene sets in a chordate, Ciona intestinalis: integrating notochord specification, morphogenesis and chordate evolution. Evolution and Development 2008, 10:1, 37-51

Acknowledgements

BEACON Center, Billie Swalla, Titus Brown, Kanchan Padvangadkar, Likit Preeyanon