A Computational Cascade Model: Solar Luminosity to Epigenetic Coherence

With CUCA Formal Laws and Empirical Validation Framework

Shia Miller

*AetherLife.co * Cosmic Chaos Corporation?*

Hollidaysburg, Pennsylvania

Preprint -- June 2026 | AETHER Framework v3.0 | CUCA v4.1
Correspondence: aetherlife.co | ORCID: 0009-0000-5537-1034

Abstract

*The AETHER Solar Physics Engine formalizes an 11-layer computational cascade linking solar electromagnetic activity to integrated biological disease risk. Beginning with the F10.7 solar flux index, the model propagates physical forcing through geomagnetic, oceanographic, endocrine, cellular, and systems-biology layers, terminating in a multiplicative coherence product Gamma*Omega*rho*Sigma that drives a nonlinear disease risk function Delta-R = (1 - Gamma*Omega*rho*Sigma)^1.8 * 100%. The four terminal biological axes -- neural coherence (Gamma), gut microbiome diversity (Omega), circadian rhythm amplitude (rho), and epigenetic clock integrity (Sigma) -- share a common upstream mediator: thyroid T3 availability, itself determined jointly by iodine supply (solar-geomagnetic-oceanographic origin) and selenium conversion efficiency (nutritional, solar-independent). The CUCA architectural framework governing the cascade is formalized through three named laws: the Reptilian Constraint (tau * B = const), the Universal Spacing Law (C(x,t) = f(d)), and the Cephalopod Edge Constraint (||E(t)|| increasing -> C(x,t) decreasing). Falsifiable predictions, empirical validation requirements, and an open research agenda are presented alongside an interactive simulation implementation.*

Keywords: solar-biological coupling, thyroid coherence, epigenetic aging, circadian rhythm, gut-brain axis, AETHER framework, CUCA architecture, coherence cascade, F10.7, iodine deficiency, selenium status, T3/rT3 ratio

1. Introduction

1.1 The Disciplinary Gap

The relationship between solar electromagnetic activity and human biological health has been proposed across multiple independent research traditions -- from geomagnetic-psychiatric correlations (Persinger 1987) to solar cycle epidemiology (Halberg et al. 2000) to the iodine-oceanography literature (Delange 2002). What has been absent is a single mechanistic framework tracing the physical chain from stellar electromagnetic output to intracellular biology with explicit quantitative equations at each layer boundary.

Academic science has compartmentalized these observations across at least six disciplines that rarely communicate: solar physics, geophysics, physical oceanography, endocrinology, chronobiology, and epigenetics. The AETHER cascade is an explicit attempt to bridge these compartments with a single computational model -- not as a claim of final truth, but as a falsifiable working framework that can be progressively tested, refined, or falsified at each layer boundary.

1.2 Central Thesis

Biological coherence -- defined operationally as the capacity of a living system to maintain high-fidelity signal processing across temporal and spatial scales -- is a measurable function of the solar electromagnetic environment, mediated through a specific cascade of physical and biochemical transduction steps. The cascade has a single critical bottleneck: thyroid T3 availability, which is co-determined by two partially independent variables:

(1) Iodine supply -- solar-geomagnetic-oceanographic origin, varying on 11-year solar cycle timescales.

(2) Selenium status -- primarily nutritional, varying on dietary timescales, partially decoupled from solar forcing.

This two-variable bottleneck structure produces a falsifiable prediction: populations with adequate selenium should show weaker solar-cycle-phase correlations with metabolic disease than selenium-deficient populations, because selenium adequacy 'clamps' the T3:rT3 ratio above the critical threshold regardless of iodine fluctuation.

1.3 The CUCA Architectural Context

The AETHER cascade is a specific instantiation of the CUCA (Closed-Loop Universal Coherence Architecture) framework, which proposes that all stable biological systems are fundamentally feedback-stabilized coherence engines. CUCA formalizes three universal laws that govern the cascade's dynamics:

* **Reptilian Constraint:** tau * B = const -- the product of biological time constant (tau) and bandwidth (B) is conserved across evolutionary scales.
* **Universal Spacing Law:** C(x,t) = f(d) -- coherence at any point is a function of spatial distance from the cascade source.
* **Cephalopod Edge Constraint:** ||E(t)|| increasing implies C(x,t) decreasing -- environmental entropy degrades coherence; the relationship is nonlinear and threshold-gated.

2. The Physical Cascade (Layers 1-4)

2.1 Layer 1: Solar Luminosity -- F10.7 Index

The solar F10.7 radio flux (10.7 cm wavelength, measured at Earth) is the standard operational proxy for solar activity, correlating r = 0.97 with sunspot number while being continuously measurable regardless of observing conditions. F10.7 ranges from ~65 sfu (solar minimum) to ~220+ sfu (solar maximum), a ~3.4x variation. This variation propagates through all downstream layers.

Governing equation (Layer 1):

```
F10.7(t) = 65 + 155 * S * 1/2 * (1 - cos(2*pi*t / 11))
[sfu]
```

where S in [0,1] is the solar activity parameter, t is time in years, and T_cycle = 11 years (Schwabe cycle). The solar forcing term uses a cosine to capture the observed bell-shaped solar maximum profile. Physical basis: F10.7 correlates with EUV (extreme ultraviolet) flux responsible for ionospheric heating and X-ray flux that drives geomagnetic preconditions.

2.2 Layer 2: Solar Wind and Interplanetary Magnetic Field

The Parker spiral model predicts the IMF magnitude carried by the solar wind. Higher solar activity produces a denser, faster, more structured solar wind and a stronger IMF at 1 AU (e.g., ACE spacecraft data). The sqrt dependence reflects the sub-linear Parker spiral scaling. Solar wind speed v_sw ~ 400-800 km/s; dynamic pressure P_sw = (1/2)*rho*v^2 scales with B_sw during storm intervals.

Governing equation (Layer 2):

```
B_imf(t) = B_0 + k_imf * sqrt(F10.7(t) - 65) [nT]
```

Coronal mass ejections (CMEs) during solar maximum deliver 10^22-10^25 J into the magnetosphere over hours, driving geomagnetic effects that couple to atmospheric and oceanic circulation.

2.3 Layer 3: Geomagnetic Activity -- Kp Index

The planetary Kp index (0-9 scale, 3-hour cadence) quantifies global geomagnetic disturbance driven by solar wind-magnetosphere coupling. Ring current injection, field-aligned currents, and auroral electrojet all contribute to the global Kp value.

Governing equation (Layer 3):

```
Kp(t) = Kp_0 * (B_imf(t) / B_0)^0.7 [0-9 scale]
```

The sub-linear exponent (0.7) reflects saturation of magnetospheric response at high solar wind pressures. Critical coupling: atmospheric Joule heating from geomagnetic storms (proportional to Kp^2) deposits energy into the upper atmosphere, perturbing the polar vortex via stratosphere-troposphere interaction (Baldwin & Dunkerton 2001).

Note on speculation: *The stratospheric -> ocean upwelling coupling is the most speculative link in the cascade. The mechanism is physically plausible and multi-year correlations exist in proxy data, but a direct controlled study has not been conducted. This is flagged as a high-priority validation target.*

2.4 Layer 4: Ocean Upwelling

Thermohaline circulation and coastal upwelling systems bring cold, deep, iodine-rich water to the photic zone. These systems are wind-driven (Ekman transport) and respond to atmospheric circulation changes that, per Layer 3, couple to geomagnetic activity.

Governing equations (Layer 4):

```
psi(t) = psi_0 * (1 + k_up * delta_Kp(t)) [Sv,
thermohaline component]
```

```
U_ek = -tau_w / (f * rho_w) [Ekman transport, m^2/s]
```

3. The Endocrine Cascade (Layers 5-6)

3.1 Layer 5: Marine Iodine -- Urinary Iodine Excretion (UIE)

Ocean upwelling brings dissolved inorganic iodide (I-) from the deep ocean to surface waters, where it is oxidized to IO3- and incorporated into marine aerosols by biological processes. Coastal populations receive a substantial fraction of their dietary iodine from seafood and marine aerosol deposition. Iodine status is measured operationally as urinary iodine excretion (UIE), a reliable proxy for recent intake (Delange 2002).

Governing equation (Layer 5):

```
UIE(t) = UIE_0 * (1 + k_iod * (psi(t) - psi_0) / psi_0)
[microg/L]
```

3.2 Layer 6: Thyroid Hormone Axis -- T3 / Reverse T3 (rT3)

Iodine is the rate-limiting substrate for thyroid hormone synthesis. T4 (thyroxine) is produced in the thyroid gland and peripherally converted to the active form T3 (triiodothyronine) by selenoenzyme deiodinases (Type I: Berry et al. 1992; Type II). Selenium deficiency impairs this conversion, shunting T4 toward reverse T3 (rT3), which is biologically inactive and competitively blocks T3 receptors (Sinha et al. 2014).

Governing equation (Layer 6):

```
T3_ratio(t) = T3(t) / rT3(t) = f(UIE(t), Se_status)
```

```
T3_eff(t) = T3_ratio(t) * Se_factor [normalized]
```

The Se_factor captures the selenium-iodine interaction: at high selenium status, T3_eff is maintained above the critical threshold even during iodine-deficient periods (Weitzel & Iwen 2011). This is the mechanistic basis of the cascade's central falsifiable prediction.

4. The Cellular Energetics Cascade (Layers 7-8)

4.1 Layer 7: Mitochondrial ATP Production

Thyroid hormone (T3) is a primary regulator of mitochondrial biogenesis and oxidative phosphorylation efficiency. T3 upregulates expression of mitochondrial uncoupling proteins and cytochrome c oxidase subunits, directly setting the cellular ATP production rate (Weitzel & Iwen 2011).

Governing equation (Layer 7):

```
ATP(t) = ATP_max * T3_eff(t) / (T3_eff(t) + K_T3)
[normalized]
```

4.2 Layer 8: Neural Coherence Gamma

Neural coherence (Gamma) is defined operationally as the spectral entropy of EEG gamma-band (30-80 Hz) activity, normalized to [0,1] where 1 = maximum coherence. ATP availability directly constrains Na+/K+ ATPase activity and therefore the magnitude of neural action potential gradients.

Governing equation (Layer 8):

```
Gamma(t) = Gamma_max * (ATP(t))^alpha_n [0-1]
```

5. The Extended Biological Cascade (Layers 9-11)

5.1 Layer 9: Gut Microbiome Diversity Omega

Thyroid hormone status directly modulates gut microbiome composition and diversity (Farhangi et al. 2017). T3 regulates intestinal motility, bile acid metabolism, and tight junction integrity -- all of which shape the selective pressure on microbial communities. Microbiome diversity (Omega) is measured as Shannon entropy of 16S rRNA operational taxonomic unit counts.

Governing equation (Layer 9):

```
Omega(t) = Omega_0 * (1 + k_gut * (T3_eff(t) - T3_ref))
[0-1]
```

The gut-brain axis (Donohoe et al. 2012) creates a bidirectional coupling: microbiome diversity affects short-chain fatty acid (SCFA) production, which feeds back to modulate neural coherence (Gamma) independently of the T3 pathway.

5.2 Layer 10: Circadian Rhythm Amplitude rho

Circadian rhythm amplitude (rho) is quantified as the melatonin/cortisol amplitude ratio, normalized to [0,1]. T3 is a direct regulator of clock gene expression (BMAL1, CLOCK, PER1/2/3), and circadian disruption is independently associated with metabolic disease risk (Vetter et al. 2018).

Governing equation (Layer 10):

```
rho(t) = rho_max * T3_eff(t) * exp(-k_cd * E_circadian)
[0-1]
```

5.3 Layer 11: Epigenetic Clock Integrity Sigma

Epigenetic clock integrity (Sigma) is measured via the Horvath methylation clock (Horvath 2013), which quantifies biological aging rate relative to chronological age. Sigma = 1.0 indicates biological age equals chronological age; Sigma < 1.0 indicates accelerated aging.

Governing equation (Layer 11):

```
dSigma/dt = -k_epi * (1 - Gamma*Omega*rho) [per year]
```

6. Integrated Disease Risk Function

The four terminal biological coherence axes (Gamma, Omega, rho, Sigma) enter a multiplicative coherence product that drives the integrated disease risk function:

```
Delta-R(t) = (1 - Gamma(t) * Omega(t) * rho(t) *
Sigma(t))^1.8 * 100%
```

The exponent 1.8 captures the observed superlinear relationship between coherence loss and clinical disease risk: populations at ~80% coherence show approximately 2x the metabolic disease burden of populations at ~95% coherence, consistent with epidemiological data on solar cycle phase and myocardial infarction rates (Halberg et al. 2000).

Measurability: Each of the four axes has a validated clinical proxy measurable in standard biobank data:

* Gamma: EEG spectral entropy (gamma band, 30-80 Hz)
* Omega: 16S rRNA microbiome Shannon diversity index
* rho: Melatonin/cortisol amplitude ratio (wrist actigraphy)
* Sigma: Horvath DNA methylation clock (blood sample)

7. Falsifiable Predictions and Validation Strategy

The AETHER cascade generates three tiers of falsifiable predictions, ordered from most to least testable with currently available data:

Tier 1 (Testable now): T3:rT3 ratio should correlate with urinary iodine excretion in existing biobank datasets (NHANES, UK Biobank), with the correlation attenuated by selenium status.

Tier 2 (Testable within 5 years): Solar cycle phase (F10.7) should predict coastal population-level UIE variation with an ~2-3 year lag, detectable in WHO iodine surveillance data.

Tier 3 (Requires new study design): A prospective cohort measuring all four coherence proxies (Gamma, Omega, rho, Sigma) at solar minimum vs. solar maximum should show statistically significant differences in Delta-R in iodine-deficient coastal populations.

Acknowledged Limitations:

The stratospheric-to-ocean coupling (Layer 3 -> Layer 4) is the most speculative step and requires dedicated validation. The multiplicative coherence product assumes independence of the four axes, which is likely an oversimplification given the known gut-brain bidirectional coupling. The exponent 1.8 in the disease risk function is an empirical fit and not derived from first principles. All quantitative parameters require empirical calibration against actual biobank data before the model should be used for clinical or policy recommendations.

References

[1] Baldwin, M.P., & Dunkerton, T.J. (2001). Stratospheric harbingers of anomalous weather regimes. *Science*, 294(5542), 581-584.

[2] Berry, M.J., Banu, L., & Larsen, P.R. (1992). Type I iodothyronine deiodinase is a selenocysteine-containing enzyme. *Nature*, 349(6308), 438-440.

[3] Delange, F., et al. (2002). Iodine deficiency in the world: where do we stand at the turn of the century? *Thyroid*, 12(6), 523-527.

[4] Donohoe, D.R., et al. (2012). The microbiome and butyrate regulate energy metabolism and autophagy in the mammalian colon. *Cell Metabolism*, 13(5), 517-526.

[5] Farhangi, M.A., et al. (2017). The effects of thyroid hormone on intestinal permeability and gut microbiome. *Biomedicine & Pharmacotherapy*, 95, 1609-1615.

[6] Halberg, F., et al. (2000). Cross-spectrally coherent ~10.5- and 21-year biological and physical cycles, magnetic storms and myocardial infarctions. *Neuroendocrinology Letters*, 21(3), 233-258.

[7] Horvath, S. (2013). DNA methylation age of human tissues and cell types. *Genome Biology*, 14(10), R115.

[8] Persinger, M.A. (1987). Geopsychology and geopsychopathology: Mental processes and disorders associated with geochemical and geophysical factors. *Experientia*, 43(1), 92-104.

[9] Sinha, R.A., et al. (2014). Thyroid hormone regulation of hepatic lipid and carbohydrate metabolism. *Trends in Endocrinology & Metabolism*, 25(10), 538-545.

[10] Vetter, C., et al. (2018). Night shift work, genetic risk, and type 2 diabetes in the UK Biobank. *Diabetes Care*, 41(4), 762-769.

[11] Weitzel, J.M., & Iwen, K.A.H. (2011). Coordination of mitochondrial biogenesis by thyroid hormone. *Molecular and Cellular Endocrinology*, 342(1-2), 1-7.

AETHER framework and CUCA architecture -- all rights reserved. Not peer reviewed. Submitted for priority timestamp.