

Integrals as anti-derivatives

Sarang S. Sane

Recall : Riemann sums and the definite integral

Recall : Riemann sums and the definite integral

Let f be a function defined on a domain containing the interval $[a, b]$ to \mathbb{R} .

Recall : Riemann sums and the definite integral

Let f be a function defined on a domain containing the interval $[a, b]$ to \mathbb{R} .

Let P consist of (i) a partition $a = x_0 < x_1 < \dots x_n = b$ of $[a, b]$ and (ii) a choice of $x_i^* \in [x_{i-1}, x_i]$.

Recall : Riemann sums and the definite integral

Let f be a function defined on a domain containing the interval $[a, b]$ to \mathbb{R} .

Let P consist of (i) a partition $a = x_0 < x_1 < \dots x_n = b$ of $[a, b]$ and (ii) a choice of $x_i^* \in [x_{i-1}, x_i]$.

Define $\Delta x_i = x_i - x_{i-1}$ and $\|P\| = \max_i \{\Delta x_i\}$.

Recall : Riemann sums and the definite integral

Let f be a function defined on a domain containing the interval $[a, b]$ to \mathbb{R} .

Let P consist of (i) a partition $a = x_0 < x_1 < \dots < x_n = b$ of $[a, b]$ and (ii) a choice of $x_i^* \in [x_{i-1}, x_i]$.

Define $\Delta x_i = x_i - x_{i-1}$ and $\|P\| = \max_i \{\Delta x_i\}$.

The **Riemann sum** of f w.r.t. the above data is defined as

$$S(P) = \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

Recall : Riemann sums and the definite integral

Let f be a function defined on a domain containing the interval $[a, b]$ to \mathbb{R} .

Let P consist of (i) a partition $a = x_0 < x_1 < \dots < x_n = b$ of $[a, b]$ and (ii) a choice of $x_i^* \in [x_{i-1}, x_i]$.

Define $\Delta x_i = x_i - x_{i-1}$ and $\|P\| = \max_i \{\Delta x_i\}$.

The **Riemann sum** of f w.r.t. the above data is defined as

$$S(P) = \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

The **(definite) integral of f from a to b** is defined as

Recall : Riemann sums and the definite integral

Let f be a function defined on a domain containing the interval $[a, b]$ to \mathbb{R} .

Let P consist of (i) a partition $a = x_0 < x_1 < \dots < x_n = b$ of $[a, b]$ and (ii) a choice of $x_i^* \in [x_{i-1}, x_i]$.

Define $\Delta x_i = x_i - x_{i-1}$ and $\|P\| = \max_i \{\Delta x_i\}$.

The **Riemann sum** of f w.r.t. the above data is defined as

$$S(P) = \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

The **(definite) integral of f from a to b** is defined as

$$\int_a^b f(x) dx = \lim_{\|P\| \rightarrow 0} S(P) = \lim_{\|P\| \rightarrow 0} \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

Anti-derivatives aka (indefinite) integrals

Let f be a function defined from a domain D in \mathbb{R} to \mathbb{R} .

Anti-derivatives aka (indefinite) integrals

Let f be a function defined from a domain D in \mathbb{R} to \mathbb{R} .

An **anti-derivative** of the function f is a function F defined on the domain D such that $F'(x) = f(x)$ for all $x \in D$.

Anti-derivatives aka (indefinite) integrals

Let f be a function defined from a domain D in \mathbb{R} to \mathbb{R} .

An **anti-derivative** of the function f is a function F defined on the domain D such that $F'(x) = f(x)$ for all $x \in D$.

Example : An anti-derivative of $f(x) = x^7 + 2x^6 - \pi x^5 + 0.5x^4 - 9$ is $F(x) = \frac{x^8}{8} + 2\frac{x^7}{7} - \pi\frac{x^6}{6} + 0.5\frac{x^5}{5} - 9x$

Anti-derivatives aka (indefinite) integrals

Let f be a function defined from a domain D in \mathbb{R} to \mathbb{R} .

An **anti-derivative** of the function f is a function F defined on the domain D such that $F'(x) = f(x)$ for all $x \in D$.

Example : An anti-derivative of $f(x) = x^7 + 2x^6 - \pi x^5 + 0.5x^4 - 9$ is $F(x) = \frac{x^8}{8} + 2\frac{x^7}{7} - \pi\frac{x^6}{6} + 0.5\frac{x^5}{5} - 9x$

Fact : If F is an anti-derivative of f , then so is $F_1(x) = F(x) + c$ where c is any constant (i.e. a real number).

Anti-derivatives aka (indefinite) integrals

Let f be a function defined from a domain D in \mathbb{R} to \mathbb{R} .

An **anti-derivative** of the function f is a function F defined on the domain D such that $F'(x) = f(x)$ for all $x \in D$.

Example : An anti-derivative of $f(x) = x^7 + 2x^6 - \pi x^5 + 0.5x^4 - 9$ is $F(x) = \frac{x^8}{8} + 2\frac{x^7}{7} - \pi\frac{x^6}{6} + 0.5\frac{x^5}{5} - 9x$

Fact : If F is an anti-derivative of f , then so is $F_1(x) = F(x) + c$ where c is any constant (i.e. a real number). In fact, every anti-derivative has this form.

Anti-derivatives aka (indefinite) integrals

Let f be a function defined from a domain D in \mathbb{R} to \mathbb{R} .

An **anti-derivative** of the function f is a function F defined on the domain D such that $F'(x) = f(x)$ for all $x \in D$.

Example : An anti-derivative of $f(x) = x^7 + 2x^6 - \pi x^5 + 0.5x^4 - 9$ is $F(x) = \frac{x^8}{8} + 2\frac{x^7}{7} - \pi\frac{x^6}{6} + 0.5\frac{x^5}{5} - 9x + C$

Fact : If F is an anti-derivative of f , then so is $F_1(x) = F(x) + c$ where c is any constant (i.e. a real number). In fact, every anti-derivative has this form.

Because of this, We will use **the** instead of **an** for the anti-derivative, since any two anti-derivatives only differ by a constant.

Anti-derivatives and integrals

The anti-derivative for a function f is more often called the (indefinite) integral of f and denoted by $\int f(x)dx$.

Anti-derivatives and integrals

The anti-derivative for a function f is more often called the (indefinite) integral of f and denoted by $\int f(x)dx$.

This is explained by :

The fundamental theorem of calculus

Anti-derivatives and integrals

The anti-derivative for a function f is more often called the **(indefinite) integral of f** and denoted by $\int f(x)dx$.

This is explained by :

The fundamental theorem of calculus

Suppose f is continuous on the domain D which includes the interval $[a, b]$. Then an anti-derivative for f on (a, b) is given by

$$F(x) = \int_a^x f(t)dt.$$

Anti-derivatives and integrals

The anti-derivative for a function f is more often called the **(indefinite) integral of f** and denoted by $\int f(x)dx$.

This is explained by :

The fundamental theorem of calculus

Suppose f is continuous on the domain D which includes the interval $[a, b]$. Then an anti-derivative for f on (a, b) is given by

$$F(x) = \int_a^x f(t)dt.$$

Conversely, if f is continuous on the domain D which includes the interval $[a, b]$ and F is the (indefinite) integral of f , then the (definite) integral from a to b of f can be computed by

$$\int_a^b f(x)dx = F(b) - F(a).$$

Tables of derivatives and integrals

Function	Derivative	Integral
1	0	x
x^r	rx^{r-1}	$\frac{x^{r+1}}{r+1} \quad r \neq -1$
		$\ln x \quad r = -1$
$\sin(x)$	$\cos(x)$	$-\cos(x)$
$\cos(x)$	$-\sin(x)$	$\sin(x)$
$\tan(x)$	$\sec^2(x)$	$\ln \sec(x) $
$\sec(x)$	$\sec(x)\tan(x)$	$\ln \sec(x) + \tan(x) $
$\cot(x)$	$-\csc^2(x)$	$\ln \sin(x) $
$\operatorname{cosec}(x)$	$-\csc^2(x)$	$\ln \operatorname{cosec}(x) - \cot(x) $

can be
obtained

Tables (contd.)

Function	Derivative	Integral
$e^{\lambda x}$	$\lambda e^{\lambda x}$	$\frac{e^{\lambda x}}{\lambda}$
a^x	$a^x \ln(a)$	$\frac{a^x}{\ln(a)}$
$\ln(x)$	$\frac{1}{x}$	$x \ln(x) - x$
$\frac{1}{\sqrt{a^2 - x^2}}$		$\sin^{-1}\left(\frac{x}{a}\right)$
$\frac{1}{\sqrt{a^2 + x^2}}$		$\tan^{-1}\left(\frac{x}{a}\right)$
$\frac{1}{a^2 - x^2}$		$\frac{1}{2a} \ln \left \frac{x+a}{x-a} \right $

Examples

$$\int_1^2 x dx = \left[\frac{x^2}{2} \right]_1^2 = \frac{2^2}{2} - \frac{1^2}{2} = \frac{4}{2} - \frac{1}{2} = \frac{3}{2}.$$

$$\begin{aligned} f(x) &= x \\ F(x) &= x^2/2 \\ F(2) - F(1) \end{aligned}$$

$$\int_0^\pi \sin(x) dx = \left[-\cos(x) \right]_0^\pi = (-\cos(\pi)) - (-\cos(0)) = (-(-1)) - (-1) = 1 + 1 = 2.$$



$$\begin{aligned} \int_0^\infty e^{-2x} dx &= \lim_{b \rightarrow \infty} \int_0^b e^{-2x} dx \\ &= \lim_{b \rightarrow \infty} \left[\frac{e^{-2x}}{-2} \right]_0^b = \lim_{b \rightarrow \infty} \frac{-1}{2} (e^{-2b} - e^0) \\ &= \lim_{b \rightarrow \infty} \frac{1}{2} - \frac{e^{-2b}}{2} = \frac{1}{2}. \end{aligned}$$

$$\int_a^\infty f(x) dx = \lim_{b \rightarrow \infty} \int_a^b f(x) dx$$

Thank you