

Statistics for Data Science -1

Lecture 9.4 Variance of a Random Variable

Usha Mohan

Indian Institute of Technology Madras

Learning objectives

Learning objectives

1. Define what is a random variable.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.
3. Probability mass function, graph, and examples.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.
3. Probability mass function, graph, and examples.
4. Cumulative distribution function, graphs, and examples.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.
3. Probability mass function, graph, and examples.
4. Cumulative distribution function, graphs, and examples.
5. Expectation and variance of a random variable.

Introduction

- ▶ The expected value of a random variable gives the weighted average of the possible values of the random variable, it does not tell us anything about the variation, or spread, of these values.

Introduction

- ▶ The expected value of a random variable gives the weighted average of the possible values of the random variable, it does not tell us anything about the variation, or spread, of these values.
- ▶ For instance, consider random variables X , Y , and Z , whose values and probabilities are as follows:

Introduction

- ▶ The expected value of a random variable gives the weighted average of the possible values of the random variable, it does not tell us anything about the variation, or spread, of these values.
- ▶ For instance, consider random variables X , Y , and Z , whose values and probabilities are as follows:
 - ▶ $X = 0$ with probability 1

Introduction

- ▶ The expected value of a random variable gives the weighted average of the possible values of the random variable, it does not tell us anything about the variation, or spread, of these values.
- ▶ For instance, consider random variables X , Y , and Z , whose values and probabilities are as follows:
 - ▶ $X = 0$ with probability 1
 - ▶ $Y = \begin{cases} -2 & \text{with probability } \frac{1}{2} \\ 2 & \text{with probability } \frac{1}{2} \end{cases}$

Introduction

- ▶ The expected value of a random variable gives the weighted average of the possible values of the random variable, it does not tell us anything about the variation, or spread, of these values.
- ▶ For instance, consider random variables X , Y , and Z , whose values and probabilities are as follows:
 - ▶ $X = 0$ with probability 1
 - ▶ $Y = \begin{cases} -2 & \text{with probability } \frac{1}{2} \\ 2 & \text{with probability } \frac{1}{2} \end{cases}$
 - ▶ $Z = \begin{cases} -20 & \text{with probability } \frac{1}{2} \\ 20 & \text{with probability } \frac{1}{2} \end{cases}$

Introduction

- ▶ The expected value of a random variable gives the weighted average of the possible values of the random variable, it does not tell us anything about the variation, or spread, of these values.
- ▶ For instance, consider random variables X , Y , and Z , whose values and probabilities are as follows:
 - ▶ $X = 0$ with probability 1
 - ▶ $Y = \begin{cases} -2 & \text{with probability } \frac{1}{2} \\ 2 & \text{with probability } \frac{1}{2} \end{cases}$
 - ▶ $Z = \begin{cases} -20 & \text{with probability } \frac{1}{2} \\ 20 & \text{with probability } \frac{1}{2} \end{cases}$
- ▶ $E(X) = E(Y) = E(Z) = 0$. However, we notice spread of Z is greater than spread of Y which is greater than spread of X

Introduction

- ▶ The expected value of a random variable gives the weighted average of the possible values of the random variable, it does not tell us anything about the variation, or spread, of these values.
- ▶ For instance, consider random variables X , Y , and Z , whose values and probabilities are as follows:
 - ▶ $X = 0$ with probability 1
 - ▶ $Y = \begin{cases} -2 & \text{with probability } \frac{1}{2} \\ 2 & \text{with probability } \frac{1}{2} \end{cases}$
 - ▶ $Z = \begin{cases} -20 & \text{with probability } \frac{1}{2} \\ 20 & \text{with probability } \frac{1}{2} \end{cases}$
- ▶ $E(X) = E(Y) = E(Z) = 0$. However, we notice spread of Z is greater than spread of Y which is greater than spread of X
- ▶ Need for a measure of spread.

Variance of a random variable

Variance of a random variable

- ▶ Let's denote expected value of a random variable X by the greek alphabet μ .

Variance of a random variable

- Let's denote expected value of a random variable X by the greek alphabet μ .

Definition

Let X be a random variable with expected value μ , then the variance of X , denoted by $\text{Var}(X)$ or $V(X)$, is defined by

$$\text{Var}(X) = E(X - \mu)^2$$

Variance of a random variable

- ▶ Let's denote expected value of a random variable X by the greek alphabet μ .

Definition

Let X be a random variable with expected value μ , then the variance of X , denoted by $\text{Var}(X)$ or $V(X)$, is defined by

$$\text{Var}(X) = E (X - \mu)^2$$

- ▶ In other words, the Variance of a random variable X measures the square of the difference of the random variable from its mean, μ , on the average.

Computational formula for $Var(X)$

Computational formula for $Var(X)$

► $Var(X) = E(X - \mu)^2$

Computational formula for $Var(X)$

- ▶ $Var(X) = E(X - \mu)^2$
- ▶ $(X - \mu)^2 = X^2 - 2X\mu + \mu^2$

Computational formula for $Var(X)$

- ▶ $Var(X) = E(X - \mu)^2$
- ▶ $(X - \mu)^2 = X^2 - 2X\mu + \mu^2$
- ▶ Using properties of expectation we know
 $E(X^2 - 2X\mu + \mu^2) = E(X^2) - 2\mu E(X) + \mu^2$ which is same as
 $E(X^2) - \mu^2$

Computational formula for $Var(X)$

- ▶ $Var(X) = E(X - \mu)^2$
- ▶ $(X - \mu)^2 = X^2 - 2X\mu + \mu^2$
- ▶ Using properties of expectation we know
 $E(X^2 - 2X\mu + \mu^2) = E(X^2) - 2\mu E(X) + \mu^2$ which is same as
 $E(X^2) - \mu^2$
- ▶ Let's compute the variance of the random variables discussed earlier.

Rolling a dice once

- ▶ Random experiment: Roll a dice once.
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Random variable X is the outcome of the roll.

Rolling a dice once

- ▶ Random experiment: Roll a dice once.
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Random variable X is the outcome of the roll.
- ▶ The probability distribution is given by

[illegible]

Rolling a dice once

- ▶ Random experiment: Roll a dice once.
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Random variable X is the outcome of the roll.
- ▶ The probability distribution is given by

[illegible]

- $E(X) = 1\frac{1}{6} + 2\frac{1}{6} + 3\frac{1}{6} + 4\frac{1}{6} + 5\frac{1}{6} + 6\frac{1}{6} = 3.5.$

Rolling a dice once

- ▶ Random experiment: Roll a dice once.
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Random variable X is the outcome of the roll.
- ▶ The probability distribution is given by

[illegible]

- ▶ $E(X) = 1\frac{1}{6} + 2\frac{1}{6} + 3\frac{1}{6} + 4\frac{1}{6} + 5\frac{1}{6} + 6\frac{1}{6} = 3.5$.
- ▶ $E(X^2) = 1\frac{1}{6} + 4\frac{1}{6} + 9\frac{1}{6} + 16\frac{1}{6} + 25\frac{1}{6} + 36\frac{1}{6} = 15.167$

Rolling a dice once

- ▶ Random experiment: Roll a dice once.
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Random variable X is the outcome of the roll.
- ▶ The probability distribution is given by

[illegible]

- $E(X) = 1\frac{1}{6} + 2\frac{1}{6} + 3\frac{1}{6} + 4\frac{1}{6} + 5\frac{1}{6} + 6\frac{1}{6} = 3.5.$
- $E(X^2) = 1\frac{1}{6} + 4\frac{1}{6} + 9\frac{1}{6} + 16\frac{1}{6} + 25\frac{1}{6} + 36\frac{1}{6} = 15.167$
- $Var(X) = 15.167 - 3.5^2 = 2.917$

Rolling a dice twice

- ▶ X is a random variable which is defined as sum of outcomes

Rolling a dice twice

- ▶ X is a random variable which is defined as sum of outcomes
- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
X^2	4	9	16	25	36	49	64	81	100	121	144
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

Rolling a dice twice

- ▶ X is a random variable which is defined as sum of outcomes
- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
X^2	4	9	16	25	36	49	64	81	100	121	144
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ $E(X) = 2 \times \frac{1}{36} + 3 \times \frac{2}{36} + \dots + 11 \times \frac{2}{36} + 12 \times \frac{1}{36} = 7$

Rolling a dice twice

- ▶ X is a random variable which is defined as sum of outcomes
- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
X^2	4	9	16	25	36	49	64	81	100	121	144
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$


- ▶ $E(X) = 2 \times \frac{1}{36} + 3 \times \frac{2}{36} + \dots + 11 \times \frac{2}{36} + 12 \times \frac{1}{36} = 7$
- ▶ $E(X^2) = 4 \times \frac{1}{36} + 9 \times \frac{2}{36} + \dots + 121 \times \frac{2}{36} + 144 \times \frac{1}{36} = 54.833$

Rolling a dice twice

- ▶ X is a random variable which is defined as sum of outcomes
- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
X^2	4	9	16	25	36	49	64	81	100	121	144
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ $E(X) = 2 \times \frac{1}{36} + 3 \times \frac{2}{36} + \dots + 11 \times \frac{2}{36} + 12 \times \frac{1}{36} = 7$
- ▶ $E(X^2) = 4 \times \frac{1}{36} + 9 \times \frac{2}{36} + \dots + 121 \times \frac{2}{36} + 144 \times \frac{1}{36} = 54.833$
- ▶ $Var(X) = 54.833 - 49 = 5.833$



Tossing a coin thrice

► $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$

Tossing a coin thrice

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

Tossing a coin thrice

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
X^2	0	1	4	9
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Tossing a coin thrice

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
X^2	0	1	4	9
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ $E(X) = \sum_{i=0}^3 x_i p(x_i) =$

Tossing a coin thrice

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
X^2	0	1	4	9
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶
$$E(X) = \sum_{i=0}^3 x_i p(x_i) =$$
$$\frac{(0 \times 1) + (1 \times 3) + (2 \times 3) + (3 \times 1)}{8} = \frac{3}{2}$$

Tossing a coin thrice

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
X^2	0	1	4	9
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ $E(X) = \sum_{i=0}^3 x_i p(x_i) =$

$$\frac{(0 \times 1) + (1 \times 3) + (2 \times 3) + (3 \times 1)}{8} = \frac{3}{2}$$

- ▶ $E(X^2) = \sum_{i=0}^3 x_i^2 p(x_i) =$

Tossing a coin thrice

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
X^2	0	1	4	9
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ $E(X) = \sum_{i=0}^3 x_i p(x_i) =$

$$\frac{(0 \times 1) + (1 \times 3) + (2 \times 3) + (3 \times 1)}{8} = \frac{3}{2}$$

- ▶ $E(X^2) = \sum_{i=0}^3 x_i^2 p(x_i) =$

$$\frac{(0 \times 1) + (1 \times 3) + (4 \times 3) + (9 \times 1)}{8} = \frac{24}{8} = 3$$

Tossing a coin thrice

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
X^2	0	1	4	9
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$


- ▶ $E(X) = \sum_{i=0}^3 x_i p(x_i) =$

$$\frac{(0 \times 1) + (1 \times 3) + (2 \times 3) + (3 \times 1)}{8} = \frac{3}{2}$$

- ▶ $E(X^2) = \sum_{i=0}^3 x_i^2 p(x_i) =$

$$\frac{(0 \times 1) + (1 \times 3) + (4 \times 3) + (9 \times 1)}{8} = \frac{24}{8} = 3$$

- ▶ $Var(X) = 3 - 2.25 = 0.75$



Bernoulli random variable

- ▶ A random variable that takes on either the value 1 or 0 is called a Bernoulli random variable.
- ▶ Let X be a Bernoulli random variable that takes on the value 1 with probability p .
- ▶ The probability distribution of the random variable is

X	0	1

Bernoulli random variable

- ▶ A random variable that takes on either the value 1 or 0 is called a Bernoulli random variable.
- ▶ Let X be a Bernoulli random variable that takes on the value 1 with probability p .
- ▶ The probability distribution of the random variable is

X	0	1
X^2	0	1

Bernoulli random variable

- ▶ A random variable that takes on either the value 1 or 0 is called a Bernoulli random variable.
- ▶ Let X be a Bernoulli random variable that takes on the value 1 with probability p .
- ▶ The probability distribution of the random variable is

X	0	1
X^2	0	1
$P(X = x_i)$	$1 - p$	p

Bernoulli random variable

- ▶ A random variable that takes on either the value 1 or 0 is called a Bernoulli random variable.
- ▶ Let X be a Bernoulli random variable that takes on the value 1 with probability p .
- ▶ The probability distribution of the random variable is

X	0	1
X^2	0	1
$P(X = x_i)$	$1 - p$	p

- ▶ Expected value of a Bernoulli random variable:
$$E(X) = 0 \times (1 - p) + 1 \times p = p$$

Bernoulli random variable

- ▶ A random variable that takes on either the value 1 or 0 is called a Bernoulli random variable.
- ▶ Let X be a Bernoulli random variable that takes on the value 1 with probability p .
- ▶ The probability distribution of the random variable is

X	0	1
X^2	0	1
$P(X = x_i)$	$1 - p$	p

- ▶ Expected value of a Bernoulli random variable:
$$E(X) = 0 \times (1 - p) + 1 \times p = p$$
- ▶ Variance of a Bernoulli random variable:
$$Var(X) = p - p^2 = p(1 - p)$$

Discrete uniform random variable

- ▶ Let X be a random variable that is equally likely to takes any of the values $1, 2, \dots, n$
- ▶ Probability mass function

X	1	2	\dots	n

Discrete uniform random variable

- ▶ Let X be a random variable that is equally likely to takes any of the values $1, 2, \dots, n$
- ▶ Probability mass function

X	1	2	\dots	n
X^2	1	4	\dots	n^2
$P(X = x_i)$	$\frac{1}{n}$	$\frac{1}{n}$	\dots	$\frac{1}{n}$

Discrete uniform random variable

- ▶ Let X be a random variable that is equally likely to takes any of the values $1, 2, \dots, n$
- ▶ Probability mass function

X	1	2	\dots	n
X^2	1	4	\dots	n^2
$P(X = x_i)$	$\frac{1}{n}$	$\frac{1}{n}$	\dots	$\frac{1}{n}$

- $E(X) = \frac{(n+1)}{2}$

Discrete uniform random variable

- ▶ Let X be a random variable that is equally likely to takes any of the values $1, 2, \dots, n$
- ▶ Probability mass function

X	1	2	\dots	n
X^2	1	4	\dots	n^2
$P(X = x_i)$	$\frac{1}{n}$	$\frac{1}{n}$	\dots	$\frac{1}{n}$

- ▶ $E(X) = \frac{(n+1)}{2}$
- ▶ $E(X^2) = \frac{(n+1)(2n+1)}{6}$

Discrete uniform random variable

- ▶ Let X be a random variable that is equally likely to takes any of the values $1, 2, \dots, n$
- ▶ Probability mass function

X	1	2	\dots	n
X^2	1	4	\dots	n^2
$P(X = x_i)$	$\frac{1}{n}$	$\frac{1}{n}$	\dots	$\frac{1}{n}$

- ▶ $E(X) = \frac{(n+1)}{2}$
- ▶ $E(X^2) = \frac{(n+1)(2n+1)}{6}$
- ▶ $Var(X) = \frac{n^2-1}{12}$

Section summary

- ▶ Definition of variance
- ▶ Computational formula of variance of a random variable.