

Statistics for Data Science -1

Lecture 8.4: Discrete Random Variable: probability mass function

Usha Mohan

Indian Institute of Technology Madras

Learning objectives

Learning objectives

1. Define what is a random variable.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.
3. Probability mass function, graph, and examples.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.
3. Probability mass function, graph, and examples.
4. Cumulative distribution function, graphs, and examples.

Learning objectives

1. Define what is a random variable.
2. Types of random variables: discrete and continuous.
3. Probability mass function, graph, and examples.
4. Cumulative distribution function, graphs, and examples.
5. Expectation and variance of a random variable.

Probability mass function, graph, and examples

Probability mass function

Probability mass function (p.m.f)

Probability mass function (p.m.f)

- ▶ A random variable that can take on at most a countable number of possible values is said to be a discrete random variable.

Probability mass function (p.m.f)

- ▶ A random variable that can take on at most a countable number of possible values is said to be a discrete random variable.
- ▶ Let X be a discrete random variable, and suppose that it has n possible values, which we will label x_1, x_2, \dots, x_n .

Probability mass function (p.m.f)

- ▶ A random variable that can take on at most a countable number of possible values is said to be a discrete random variable.
- ▶ Let X be a discrete random variable, and suppose that it has n possible values, which we will label x_1, x_2, \dots, x_n .
- ▶ For a discrete random variable X , we define the probability mass function $p(x)$ of X by

$$p(x_i) = P(X = x_i)$$

Probability mass function (p.m.f)

- ▶ A random variable that can take on at most a countable number of possible values is said to be a discrete random variable.
- ▶ Let X be a discrete random variable, and suppose that it has n possible values, which we will label x_1, x_2, \dots, x_n .
- ▶ For a discrete random variable X , we define the probability mass function $p(x)$ of X by

$$p(x_i) = P(X = x_i)$$

- ▶ Represent it in tabular form

X	x_1	x_2	x_3	\dots	\dots	x_n
$P(X = x_i)$	$p(x_1)$	$p(x_2)$	$p(x_3)$	\dots	\dots	$p(x_n)$

Properties of p.m.f

Properties of p.m.f

- ▶ The probability mass function $p(x)$ is positive for at most a countable number of values of x . That is, if X must assume one of the values x_1, x_2, \dots , then

Properties of p.m.f

- ▶ The probability mass function $p(x)$ is positive for at most a countable number of values of x . That is, if X must assume one of the values x_1, x_2, \dots , then
 1. $p(x_i) \geq 0, i = 1, 2, \dots$

Properties of p.m.f

- ▶ The probability mass function $p(x)$ is positive for at most a countable number of values of x . That is, if X must assume one of the values x_1, x_2, \dots , then
 1. $p(x_i) \geq 0, i = 1, 2, \dots$
 2. $p(x) = 0$ for all other values of x

Properties of p.m.f

- ▶ The probability mass function $p(x)$ is positive for at most a countable number of values of x . That is, if X must assume one of the values x_1, x_2, \dots , then
 1. $p(x_i) \geq 0, i = 1, 2, \dots$
 2. $p(x) = 0$ for all other values of x
- ▶ Represent it in tabular form

X	x_1	x_2	x_3			
$P(X = x_i)$	$p(x_1)$	$p(x_2)$	$p(x_3)$			

Properties of p.m.f

- ▶ The probability mass function $p(x)$ is positive for at most a countable number of values of x . That is, if X must assume one of the values x_1, x_2, \dots , then
 1. $p(x_i) \geq 0, i = 1, 2, \dots$
 2. $p(x) = 0$ for all other values of x
- ▶ Represent it in tabular form

X	x_1	x_2	x_3			
$P(X = x_i)$	$p(x_1)$	$p(x_2)$	$p(x_3)$			

- ▶ Since X must take one of the values x_i , we have

$$\sum_{i=1}^{\infty} p(x_i) = 1$$

Example

Example

- ▶ Suppose X is a random variable that takes three values, 0, 1, and 2 with probabilities

Example

- ▶ Suppose X is a random variable that takes three values, 0, 1, and 2 with probabilities
 - ▶ $p(0) = P(X = 0) = \frac{1}{4}$
 - ▶ $p(1) = P(X = 1) = \frac{1}{2}$
 - ▶ $p(2) = P(X = 2) = \frac{1}{4}$

Example

- ▶ Suppose X is a random variable that takes three values, 0, 1, and 2 with probabilities
 - ▶ $p(0) = P(X = 0) = \frac{1}{4}$
 - ▶ $p(1) = P(X = 1) = \frac{1}{2}$
 - ▶ $p(2) = P(X = 2) = \frac{1}{4}$
- ▶ Tabular form

X	0	1	2
$P(X = x_i)$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

Example

- ▶ Suppose X is a random variable that takes three values, 0, 1, and 2 with probabilities
 - ▶ $p(0) = P(X = 0) = \frac{1}{4}$
 - ▶ $p(1) = P(X = 1) = \frac{1}{2}$
 - ▶ $p(2) = P(X = 2) = \frac{1}{4}$
- ▶ Tabular form

X	0	1	2
$P(X = x_i)$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

- ▶ Verify that $\sum_{i=1}^3 p(x_i) = \frac{1}{4} + \frac{1}{2} + \frac{1}{4} = 1$

Example

Let X be a random variable that takes values 1,2,3,4,5. Which of the following are probability mass functions?

1.

X	1	2	3	4	5
$P(X = x_i)$	0.4	0.1	0.2	0.1	0.3

Example

Let X be a random variable that takes values 1,2,3,4,5. Which of the following are probability mass functions?

1.

X	1	2	3	4	5
$P(X = x_i)$	0.4	0.1	0.2	0.1	0.3

NO

Example

Let X be a random variable that takes values 1,2,3,4,5. Which of the following are probability mass functions?

1.

X	1	2	3	4	5
$P(X = x_i)$	0.4	0.1	0.2	0.1	0.3

NO

2.

X	1	2	3	4	5
$P(X = x_i)$	0.2	0.3	0.4	-0.1	0.2

Example

Let X be a random variable that takes values 1,2,3,4,5. Which of the following are probability mass functions?

1.

X	1	2	3	4	5
$P(X = x_i)$	0.4	0.1	0.2	0.1	0.3

NO

2.

X	1	2	3	4	5
$P(X = x_i)$	0.2	0.3	0.4	-0.1	0.2

NO

Example

Let X be a random variable that takes values 1,2,3,4,5. Which of the following are probability mass functions?

1.

X	1	2	3	4	5
$P(X = x_i)$	0.4	0.1	0.2	0.1	0.3

NO

2.

X	1	2	3	4	5
$P(X = x_i)$	0.2	0.3	0.4	-0.1	0.2

NO

3.

X	1	2	3	4	5
$P(X = x_i)$	0.3	0.1	0.2	0.4	0.0

Example

Let X be a random variable that takes values 1,2,3,4,5. Which of the following are probability mass functions?

1.

X	1	2	3	4	5
$P(X = x_i)$	0.4	0.1	0.2	0.1	0.3

NO

2.

X	1	2	3	4	5
$P(X = x_i)$	0.2	0.3	0.4	-0.1	0.2

NO

3.

X	1	2	3	4	5
$P(X = x_i)$	0.3	0.1	0.2	0.4	0.0

YES

Example

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities
 - ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities
 - ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ
- ▶ What is the value of c ?

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities
 - ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ
- ▶ What is the value of c ?
 - ▶ $\sum_{i=0}^{\infty} p(x_i) = 1$

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities
 - ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ
- ▶ What is the value of c ?
 - ▶ $\sum_{i=0}^{\infty} p(x_i) = 1$
 - ▶ $\sum_{i=0}^{\infty} c \frac{\lambda^i}{i!} = 1$

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities
 - ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ
- ▶ What is the value of c ?
 - ▶ $\sum_{i=0}^{\infty} p(x_i) = 1$
 - ▶ $\sum_{i=0}^{\infty} c \frac{\lambda^i}{i!} = 1$
 - ▶ $c \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = 1$

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities
 - ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ
- ▶ What is the value of c ?
 - ▶ $\sum_{i=0}^{\infty} p(x_i) = 1$
 - ▶ $\sum_{i=0}^{\infty} c \frac{\lambda^i}{i!} = 1$
 - ▶ $c \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = 1$
 - ▶ Recall, $e^x = \sum_{i=0}^{\infty} \frac{x^i}{i!}$, hence

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities
 - ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ
- ▶ What is the value of c ?
 - ▶ $\sum_{i=0}^{\infty} p(x_i) = 1$
 - ▶ $\sum_{i=0}^{\infty} c \frac{\lambda^i}{i!} = 1$
 - ▶ $c \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = 1$
 - ▶ Recall, $e^x = \sum_{i=0}^{\infty} \frac{x^i}{i!}$, hence $c \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = ce^{\lambda}$

Example

- ▶ Suppose X is a random variable that takes values, 0, 1, 2, ... with probabilities

- ▶ $p(i) = c \frac{\lambda^i}{i!}$, for some positive λ

- ▶ What is the value of c ?

- ▶ $\sum_{i=0}^{\infty} p(x_i) = 1$

- ▶ $\sum_{i=0}^{\infty} c \frac{\lambda^i}{i!} = 1$

- ▶ $c \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = 1$

- ▶ Recall, $e^x = \sum_{i=0}^{\infty} \frac{x^i}{i!}$, hence $c \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = ce^{\lambda}$

- ▶ Hence, $c \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = ce^{\lambda} = 1$ which gives $c = e^{-\lambda}$

Example: Rolling a dice twice

$$\blacktriangleright S = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$$

Example: Rolling a dice twice

- ▶ $S = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes

Example: Rolling a dice twice

- ▶ $S = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes
- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

Example: Rolling a dice twice

- ▶ $S = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes
- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^{11} p(x_i) =$

Example: Rolling a dice twice

- ▶ $S = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes
- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^{11} p(x_i) = \frac{36}{36} = 1$

Example: Rolling a dice twice

- ▶ $S = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes

- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^{11} p(x_i) = \frac{36}{36} = 1$

- ▶ Y is the random variable which takes the lesser of the values of the outcomes

Example: Rolling a dice twice

- ▶ $s = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes

- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^{11} p(x_i) = \frac{36}{36} = 1$

- ▶ Y is the random variable which takes the lesser of the values of the outcomes

- ▶ Probability mass function

Y	1	2	3	4	5	6
$P(Y = y_i)$	$\frac{11}{36}$	$\frac{9}{36}$	$\frac{7}{36}$	$\frac{5}{36}$	$\frac{3}{36}$	$\frac{1}{36}$

Example: Rolling a dice twice

- ▶ $s = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes

- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^{11} p(x_i) = \frac{36}{36} = 1$

- ▶ Y is the random variable which takes the lesser of the values of the outcomes

- ▶ Probability mass function

Y	1	2	3	4	5	6
$P(Y = y_i)$	$\frac{11}{36}$	$\frac{9}{36}$	$\frac{7}{36}$	$\frac{5}{36}$	$\frac{3}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^6 p(y_i) =$

Example: Rolling a dice twice

- ▶ $s = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$
- ▶ X is a random variable which is defined as sum of outcomes

- ▶ Probability mass function

X	2	3	4	5	6	7	8	9	10	11	12
$P(X = x_i)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^{11} p(x_i) = \frac{36}{36} = 1$

- ▶ Y is the random variable which takes the lesser of the values of the outcomes

- ▶ Probability mass function

Y	1	2	3	4	5	6
$P(Y = y_i)$	$\frac{11}{36}$	$\frac{9}{36}$	$\frac{7}{36}$	$\frac{5}{36}$	$\frac{3}{36}$	$\frac{1}{36}$

- ▶ Verify: $\sum_{i=1}^6 p(y_i) = \frac{36}{36} = 1$

Example: Tossing a coin three times

$$\blacktriangleright S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$$

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses
- ▶ Probability mass function

X	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(x_i) =$

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(x_i) = \frac{8}{8} = 1$

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(x_i) = \frac{8}{8} = 1$

- ▶ Y is the random variable which counts the toss in which heads appears first

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(x_i) = \frac{8}{8} = 1$

- ▶ Y is the random variable which counts the toss in which heads appears first

- ▶ Probability mass function

Y	1	2	3	NIL
$P(Y = y_i)$	$\frac{4}{8}$	$\frac{2}{8}$	$\frac{1}{8}$	$\frac{1}{8}$

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(x_i) = \frac{8}{8} = 1$

- ▶ Y is the random variable which counts the toss in which heads appears first

- ▶ Probability mass function

Y	1	2	3	NIL
$P(Y = y_i)$	$\frac{4}{8}$	$\frac{2}{8}$	$\frac{1}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(y_i) =$

Example: Tossing a coin three times

- ▶ $S = \{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
- ▶ X is the random variable which counts the number of heads in the tosses

- ▶ Probability mass function

X	0	1	2	3
$P(X = x_i)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(x_i) = \frac{8}{8} = 1$

- ▶ Y is the random variable which counts the toss in which heads appears first

- ▶ Probability mass function

Y	1	2	3	NIL
$P(Y = y_i)$	$\frac{4}{8}$	$\frac{2}{8}$	$\frac{1}{8}$	$\frac{1}{8}$

- ▶ Verify: $\sum_{i=1}^4 p(y_i) = \frac{8}{8} = 1$

Section summary

- ▶ Probability mass function.
- ▶ Properties of probability mass function.