

# Statistics for Data Science -1

## Lecture 7.3: Conditional Probability: Multiplication rule

Usha Mohan

Indian Institute of Technology Madras

## Learning objectives

1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.

## Learning objectives

1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
2. Distinguish between independent and dependent events.

## Learning objectives

1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
2. Distinguish between independent and dependent events.
3. Solve applications of probability.

## Multiplication rule

# Multiplication rule

## Multiplication rule

- ▶ Conditional probability formula:

$$P(E|F) = \frac{P(E \cap F)}{P(F)}$$

Multiply both sides of the formula with  $P(F)$ , we get

## Multiplication rule

- ▶ Conditional probability formula:

$$P(E|F) = \frac{P(E \cap F)}{P(F)}$$

Multiply both sides of the formula with  $P(F)$ , we get



$$P(E \cap F) = P(F)P(E|F)$$



## Multiplication rule

- ▶ Conditional probability formula:

$$P(E|F) = \frac{P(E \cap F)}{P(F)}$$

Multiply both sides of the formula with  $P(F)$ , we get



$$P(E \cap F) = P(F)P(E|F)$$

- ▶ This rule states that the probability that both  $E$  and  $F$  occur is equal to the probability that  $F$  occurs multiplied by the conditional probability of  $E$  given that  $F$  occurs.

## Multiplication rule

- ▶ Conditional probability formula:

$$P(E|F) = \frac{P(E \cap F)}{P(F)}$$

Multiply both sides of the formula with  $P(F)$ , we get



$$P(E \cap F) = P(F)P(E|F)$$

- ▶ This rule states that the probability that both  $E$  and  $F$  occur is equal to the probability that  $F$  occurs multiplied by the conditional probability of  $E$  given that  $F$  occurs.
- ▶ It is often quite useful for computing the probability of an intersection.

## Example: application of multiplication rule

## Example: application of multiplication rule

- ▶ In an introductory statistics class of forty students, the number of males is equal to 23 and number of females is equal to 17. Two students are selected at random from the class. The first student selected is not returned to the class for possible reselection; that is, the sampling is without replacement. Find the probability that the first student selected is female and the second is male.

## Example: application of multiplication rule

- ▶ In an introductory statistics class of forty students, the number of males is equal to 23 and number of females is equal to 17. Two students are selected at random from the class. The first student selected is not returned to the class for possible reselection; that is, the sampling is without replacement. Find the probability that the first student selected is female and the second is male.
- ▶ Experiment: Selecting two students from forty students.

## Example: application of multiplication rule

- ▶ In an introductory statistics class of forty students, the number of males is equal to 23 and number of females is equal to 17. Two students are selected at random from the class. The first student selected is not returned to the class for possible reselection; that is, the sampling is without replacement. Find the probability that the first student selected is female and the second is male.
- ▶ Experiment: Selecting two students from forty students.
- ▶ Sample space:  $S = \{M_1M_2, M_1F_2, F_1M_2, F_1F_2\}$ ; where  $M_1M_2$  represents the outcome the first student is male and the second student is male. Other outcomes can be interpreted similarly.

## Example: application of multiplication rule-cont.

## Example: application of multiplication rule-cont.

- ▶ Event: First student female and second is male.



## Example: application of multiplication rule-cont.

- ▶ Event: First student female and second is male.
- ▶  $P(F_1 M_2) =$   
 $P(\text{First student is female} \cap \text{Second student is male})$

## Example: application of multiplication rule-cont.

- ▶ Event: First student female and second is male.
- ▶  $P(F_1 M_2) =$   
 $P(\text{First student is female} \cap \text{Second student is male})$
- ▶  $P(\text{First student is female}) = \frac{17}{40}$ .

## Example: application of multiplication rule-cont.

- ▶ Event: First student female and second is male.
- ▶  $P(F_1 M_2) =$   
 $P(\text{First student is female} \cap \text{Second student is male})$
- ▶  $P(\text{First student is female}) = \frac{17}{40}$ .
- ▶ Given sampling without replacement; first student selected is not returned to the class for reselection.

## Example: application of multiplication rule-cont.

- ▶ Event: First student female and second is male.
- ▶  $P(F_1 M_2) =$   
 $P(\text{First student is female} \cap \text{Second student is male})$
- ▶  $P(\text{First student is female}) = \frac{17}{40}$ .
- ▶ Given sampling without replacement; first student selected is not returned to the class for reselection.
- ▶ Given that the first student selected is female, of the 39 students remaining in the class 23 are male, so  
 $P(\text{Second student is male} | \text{First student is Female}) = \frac{23}{39}$

## Example: application of multiplication rule-cont.

- ▶ Event: First student female and second is male.
- ▶  $P(F_1 M_2) =$   
 $P(\text{First student is female} \cap \text{Second student is male})$
- ▶  $P(\text{First student is female}) = \frac{17}{40}$ .
- ▶ Given sampling without replacement; first student selected is not returned to the class for reselection.
- ▶ Given that the first student selected is female, of the 39 students remaining in the class 23 are male, so  
 $P(\text{Second student is male} | \text{First student is Female}) = \frac{23}{39}$
- ▶ Hence,  $P(\text{First student female and second is male}) =$   
 $P(\text{First student is female}) \times$   
 $P(\text{Second student is male} | \text{First student is Female}) = \frac{17}{40} \frac{23}{39} =$   
0.251

## Generalized multiplication rule

A generalization of the multiplication rule, which provides an expression for the probability of the intersection of an arbitrary number of events, is referred to as the generalized multiplication rule and is given by

## Generalized multiplication rule

A generalization of the multiplication rule, which provides an expression for the probability of the intersection of an arbitrary number of events, is referred to as the generalized multiplication rule and is given by  $P(E_1 \cap E_2 \cap \dots \cap E_n) = P(E_1)P(E_2|E_1)P(E_3|E_1 \cap E_2) \dots P(E_n|E_1 \cap E_2 \dots \cap E_{n-1})$

## Example: deck of cards

- ▶ An ordinary deck of 52 playing cards is randomly divided into 4 piles of 13 cards each. Compute the probability that each pile has exactly 1 ace.



## Example: deck of cards

- ▶ An ordinary deck of 52 playing cards is randomly divided into 4 piles of 13 cards each. Compute the probability that each pile has exactly 1 ace.
- ▶ Define events  $E_i; i = 1, 2, 3, 4$  as follows
  1.  $E_1 = \{\text{the ace of spades is in any one of the piles}\}$
  2.  $E_2 = \{\text{the ace of spades and the ace of hearts are in different pile}\}$
  3.  $E_3 = \{\text{the aces of spades, heart, and diamonds are in different piles}\}$
  4.  $E_4 = \{\text{all four aces are in different piles}\}$

## Example: deck of cards

- ▶ An ordinary deck of 52 playing cards is randomly divided into 4 piles of 13 cards each. Compute the probability that each pile has exactly 1 ace.
- ▶ Define events  $E_i; i = 1, 2, 3, 4$  as follows
  1.  $E_1 = \{\text{the ace of spades is in any one of the piles}\}$
  2.  $E_2 = \{\text{the ace of spades and the ace of hearts are in different pile}\}$
  3.  $E_3 = \{\text{the aces of spades, heart, and diamonds are in different piles}\}$
  4.  $E_4 = \{\text{all four aces are in different piles}\}$
- ▶ What we require is  $P(E_1 \cap E_2 \cap E_3 \cap E_4)$

# Solution

## Solution

1.  $P(E_1) = 1$

## Solution

1.  $P(E_1) = 1$

2.  $P(E_2|E_1) = \frac{39}{51}$

## Solution

1.  $P(E_1) = 1$
2.  $P(E_2|E_1) = \frac{39}{51}$
3.  $P(E_3|E_1 \cap E_2) = \frac{26}{50}$

## Solution

1.  $P(E_1) = 1$
2.  $P(E_2|E_1) = \frac{39}{51}$
3.  $P(E_3|E_1 \cap E_2) = \frac{26}{50}$
4.  $P(E_4|E_1 \cap E_2 \cap E_3) = \frac{13}{49}$

## Solution

1.  $P(E_1) = 1$
2.  $P(E_2|E_1) = \frac{39}{51}$
3.  $P(E_3|E_1 \cap E_2) = \frac{26}{50}$
4.  $P(E_4|E_1 \cap E_2 \cap E_3) = \frac{13}{49}$
5.  $P(E_1 \cap E_2 \cap E_3 \cap E_4) = \frac{39}{51} \frac{26}{50} \frac{13}{49} \approx 0.105$



## Section summary

1. Multiplication rule and its application to find probability of intersection of events.