

# Statistics for Data Science -1

## Lecture 5.4: Permutations formula: objects are not distinct & Circular permutations

Usha Mohan

Indian Institute of Technology Madras

## Learning objectives

1. Understand basic principles of counting.
2. Concept of factorials.
3. Understand differences between counting with order (permutation) and counting without regard to order (combination).
4. Use permutations and combinations to answer real life applications.

## Permutations

Permutation when objects are not distinct

## Circular permutations

Solving of  $n$  and  $r$  using permutation formula

## Example: Rearranging letters

## Example: Rearranging letters

- ▶ Suppose we want to rearrange the letters in the word "DATA". How many ways can it be done?

- ▶ Suppose we want to rearrange the letters in the word "DATA". How many ways can it be done?
- ▶ There are three distinct letters : D, A, T.

## Example: Rearranging letters

- ▶ Suppose we want to rearrange the letters in the word "DATA". How many ways can it be done?
- ▶ There are three distinct letters : D, A, T.
- ▶ Hence the possible arrangements taking all the four letters at a time are

## Example: Rearranging letters

- ▶ Suppose we want to rearrange the letters in the word "DATA". How many ways can it be done?
- ▶ There are three distinct letters : D, A, T.
- ▶ Hence the possible arrangements taking all the four letters at a time are

First place	Second place	Third place	Fourth place
A	D	T	A
A	D	A	T
A	T	D	A
A	T	A	D
A	A	D	T
A	A	T	D
D	A	T	A
D	A	A	T
D	T	A	A
T	A	D	A
T	A	A	D
T	D	A	A



## Permutations when objects are not distinct

- ▶ As seen in the example, we can treat the two A's in DATA as distinct. Say,  $A_1$  and  $A_2$ .

## Permutations when objects are not distinct

- ▶ As seen in the example, we can treat the two A's in DATA as distinct. Say,  $A_1$  and  $A_2$ .
- ▶ If they are treated as distinct objects, then based on the earlier formula, total number of arrangements =  $4!$ .

## Permutations when objects are not distinct

- ▶ As seen in the example, we can treat the two A's in DATA as distinct. Say,  $A_1$  and  $A_2$ .
- ▶ If they are treated as distinct objects, then based on the earlier formula, total number of arrangements =  $4!$ .
- ▶ Now  $A_1$  and  $A_2$  can be arranged among themselves in  $2!$  ways.

## Permutations when objects are not distinct

- ▶ As seen in the example, we can treat the two A's in DATA as distinct. Say,  $A_1$  and  $A_2$ .
- ▶ If they are treated as distinct objects, then based on the earlier formula, total number of arrangements =  $4!$ .
- ▶ Now  $A_1$  and  $A_2$  can be arranged among themselves in  $2!$  ways.
- ▶  $A_1$  and  $A_2$  are essentially the same. Hence, the total number of ways the letters in "DATA" can be arranged is  $\frac{4!}{2!} = 12$

## Permutation formula

- ▶ The number of permutations of  $n$  objects when  $p$  of them are of one kind and rest distinct is equal to

$$\frac{n!}{p!}$$

## Example

- ▶ Suppose we want to rearrange the letters in the word “STATISTICS”. How many ways can it be done?

## Example

- ▶ Suppose we want to rearrange the letters in the word “STATISTICS”. How many ways can it be done?

## Example

- ▶ Suppose we want to rearrange the letters in the word “STATISTICS”. How many ways can it be done?
- ▶ Total of ten letters of which there are five distinct letters : S,T,A,I,C.



## Example

- ▶ Suppose we want to rearrange the letters in the word “STATISTICS”. How many ways can it be done?
- ▶ Total of ten letters of which there are five distinct letters : S,T,A,I,C.

## Example

- ▶ Suppose we want to rearrange the letters in the word “STATISTICS”. How many ways can it be done?
- ▶ Total of ten letters of which there are five distinct letters : S,T,A,I,C.
- ▶ “S” appears 3 times; “T” appears 3 times, “A” once, “I” twice, and “C” once

## Permutation formula

- ▶ The number of permutations of  $n$  objects where  $p_1$  is of one kind,  $p_2$  is of second kind, and so on  $p_k$  of  $k^{th}$  kind is given by

## Permutation formula

- ▶ The number of permutations of  $n$  objects where  $p_1$  is of one kind,  $p_2$  is of second kind, and so on  $p_k$  of  $k^{th}$  kind is given by

$$\frac{n!}{p_1! p_2! \dots p_k!}$$

## Permutation formula

- ▶ The number of permutations of  $n$  objects where  $p_1$  is of one kind,  $p_2$  is of second kind, and so on  $p_k$  of  $k^{th}$  kind is given by

$$\frac{n!}{p_1! p_2! \dots p_k!}$$

- ▶ Applying the above formula to the word “STATISTICS”;

## Permutation formula

- ▶ The number of permutations of  $n$  objects where  $p_1$  is of one kind,  $p_2$  is of second kind, and so on  $p_k$  of  $k^{th}$  kind is given by

$$\frac{n!}{p_1! p_2! \dots p_k!}$$

- ▶ Applying the above formula to the word “STATISTICS”;  $n = 10, p_1 = 3, p_2 = 3, p_3 = 1, p_4 = 2, p_5 = 1$ .

## Permutation formula

- ▶ The number of permutations of  $n$  objects where  $p_1$  is of one kind,  $p_2$  is of second kind, and so on  $p_k$  of  $k^{th}$  kind is given by

$$\frac{n!}{p_1! p_2! \dots p_k!}$$

- ▶ Applying the above formula to the word "STATISTICS";  $n = 10, p_1 = 3, p_2 = 3, p_3 = 1, p_4 = 2, p_5 = 1$ . Hence, total number of ways =

$$\frac{10!}{3!3!1!2!1!} =$$

## Permutation formula

- ▶ The number of permutations of  $n$  objects where  $p_1$  is of one kind,  $p_2$  is of second kind, and so on  $p_k$  of  $k^{th}$  kind is given by

$$\frac{n!}{p_1! p_2! \dots p_k!}$$

- ▶ Applying the above formula to the word “STATISTICS”;  $n = 10, p_1 = 3, p_2 = 3, p_3 = 1, p_4 = 2, p_5 = 1$ . Hence, total number of ways =

$$\frac{10!}{3!3!1!2!1!} = 50,400$$



## Section summary





1. The number of permutations of  $n$  objects when  $p$  of them are of one kind and rest distinct is equal to  $\frac{n!}{p!}$
2. The number of permutations of  $n$  objects where  $p_1$  is of one kind,  $p_2$  is of second kind, and so on  $p_k$  of  $k^{th}$  kind is given by  $\frac{n!}{p_1!p_2!\dots p_k!}$

## Example


- ▶ How many ways can four people sit in a round table?
- ▶ We consider two cases: each selection is called a combination of 3 different objects taken 2 at a time.
  - ▶ Clockwise and anticlockwise are different
  - ▶ Clockwise and anticlockwise are same.

## Circular permutation: Clockwise and anticlockwise are different

- ▶ Consider the linear permutations of  $A, B, C$  and  $D$
- ▶ The arrangements  $ABCD, BCDA, CDAB$ , and  $DABC$  are different when the people are seated in a row.
- ▶ However, when they are seated in a circle as shown below:




## Circular permutation: Clockwise and anticlockwise are different



The number of ways  $n$  distinct objects can be arranged in a circle (clockwise and anticlockwise are different) is equal to  $(n - 1)!$

Circular permutation: Clockwise and anticlockwise are same



The number of ways  $n$  distinct objects can be arranged in a circle (clockwise and anticlockwise are same) is equal to  $\frac{(n-1)!}{2}$

## Example : Solving for $n$

- Find value of  $n$  if  ${}^nP_4 = 20{}^nP_2$

## Example : Solving for $n$

- Find value of  $n$  if  ${}^nP_4 = 20{}^nP_2$   
Answer:  $\frac{n!}{(n-4)!} = 20 \times \frac{n!}{(n-2)!}$

## Example : Solving for $n$

- Find value of  $n$  if  ${}^nP_4 = 20{}^nP_2$

Answer:  $\frac{n!}{(n-4)!} = 20 \times \frac{n!}{(n-2)!}$

Solving  $(n-2) \times (n-3) = 20$ , we get  $n = -2$  or  $n = 7$ .



## Example : Solving for $n$

- Find value of  $n$  if  ${}^nP_4 = 20 {}^nP_2$

Answer:  $\frac{n!}{(n-4)!} = 20 \times \frac{n!}{(n-2)!}$

Solving  $(n-2) \times (n-3) = 20$ , we get  $n = -2$  or  $n = 7$ .

Eliminating  $n = -2$ , we get  $n = 7$ .

## Example : Solving for $n$

$$\blacktriangleright \frac{{}^nP_4}{{}^{n-1}P_4} = \frac{5}{3}$$

$$\text{Answer: } \frac{n!}{(n-4)!} \times \frac{(n-5)!}{(n-1)!} = \frac{5}{3}$$

## Example : Solving for $n$

$$\blacktriangleright \frac{{}^nP_4}{{}^{n-1}P_4} = \frac{5}{3}$$

$$\text{Answer: } \frac{n!}{(n-4)!} \times \frac{(n-5)!}{(n-1)!} = \frac{5}{3}$$

$$\frac{n}{(n-4)} = \frac{5}{3}$$

## Example : Solving for $n$

$$\blacktriangleright \frac{{}^nP_4}{{}^{n-1}P_4} = \frac{5}{3}$$

$$\text{Answer: } \frac{n!}{(n-4)!} \times \frac{(n-5)!}{(n-1)!} = \frac{5}{3}$$

$$\frac{n}{(n-4)} = \frac{5}{3}$$

Solving for  $n$  gives us  $n = 10$ .

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻ 16/ 17

## Example : Solving for $r$

- Find  $r$ , if  ${}^5P_r = 2 \cdot {}^6P_{r-1}$   
Answer:  $\frac{5!}{(5-r)!} = 2 \cdot \frac{6!}{(7-r)!}$

## Example : Solving for $r$

► Find  $r$ , if  ${}^5P_r = 2 \cdot {}^6P_{r-1}$

Answer:  $\frac{5!}{(5-r)!} = 2 \cdot \frac{6!}{(7-r)!}$

$$\frac{5!}{(5-r)!} = 2 \cdot \frac{6!}{(7-r)(6-r)(5-r)!}$$

## Example : Solving for $r$

► Find  $r$ , if  ${}^5P_r = 2 \cdot {}^6P_{r-1}$

$$\text{Answer: } \frac{5!}{(5-r)!} = 2 \cdot \frac{6!}{(7-r)!}$$

$$\frac{5!}{(5-r)!} = 2 \cdot \frac{6!}{(7-r)(6-r)(5-r)!}$$

Solving  $(7-r)(6-r) = 12$  gives  $r = 10$  or  $r = 3$ .

Since  $r \leq n$ , the option  $r = 10$  is eliminated and we get  $r = 3$ .



## Topic summary

1. Permutations when objects are distinct
  - 1.1 repetitions not allowed.
  - 1.2 repetitions allowed.
2. Permutations when objects are not distinct.
3. Circular permutations:
  - 3.1 Clockwise and anticlockwise are different.
  - 3.2 Clockwise and anticlockwise are same.
4. Solving for  $r$  and  $n$  using the permutation formula.