Statistics for Data Science -1

Lecture 10.5: Expectation and Variance of Binomial distribution

Usha Mohan

Indian Institute of Technology Madras

1. Derive the formula for the probability mass function for Binomial distribution.

1. Derive the formula for the probability mass function for Binomial distribution.

- Derive the formula for the probability mass function for Binomial distribution.
- 2. Understand the effect of parameters n and p on the shape of the Binomial distribution.

- 1. Derive the formula for the probability mass function for Binomial distribution.
- 2. Understand the effect of parameters n and p on the shape of the Binomial distribution.
- 3. Expectation and variance of the binomial distribution.

- Derive the formula for the probability mass function for Binomial distribution.
- 2. Understand the effect of parameters n and p on the shape of the Binomial distribution.
- 3. Expectation and variance of the binomial distribution.
- 4. To understand situations that can be modeled as a Binomial distribution.

▶ A binomial random variable $X \sim Bin(n, p)$ is equal to the number of successes in n independent trials when each trial is a success with probability p.

- A binomial random variable $X \sim Bin(n, p)$ is equal to the number of successes in n independent trials when each trial is a success with probability p.
- \triangleright We can represent X as

$$X = X_1 + X_2 + \ldots + X_n$$

where X_i is equal to 1 if trial i is a success and is equal to 0 if trial i is a failure.

- A binomial random variable $X \sim Bin(n, p)$ is equal to the number of successes in n independent trials when each trial is a success with probability p.
- ightharpoonup We can represent X as

$$X = X_1 + X_2 + \ldots + X_n$$

where X_i is equal to 1 if trial i is a success and is equal to 0 if trial i is a failure.

 $P(X_i = 1) = p \text{ and } P(X_i = 0) = (1 - p)$

$$\triangleright X = X_1 + X_2 + \ldots + X_n$$

- $X = X_1 + X_2 + \ldots + X_n$
- $E(X) = E(X_1 + X_2 + \ldots + X_n) = E(X_1) + E(X_2) + \ldots + E(X_n)$

- $X = X_1 + X_2 + \ldots + X_n$
- $E(X) = E(X_1 + X_2 + \ldots + X_n) = E(X_1) + E(X_2) + \ldots + E(X_n)$
 - ightharpoonup E(X) = p + p + ... + p = np

- $X = X_1 + X_2 + ... + X_n$
- $E(X) = E(X_1 + X_2 + ... + X_n) = E(X_1) + E(X_2) + ... + E(X_n)$
 - $ightharpoonup E(X) = p + p + \ldots + p = np$
- $V(X) = V(X_1 + X_2 + ... + X_n) = V(X_1) + V(X_2) + ... + V(X_n)$

- $X = X_1 + X_2 + ... + X_n$
- $E(X) = E(X_1 + X_2 + \ldots + X_n) = E(X_1) + E(X_2) + \ldots + E(X_n)$
 - ► E(X) = p + p + ... + p = np
- $V(X) = V(X_1 + X_2 + ... + X_n) = V(X_1) + V(X_2) + ... + V(X_n)$
 - V(X) = p(1-p) + p(1-p) + ... + p(1-p) = np(1-p)

- $X = X_1 + X_2 + ... + X_n$
- $E(X) = E(X_1 + X_2 + ... + X_n) = E(X_1) + E(X_2) + ... + E(X_n)$ E(X) = p + p + ... + p = np
- $V(X) = V(X_1 + X_2 + ... + X_n) = V(X_1) + V(X_2) + ... + V(X_n)$
 - V(X) = p(1-p) + p(1-p) + ... + p(1-p) = np(1-p)

Result

Using the fact that the expectation of the sum of random variables is equal to the sum of their expectations, we see that Expectation of a Binomial random variable X is

$$E(X) = np$$

Also, since the variance of the sum of independent random variables is equal to the sum of their variances, we have variance of a Binomial random variable is

$$Var(X) = np(1-p)$$

If a fair coin is tossed 500 times, what is the standard deviation of the number of times that a head appears?

If a fair coin is tossed 500 times, what is the standard deviation of the number of times that a head appears?

Let X = the number of heads in 500 tosses of a fair coin.

If a fair coin is tossed 500 times, what is the standard deviation of the number of times that a head appears?

Let X = the number of heads in 500 tosses of a fair coin. Then $X \sim B(500, 1/2)$.

If a fair coin is tossed 500 times, what is the standard deviation of the number of times that a head appears?

Let X = the number of heads in 500 tosses of a fair coin. Then $X = R(500, 1/2) V(X) = 125 SD(X) = \sqrt{125} = 11,1803$

$$X \sim B(500, 1/2).V(X) = 125, SD(X) = \sqrt{125} = 11.1803$$

The expected number of heads in a series of 10 tosses of a coin is 6. What is the probability there are 8 heads?

The expected number of heads in a series of 10 tosses of a coin is 6. What is the probability there are 8 heads? Let X be number of heads.

The expected number of heads in a series of 10 tosses of a coin is 6. What is the probability there are 8 heads? Let X be number of heads. $X \sim B(10, p)$.

The expected number of heads in a series of 10 tosses of a coin is 6. What is the probability there are 8 heads? Let X be number of heads. $X \sim B(10, p)$.

1. Since
$$E(X) = np$$
; $10p = 6$, hence $p = 0.6$

The expected number of heads in a series of 10 tosses of a coin is 6. What is the probability there are 8 heads? Let X be number of heads. $X \sim B(10, p)$.

- 1. Since E(X) = np; 10p = 6, hence p = 0.6
- 2. Prob there are 8 heads; P(X = 8) = 0.121

a
$$P(X = 3)$$

- a P(X = 3)
- b $P(X \ge 4)$

- a P(X = 3)
- b $P(X \ge 4)$
- \triangleright $X \sim B(n, p)$

- a P(X = 3)
- b $P(X \ge 4)$
- $\triangleright X \sim B(n,p)$
- ightharpoonup np = 4.5, np(1-p) = 0.45.

a
$$P(X = 3)$$

b
$$P(X \ge 4)$$

$$\triangleright X \sim B(n,p)$$

$$ightharpoonup np = 4.5, np(1-p) = 0.45.$$

► Solving gives
$$n = 5$$
 and $p = 0.9$

a
$$P(X = 3) = 0.0729$$

b
$$P(X > 4) = 0.9185$$

Section summary

- Expectation and variance of Binomial random variable
- Applications