

Statistics for Data Science -1

Lecture 6.6: Probability- Equally likely outcomes

Usha Mohan

Indian Institute of Technology Madras

Learning objectives

Learning objectives

1. Understand uncertainty and concept of a random experiment.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.
6. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.
6. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
7. Distinguish between independent and dependent events.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.
6. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
7. Distinguish between independent and dependent events.
8. Solve applications of probability.

Equally likely outcomes

Equally likely outcomes

Equally likely outcomes

- ▶ For certain experiments it is natural to assume that each outcome in the sample space S is equally likely to occur.

Equally likely outcomes

- ▶ For certain experiments it is natural to assume that each outcome in the sample space S is equally likely to occur.
- ▶ That is, if sample space S consists of N outcomes, say, $S = \{1, 2, \dots, N\}$, then it is often reasonable to suppose that

Equally likely outcomes

- ▶ For certain experiments it is natural to assume that each outcome in the sample space S is equally likely to occur.
- ▶ That is, if sample space S consists of N outcomes, say, $S = \{1, 2, \dots, N\}$, then it is often reasonable to suppose that

$$P(\{1\}) = P(\{2\}) = \dots = P(\{N\})$$

- ▶ In this expression, $P(\{i\})$ is the probability of the event consisting of the single outcome i .

Equally likely outcomes

- ▶ For certain experiments it is natural to assume that each outcome in the sample space S is equally likely to occur.
- ▶ That is, if sample space S consists of N outcomes, say, $S = \{1, 2, \dots, N\}$, then it is often reasonable to suppose that

$$P(\{1\}) = P(\{2\}) = \dots = P(\{N\})$$

- ▶ In this expression, $P(\{i\})$ is the probability of the event consisting of the single outcome i .
- ▶ Using the properties of probability, we can show that the foregoing implies that the probability of any event A is equal to the proportion of the outcomes in the sample space that is in A .

Equally likely outcomes

- ▶ For certain experiments it is natural to assume that each outcome in the sample space S is equally likely to occur.
- ▶ That is, if sample space S consists of N outcomes, say, $S = \{1, 2, \dots, N\}$, then it is often reasonable to suppose that

$$P(\{1\}) = P(\{2\}) = \dots = P(\{N\})$$

- ▶ In this expression, $P(\{i\})$ is the probability of the event consisting of the single outcome i .
- ▶ Using the properties of probability, we can show that the foregoing implies that the probability of any event A is equal to the proportion of the outcomes in the sample space that is in A .
- ▶ That is, $P(A) =$

Equally likely outcomes

- ▶ For certain experiments it is natural to assume that each outcome in the sample space S is equally likely to occur.
- ▶ That is, if sample space S consists of N outcomes, say, $S = \{1, 2, \dots, N\}$, then it is often reasonable to suppose that

$$P(\{1\}) = P(\{2\}) = \dots = P(\{N\})$$

- ▶ In this expression, $P(\{i\})$ is the probability of the event consisting of the single outcome i .
- ▶ Using the properties of probability, we can show that the foregoing implies that the probability of any event A is equal to the proportion of the outcomes in the sample space that is in A .
- ▶ That is, $P(A) = \frac{\text{number of outcomes in } S \text{ that are in } A}{N}$

Example: Rolling a dice

Example: Rolling a dice

- ▶ Experiment: Roll a fair dice

Example: Rolling a dice

- ▶ Experiment: Roll a fair dice
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$

Example: Rolling a dice

- ▶ Experiment: Roll a fair dice
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Let E_i denote the event of outcome i . Since the dice is fair,
 $P(E_i) = \frac{1}{6}$.

Example: Rolling a dice

- ▶ Experiment: Roll a fair dice
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Let E_i denote the event of outcome i . Since the dice is fair, $P(E_i) = \frac{1}{6}$.
- ▶ Define A to be the event the outcome is odd $A = \{1, 3, 5\}$

Example: Rolling a dice

- ▶ Experiment: Roll a fair dice
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Let E_i denote the event of outcome i . Since the dice is fair, $P(E_i) = \frac{1}{6}$.
- ▶ Define A to be the event the outcome is odd $A = \{1, 3, 5\}$
 $P(A) = P(E_1) + P(E_3) + P(E_5) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2}$

Example: Rolling a dice

- ▶ Experiment: Roll a fair dice
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Let E_i denote the event of outcome i . Since the dice is fair, $P(E_i) = \frac{1}{6}$.
- ▶ Define A to be the event the outcome is odd $A = \{1, 3, 5\}$
 $P(A) = P(E_1) + P(E_3) + P(E_5) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2}$
- ▶ Let B be the event that the outcome is greater than 4.

Example: Rolling a dice


- ▶ Experiment: Roll a fair dice
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Let E_i denote the event of outcome i . Since the dice is fair, $P(E_i) = \frac{1}{6}$.
- ▶ Define A to be the event the outcome is odd $A = \{1, 3, 5\}$
 $P(A) = P(E_1) + P(E_3) + P(E_5) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2}$
- ▶ Let B be the event that the outcome is greater than 4.
 $B = \{5, 6\}$ $P(B) = \frac{2}{6}$

Example: Rolling a dice

- ▶ Experiment: Roll a fair dice
- ▶ Sample space: $S = \{1, 2, 3, 4, 5, 6\}$
- ▶ Let E_i denote the event of outcome i . Since the dice is fair, $P(E_i) = \frac{1}{6}$.
- ▶ Define A to be the event the outcome is odd $A = \{1, 3, 5\}$
 $P(A) = P(E_1) + P(E_3) + P(E_5) = \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2}$
- ▶ Let B be the event that the outcome is greater than 4.
 $B = \{5, 6\}$ $P(B) = \frac{2}{6}$
- ▶ Let C be the event that the outcome is either odd or greater than 4.
 $P(C) = P(A \cup B) = P(A) + P(B) - P(A \cap B) = \frac{3}{6} + \frac{2}{6} - \frac{1}{6} = \frac{4}{6}$


Example: playing cards

When drawing a card from a standard deck of 52 playing cards, what is the probability that the card is either red or a queen?



Example: playing cards

When drawing a card from a standard deck of 52 playing cards, what is the probability that the card is either red or a queen?




- Let R be the event that the card drawn is Red.

$$P(R) = \frac{26}{52} = \frac{1}{2}$$

Example: playing cards

When drawing a card from a standard deck of 52 playing cards, what is the probability that the card is either red or a queen?



- ▶ Let R be the event that the card drawn is Red.

$$P(R) = \frac{26}{52} = \frac{1}{2}$$

- ▶ Let Q be the event that the card drawn is Queen.

$$P(Q) = \frac{4}{52} = \frac{1}{13}$$

Example: playing cards-contd.

Probability that the card is either red or a queen = $P(R \cup Q)$

Example: playing cards-contd.

Probability that the card is either red or a queen = $P(R \cup Q)$

- ▶ Applying addition rule: $P(R \cup Q) = P(R) + P(Q) - P(R \cap Q)$

Example: playing cards-contd.

Probability that the card is either red or a queen = $P(R \cup Q)$

- ▶ Applying addition rule: $P(R \cup Q) = P(R) + P(Q) - P(R \cap Q)$
- ▶ $R \cap Q$ describes the event that the card drawn is a Red Queen. $P(R \cap Q) = \frac{2}{56}$

Example: playing cards-contd.

Probability that the card is either red or a queen = $P(R \cup Q)$

- ▶ Applying addition rule: $P(R \cup Q) = P(R) + P(Q) - P(R \cap Q)$
- ▶ $R \cap Q$ describes the event that the card drawn is a Red Queen. $P(R \cap Q) = \frac{2}{52}$
- ▶ Hence $P(R \cup Q) = \frac{26}{52} + \frac{4}{52} - \frac{2}{52} = \frac{28}{52} = \frac{7}{13}$

Topic summary

1. Interpretations of probability
2. Probability axioms
3. Addition rule of probability.
4. Equally likely outcomes.