

# Statistics for Data Science -1

## Continuous Random Variables-Continuous distributions

Usha Mohan

Indian Institute of Technology Madras


## Exponential distribution

A continuous random variable whose probability density function is given, for some  $\lambda > 0$ , by

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

is said to be an exponential random variable (or, more simply, is said to be exponentially distributed) with parameter  $\lambda$ .

## Graph of pdf for different values of $\lambda$



## cdf of Exponential distribution

$$F(a) = P(X \leq a)$$

## cdf of Exponential distribution

$$\begin{aligned} F(a) &= P(X \leq a) \\ &= \int_0^a \lambda e^{-\lambda x} dx \end{aligned}$$


## cdf of Exponential distribution

$$\begin{aligned} F(a) &= P(X \leq a) \\ &= \int_0^a \lambda e^{-\lambda x} dx \\ &= -e^{-\lambda x} \Big|_0^a \end{aligned}$$

## cdf of Exponential distribution

$$\begin{aligned} F(a) &= P(X \leq a) \\ &= \int_0^a \lambda e^{-\lambda x} dx \\ &= -e^{-\lambda x} \Big|_0^a \\ &= 1 - e^{-\lambda a} \end{aligned}$$

### Graph of cdf for different values of $\lambda$





## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

$$\blacktriangleright E(X) = \frac{1}{\lambda}$$

## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

$$\blacktriangleright E(X) = \frac{1}{\lambda}$$

$$\blacktriangleright \text{Var}(X) = \frac{1}{\lambda^2}$$

## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $\text{Var}(X) = \frac{1}{\lambda^2}$
- ▶ It can be shown through integration by parts

$$E(X^n) = \frac{n}{\lambda} E(X^{n-1})$$

## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $\text{Var}(X) = \frac{1}{\lambda^2}$
- ▶ It can be shown through integration by parts

$$E(X^n) = \frac{n}{\lambda} E(X^{n-1})$$

- ▶  $E(X) = \frac{1}{\lambda}$

## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $\text{Var}(X) = \frac{1}{\lambda^2}$
- ▶ It can be shown through integration by parts

$$E(X^n) = \frac{n}{\lambda} E(X^{n-1})$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $E(X^2) = \frac{2}{\lambda} \frac{1}{\lambda} = \frac{2}{\lambda^2}$

## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $\text{Var}(X) = \frac{1}{\lambda^2}$
- ▶ It can be shown through integration by parts

$$E(X^n) = \frac{n}{\lambda} E(X^{n-1})$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $E(X^2) = \frac{2}{\lambda} \frac{1}{\lambda} = \frac{2}{\lambda^2}$
- ▶ Hence  $\text{Var}(X) = \frac{2}{\lambda^2} - \frac{1}{\lambda^2} = \frac{1}{\lambda^2}$

## Expectation and variance of exponential distribution

$$X \sim \exp(\lambda)$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $\text{Var}(X) = \frac{1}{\lambda^2}$
- ▶ It can be shown through integration by parts

$$E(X^n) = \frac{n}{\lambda} E(X^{n-1})$$

- ▶  $E(X) = \frac{1}{\lambda}$
- ▶  $E(X^2) = \frac{2}{\lambda} \frac{1}{\lambda} = \frac{2}{\lambda^2}$
- ▶ Hence  $\text{Var}(X) = \frac{2}{\lambda^2} - \frac{1}{\lambda^2} = \frac{1}{\lambda^2}$

Thus, the mean of the exponential is the reciprocal of its parameter  $\lambda$ , and the variance is the mean squared.



## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

Solution Let  $X$  denote the length of the call made by the person in the booth.  $X \sim \exp(0.5)$

## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

Solution Let  $X$  denote the length of the call made by the person in the booth.  $X \sim \exp(0.5)$

- a more than 10 minutes =

## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

Solution Let  $X$  denote the length of the call made by the person in the booth.  $X \sim \text{exp}(0.1)$

a more than 10 minutes =  $P(X > 10) = e^{-1} \approx 0.368$

## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

Solution Let  $X$  denote the length of the call made by the person in the booth.  $X \sim \exp(0.1)$

- a more than 10 minutes =  $P(X > 10) = e^{-1} \approx 0.368$
- b between 10 and 20 minutes =

## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

Solution Let  $X$  denote the length of the call made by the person in the booth.  $X \sim \exp(0.1)$

- a more than 10 minutes =  $P(X > 10) = e^{-1} \approx 0.368$
- b between 10 and 20 minutes =  
 $P(10 < X < 20) = F(20) - F(10)$

## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

Solution Let  $X$  denote the length of the call made by the person in the booth.  $X \sim \exp(0.1)$

a more than 10 minutes =  $P(X > 10) = e^{-1} \approx 0.368$

b between 10 and 20 minutes =  
 $P(10 < X < 20) = F(20) - F(10) = e^{-1} - e^{-2} \approx$

## Application

In practice, the exponential distribution often arises as the distribution of the amount of time until some specific event occurs.

- ▶ Suppose that the length of a phone call in minutes is an exponential random variable with parameter  $\lambda = 0.1$ . If someone arrives immediately ahead of you at a public telephone booth, find the probability that you will have to wait
  - a more than 10 minutes
  - b between 10 and 20 minutes.

Solution Let  $X$  denote the length of the call made by the person in the booth.  $X \sim \exp(0.1)$

- a more than 10 minutes =  $P(X > 10) = e^{-1} \approx 0.368$
- b between 10 and 20 minutes =  
 $P(10 < X < 20) = F(20) - F(10) = e^{-1} - e^{-2} \approx 0.233$



## Section summary

- ▶ Exponential distribution and its applications.