

Statistics for Data Science -1

Lecture 7.2: Conditional Probability: Definition

Usha Mohan

Indian Institute of Technology Madras

Learning objectives

1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.

Learning objectives

1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
2. Distinguish between independent and dependent events.

Learning objectives

1. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
2. Distinguish between independent and dependent events.
3. Solve applications of probability.

Conditional Probability

Multiplication rule

Independent events

Bayes' rule

Introduction

- ▶ We are often interested in determining probabilities when some partial information concerning the outcome of the experiment is available. In such situations, the probabilities are called **conditional probabilities**.

Example: Roll a dice twice

Example: Roll a dice twice

- ▶ Experiment: Roll a dice twice

Example: Roll a dice twice

- ▶ Experiment: Roll a dice twice

- ▶ Sample space:

$$S = \left\{ \begin{array}{l} (1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6), \\ (2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), \\ (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6), \\ (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6), \\ (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6), \\ (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6), \end{array} \right\}$$

- ▶ Each outcome is **equally likely** to occur with a probability of $\frac{1}{36}$

Example: Rolling a dice twice- contd.

Example: Rolling a dice twice- contd.

- ▶ Suppose further that the first roll of the dice lands on 4.

Example: Rolling a dice twice- contd.

- ▶ Suppose further that the first roll of the dice lands on 4.
- ▶ Given this information, what is the resulting probability that the sum of the dice is 10?

Example: Rolling a dice twice- contd.

- ▶ Suppose further that the first roll of the dice lands on 4.
- ▶ Given this information, what is the resulting probability that the sum of the dice is 10?
- ▶ In other words, the restricted sample space if the first dice lands of a four $F = \{(4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6)\}$
- ▶ If each outcome of a finite sample space S is equally

Example: Rolling a dice twice- contd.

- ▶ Suppose further that the first roll of the dice lands on 4.
- ▶ Given this information, what is the resulting probability that the sum of the dice is 10?
- ▶ In other words, the restricted sample space if the first dice lands of a four $F = \{(4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6)\}$
- ▶ If each outcome of a finite sample space S is equally likely, then, conditional on the event that the outcome lies in a subset F , all outcomes in F become equally likely.

Example: Rolling a dice twice- contd.

- ▶ Suppose further that the first roll of the dice lands on 4.
- ▶ Given this information, what is the resulting probability that the sum of the dice is 10?
- ▶ In other words, the restricted sample space if the first dice lands of a four $F = \{(4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6)\}$
- ▶ If each outcome of a finite sample space S is equally likely, then, conditional on the event that the outcome lies in a subset F , all outcomes in F become equally likely. In such cases, it is often convenient to compute conditional probabilities of the form $P(E|F)$ by using F as the sample space.

Example: Rolling a dice twice- contd.

- ▶ Suppose further that the first roll of the dice lands on 4.
- ▶ Given this information, what is the resulting probability that the sum of the dice is 10?
- ▶ In other words, the restricted sample space if the first dice lands of a four $F = \{(4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6)\}$
- ▶ If each outcome of a finite sample space S is equally likely, then, conditional on the event that the outcome lies in a subset F , all outcomes in F become equally likely. In such cases, it is often convenient to compute conditional probabilities of the form $P(E|F)$ by using F as the sample space.
- ▶ Among outcomes in the restricted sample space, the outcome that satisfies the sum of dice is 10 is outcome $(4, 6)$. And this happens with Probability $\frac{1}{6}$

Conditional Probability: formula

Conditional Probability: formula

- ▶ Let E denote the event that the sum of the dice is 10 and let F denote the event that the first die lands on 4, then the probability obtained is called the **conditional probability** of E given that F has occurred. It is denoted by

$$P(E|F)$$

Conditional Probability: formula

- ▶ Let E denote the event that the sum of the dice is 10 and let F denote the event that the first die lands on 4, then the probability obtained is called the **conditional probability** of E given that F has occurred. It is denoted by

$$P(E|F)$$

- ▶ The probability that event E occurs given that event F occurs (or conditional on event F occurring) is given by

Conditional Probability: formula


- ▶ Let E denote the event that the sum of the dice is 10 and let F denote the event that the first die lands on 4, then the probability obtained is called the **conditional probability** of E given that F has occurred. It is denoted by

$$P(E|F)$$

- ▶ The probability that event E occurs given that event F occurs (or conditional on event F occurring) is given by

$$P(E|F) = \frac{P(E \cap F)}{P(F)}; P(F) > 0$$

Conditional probability: Venn diagram illustration



$$P(E|F) = \frac{P(E \cap F)}{P(F)}$$

Apply the formula to the example

Apply the formula to the example

- ▶ As a further check of the preceding formula for the conditional probability, use it to compute the conditional probability that the sum of a pair of rolled dice is 10, given that the first die lands on 4.

Apply the formula to the example

- ▶ As a further check of the preceding formula for the conditional probability, use it to compute the conditional probability that the sum of a pair of rolled dice is 10, given that the first die lands on 4.
- ▶
$$P(E|F) = \frac{P(E \cap F)}{P(F)}$$

Apply the formula to the example

- ▶ As a further check of the preceding formula for the conditional probability, use it to compute the conditional probability that the sum of a pair of rolled dice is 10, given that the first die lands on 4.
- ▶ $P(E|F) = \frac{P(E \cap F)}{P(F)}$
- ▶ $\frac{P(E \cap F)}{P(F)} = \frac{P(\{(4,6)\})}{P(\{(4,1),(4,2),(4,3),(4,4),(4,5),(4,6)\})} = \frac{1/36}{6/36} = \frac{1}{6}$

Section summary

1. Introduced notion of conditional probability
2. Formula: $P(E|F) = \frac{P(E \cap F)}{P(F)}$