

Riemann sums and the integral

Sarang S. Sane




Strategy to compute areas



Strategy to compute areas


Suppose you want to compute the area of a given shape.



Strategy to compute areas

Suppose you want to compute the area of a given shape.


1. Break the shape you have into rectangles. Some part of the shape may be left out, some extra part may get included.



Strategy to compute areas

Suppose you want to compute the area of a given shape.

1. Break the shape you have into rectangles. Some part of the shape may be left out, some extra part may get included.
2. Calculate the area covered by the rectangles.



Strategy to compute areas

Suppose you want to compute the area of a given shape.

1. Break the shape you have into rectangles. Some part of the shape may be left out, some extra part may get included.
2. Calculate the area covered by the rectangles.
3. Decrease the sizes of the rectangles used. The part of the shape left out and the extra part keep decreasing and recalculate.

Strategy to compute areas

Suppose you want to compute the area of a given shape.


1. Break the shape you have into rectangles. Some part of the shape may be left out, some extra part may get included.
2. Calculate the area covered by the rectangles.
3. Decrease the sizes of the rectangles used. The part of the shape left out and the extra part keep decreasing and recalculate.
4. As the sizes decrease, the area of the shape is better and better approximated.

Strategy to compute areas

Suppose you want to compute the area of a given shape.


1. Break the shape you have into rectangles. Some part of the shape may be left out, some extra part may get included.
2. Calculate the area covered by the rectangles.
3. Decrease the sizes of the rectangles used. The part of the shape left out and the extra part keep decreasing and recalculate.
4. As the sizes decrease, the area of the shape is better and better approximated.
5. In the **limit**, you will get the area of the shape.

The area under the graph of a function




Graph of $y = e^{-x}$

Riemann sums



Riemann sums


Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.



Riemann sums

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.

Suppose the interval $[a, b]$ is in the domain D .




Riemann sums

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.

Suppose the interval $[a, b]$ is in the domain D .

Let P consist of the following data :



Riemann sums

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.

Suppose the interval $[a, b]$ is in the domain D .

Let P consist of the following data :

- ▶ a partition of $[a, b]$, i.e. a choice of intermediate points
 $a = x_0 < x_1 < \dots < x_n = b$

Riemann sums

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.

Suppose the interval $[a, b]$ is in the domain D .

Let P consist of the following data :

- ▶ a partition of $[a, b]$, i.e. a choice of intermediate points
 $a = x_0 < x_1 < \dots < x_n = b$
- ▶ a choice of $x_i^* \in [x_{i-1}, x_i]$

Riemann sums

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.

Suppose the interval $[a, b]$ is in the domain D .

Let P consist of the following data :

- ▶ a partition of $[a, b]$, i.e. a choice of intermediate points
 $a = x_0 < x_1 < \dots < x_n = b$
- ▶ a choice of $x_i^* \in [x_{i-1}, x_i]$

Define $\Delta x_i = x_i - x_{i-1}$ and $\|P\| = \max_i \{\Delta x_i\}$.

Riemann sums

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.

Suppose the interval $[a, b]$ is in the domain D .

Let P consist of the following data :

- ▶ a partition of $[a, b]$, i.e. a choice of intermediate points
 $a = x_0 < x_1 < \dots x_n = b$
- ▶ a choice of $x_i^* \in [x_{i-1}, x_i]$

Define $\Delta x_i = x_i - x_{i-1}$ and $\|P\| = \max_i \{\Delta x_i\}$.

The **Riemann sum** of f w.r.t. the above data is defined as


$$S(P) = \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

The definite integral of a function



The definite integral of a function


Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.
Suppose the interval $[a, b]$ is in the domain D .



The definite integral of a function

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.
Suppose the interval $[a, b]$ is in the domain D .

The (definite) integral of f from a to b is defined as



The definite integral of a function

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.
Suppose the interval $[a, b]$ is in the domain D .

The (definite) integral of f from a to b is defined as

$$\lim_{\|P\| \rightarrow \infty} S(P) = \lim_{\|P\| \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

The definite integral of a function

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.
Suppose the interval $[a, b]$ is in the domain D .

The (definite) integral of f from a to b is defined as

$$\lim_{\|P\| \rightarrow \infty} S(P) = \lim_{\|P\| \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

It is denoted by $\int_a^b f(x) dx$.

The definite integral of a function

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.
Suppose the interval $[a, b]$ is in the domain D .

The (definite) integral of f from a to b is defined as

$$\lim_{\|P\| \rightarrow \infty} S(P) = \lim_{\|P\| \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

It is denoted by $\int_a^b f(x) dx$.

Let $f \geq 0$ (resp. $f \leq 0$) be piecewise continuous on this interval.

The definite integral of a function

Let f be a function from D to \mathbb{R} for some domain $D \subseteq \mathbb{R}$.
Suppose the interval $[a, b]$ is in the domain D .


The (definite) integral of f from a to b is defined as

$$\lim_{\|P\| \rightarrow \infty} S(P) = \lim_{\|P\| \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x_i.$$

It is denoted by $\int_a^b f(x) dx$.

Let $f \geq 0$ (resp. $f \leq 0$) be piecewise continuous on this interval.
Then the area under the graph of the function $\Gamma(f)$ above the
interval $[a, b]$ is measured by $\int_a^b f(x) dx$ (resp. $-\int_a^b f(x) dx$).

Example



$$P_n = \left\{ x_0 = 1, x_1 = 1 + \frac{1}{n}, x_2 = 1 + \frac{2}{n}, \dots, x_n = 2 \right.$$

$$x_i = 1 + \frac{i}{n}.$$

$1 = x_0 < x_1 < x_2 < \dots < x_n = 2.$

$$\Delta x_i = \frac{1}{n}. \quad \|P_n\| = \frac{1}{n}.$$

$$x_i^* = x_i = 1 + \frac{i}{n}$$

$$S(P_n) = \sum_{i=1}^n f(x_i^*) \Delta x_i$$

$$\begin{aligned}
 &= \sum_{i=1}^n \left\{ 2 \left(1 + \frac{i}{n} \right) - 1 \right\} \frac{1}{n} \\
 &= \sum_{i=1}^n \left(1 + \frac{2i}{n} \right) \frac{1}{n} \\
 &= \sum_{i=1}^n \frac{1}{n} + \sum_{i=1}^n \frac{2i}{n^2}
 \end{aligned}$$

$$\begin{aligned}
 &= 1 + \frac{2}{n^2} \sum_{i=1}^n i \\
 &= 1 + \frac{2}{n^2} \frac{n(n+1)}{2} \\
 &= 1 + \frac{n+1}{n} = 2 + \frac{1}{n} \\
 \lim_{\|P_n\| \rightarrow 0} S(P_n) &= \lim_{n \rightarrow \infty} 2 + \frac{1}{n} = 2.
 \end{aligned}$$

Thank you

