

Computing derivatives and L'Hôpital's rule

Sarang S. Sane

Recall

Definition

Let f be a function defined on an open interval around a .

Recall

Definition

Let f be a function defined on an open interval around a . Then f is **differentiable at a** if $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ exists.

Recall

Definition

Let f be a function defined on an open interval around a . Then f is **differentiable at a** if $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ exists.

Definition

For a function $f(x)$ its **derivative** function, $f'(x)$ or $\frac{df}{dx}(x)$ is

$$f'(x) = \frac{df}{dx}(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

Recall

Definition

Let f be a function defined on an open interval around a . Then f is **differentiable at a** if $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ exists.

Definition

For a function $f(x)$ its **derivative** function, $f'(x)$ or $\frac{df}{dx}(x)$ is

$$f'(x) = \frac{df}{dx}(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

Its domain consists of those points at which the function $f(x)$ is differentiable.

Recall

Definition

Let f be a function defined on an open interval around a . Then f is **differentiable at a** if $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ exists.

Definition

For a function $f(x)$ its **derivative** function, $f'(x)$ or $\frac{df}{dx}(x)$ is

$$f'(x) = \frac{df}{dx}(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

Its domain consists of those points at which the function $f(x)$ is differentiable.

Fact

If f is differentiable at a , then it is continuous at a .

Useful rules about derivatives

Useful rules about derivatives

Linearity :

Useful rules about derivatives

Linearity :

1. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f + g)(x)$ and $(f + g)'(a) = f'(a) + g'(a)$.

Useful rules about derivatives

Linearity :

1. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f + g)(x)$ and $(f + g)'(a) = f'(a) + g'(a)$.
2. If $f(x)$ is differentiable at the point a and $c \in \mathbb{R}$, then $(cf)(x)$ is also differentiable at the point a and $(cf)'(a) = cf'(a)$.

Useful rules about derivatives

Linearity :

1. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f + g)(x)$ and $(f + g)'(a) = f'(a) + g'(a)$.
2. If $f(x)$ is differentiable at the point a and $c \in \mathbb{R}$, then $(cf)(x)$ is also differentiable at the point a and $(cf)'(a) = cf'(a)$.
3. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f - g)(x)$ and $(f - g)'(a) = f'(a) - g'(a)$.

Useful rules about derivatives

Linearity :

1. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f + g)(x)$ and $(f + g)'(a) = f'(a) + g'(a)$.
2. If $f(x)$ is differentiable at the point a and $c \in \mathbb{R}$, then $(cf)(x)$ is also differentiable at the point a and $(cf)'(a) = cf'(a)$.
3. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f - g)(x)$ and $(f - g)'(a) = f'(a) - g'(a)$.

The product rule

Useful rules about derivatives

Linearity :

1. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f + g)(x)$ and $(f + g)'(a) = f'(a) + g'(a)$.
2. If $f(x)$ is differentiable at the point a and $c \in \mathbb{R}$, then $(cf)(x)$ is also differentiable at the point a and $(cf)'(a) = cf'(a)$.
3. If $f(x)$ and $g(x)$ are differentiable at the point a , then so is $(f - g)(x)$ and $(f - g)'(a) = f'(a) - g'(a)$.

The product rule

If $f(x)$ and $g(x)$ are differentiable at a , then so is $(fg)(x)$ and

$$(fg)'(a) = f'(a)g(a) + f(a)g'(a).$$
$$\lim_{h \rightarrow 0} \frac{f(a+h)g(a+h) - f(a)g(a)}{h} = \lim_{h \rightarrow 0} \frac{f(a+h)(g(a+h) - g(a)) + (f(a+h) - f(a))g(a)}{h}$$

Useful rules about derivatives (contd.)

Useful rules about derivatives (contd.)

The quotient rule

Useful rules about derivatives (contd.)

The quotient rule

If $f(x)$ and $g(x)$ are differentiable at the point a and $g(a) \neq 0$, then so is $\frac{f}{g}(x)$ and

$$\left(\frac{f}{g}\right)'(a) = \frac{f'(a)g(a) - f(a)g'(a)}{g(a)^2}.$$

Useful rules about derivatives (contd.)

The quotient rule

If $f(x)$ and $g(x)$ are differentiable at the point a and $g(a) \neq 0$, then so is $\frac{f}{g}(x)$ and

$$\left(\frac{f}{g}\right)'(a) = \frac{f'(a)g(a) - f(a)g'(a)}{g(a)^2}.$$

Composition : the chain rule

Useful rules about derivatives (contd.)

The quotient rule

If $f(x)$ and $g(x)$ are differentiable at the point a and $g(a) \neq 0$, then so is $\frac{f}{g}(x)$ and

$$\left(\frac{f}{g}\right)'(a) = \frac{f'(a)g(a) - f(a)g'(a)}{g(a)^2}.$$

Composition : the chain rule

If $f(x)$ and $g(x)$ are differentiable functions, then so is the function $f(g(x))$ and its derivative is :

$$(f(g))'(x) = f'(g(x))g'(x).$$

Examples

Examples

Guess: $f(x) = x^n$
 $f'(x) = nx^{n-1}$

$$f(x) = x^n; n \in \mathbb{N}$$

$$= x^{n-1} \times x$$

$$f'(x) = (n-1)x^{n-2} \times x + x^{n-1} \times 1$$
$$= (n-1)x^{n-1} + x^{n-1} = nx^{n-1}$$

$$f(x) = 5x^3 - 17x^2 + \pi x - 0.5$$

$$f'(x) = 5 \times 3x^2 - 17 \times 2x + \pi \times 1 - 0$$
$$= 15x^2 - 34x + \pi$$

$$f(x) = x^7 \sin(x)$$

$$f'(x) = 7x^6 \times \sin(x) + x^7 \times \cos(x)$$
$$= 7x^6 \sin(x) + x^7 \cos(x)$$

$$f(x) = x^a$$
$$f'(x) = a x^{a-1}$$

$$f(x) = x^2 = x \times x$$
$$f'(x) = (x)' \times x + x \times (x)'$$
$$= 1 \times x + x \times 1$$
$$= x + x = 2x$$

$$f(x) = x^3 = x^2 \times x$$
$$f'(x) = 2x \times x + x^2 \times 1$$
$$= 2x^2 + x^2$$
$$= 3x^2$$

Gen. statement for a polynomial.

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$$
$$\Rightarrow f'(x) = n a_n x^{n-1} + (n-1) a_{n-1} x^{n-2} + \dots + a_1$$

Examples

Examples

$$f(x) = \tan(x) = \frac{\sin(x)}{\cos(x)}$$

$$\begin{aligned} f'(x) &= \frac{\cos(x) \times \cos(x) - \sin(x) \times (-\sin(x))}{\cos^2(x)} \\ &= \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} = \frac{1}{\cos^2(x)} \\ &= \sec^2(x). \end{aligned}$$

$$f(x) = \frac{1}{x^r} \quad ; r > 0$$

$$= x^{-r}$$

$$\begin{aligned} f'(x) &= \frac{\frac{d}{dx}(1) \times x^r - 1 \times \frac{d}{dx}(x^r)}{(x^r)^2} \\ &= \frac{-r x^{r-1}}{x^{2r}} = -r x^{r-1-2r} \\ &= -r x^{-r-1} \end{aligned}$$

$$f(x) = \tan(2x)$$

$$= \frac{\sin(2x)}{\cos(2x)} = \frac{2 \tan(x)}{1 - \tan^2(x)}$$

$$\begin{aligned} g(x) &= 2x. & h(x) &= \tan(x) \\ f(x) &= h(g(x)). \\ f'(x) &= h'(g(x)) g'(x) \\ &= \sec^2(2x) \times 2 \\ &= 2 \sec^2(2x) \end{aligned}$$

Indeterminate limits

Indeterminate limits

Let $f(x)$ and $g(x)$ be functions and suppose $f(x)$ and $g(x)$ are defined on an interval around the point c .

Indeterminate limits

Let $f(x)$ and $g(x)$ be functions and suppose $f(x)$ and $g(x)$ are defined on an interval around the point c .

Further suppose $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or both limits diverge to ∞ or both limits diverge to $-\infty$.

Indeterminate limits

Let $f(x)$ and $g(x)$ be functions and suppose $f(x)$ and $g(x)$ are defined on an interval around the point c .

Further suppose $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or both limits diverge to ∞ or both limits diverge to $-\infty$.

Suppose we are interested in computing $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$.

Indeterminate limits

Let $f(x)$ and $g(x)$ be functions and suppose $f(x)$ and $g(x)$ are defined on an interval around the point c .

Further suppose $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or both limits diverge to ∞ or both limits diverge to $-\infty$.

Suppose we are interested in computing $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$.

Then we cannot use the quotient rule to compute it since the quotient rule yields an *indeterminate form*

Indeterminate limits

Let $f(x)$ and $g(x)$ be functions and suppose $f(x)$ and $g(x)$ are defined on an interval around the point c .

Further suppose $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or both limits diverge to ∞ or both limits diverge to $-\infty$.

Suppose we are interested in computing $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$.

Then we cannot use the quotient rule to compute it since the quotient rule yields an *indeterminate form* e.g. $\lim_{x \rightarrow 0} \frac{\log_e(1+x)}{x} =$

$$\lim_{x \rightarrow 0} \log_e(1+x) = \log_e(1+0) = \log_e(1) = 0.$$
$$\lim_{x \rightarrow 0} x = 0.$$

Indeterminate limits

Let $f(x)$ and $g(x)$ be functions and suppose $f(x)$ and $g(x)$ are defined on an interval around the point c .

Further suppose $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or both limits diverge to ∞ or both limits diverge to $-\infty$.

Suppose we are interested in computing $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$.

Then we cannot use the quotient rule to compute it since the quotient rule yields an *indeterminate form* e.g. $\lim_{x \rightarrow 0} \frac{\log_e(1+x)}{x} =$

In this situation, we can try and use **L'Hôpital's rule**.

Indeterminate limits : L'Hôpital's rule

Indeterminate limits : L'Hôpital's rule

In the situation of the indeterminate form, suppose the following conditions hold :

Indeterminate limits : L'Hôpital's rule

In the situation of the indeterminate form, suppose the following conditions hold :

1. $f'(x)$ and $g'(x)$ exist on this interval (except possibly at c).

Indeterminate limits : L'Hôpital's rule

In the situation of the indeterminate form, suppose the following conditions hold :

1. $f'(x)$ and $g'(x)$ exist on this interval (except possibly at c).
2. $g'(x) \neq 0$ on this interval (except possibly at c).

Indeterminate limits : L'Hôpital's rule

In the situation of the indeterminate form, suppose the following conditions hold :

1. $f'(x)$ and $g'(x)$ exist on this interval (except possibly at c).
2. $g'(x) \neq 0$ on this interval (except possibly at c).
3. $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = L$.

Indeterminate limits : L'Hôpital's rule

In the situation of the indeterminate form, suppose the following conditions hold :

1. $f'(x)$ and $g'(x)$ exist on this interval (except possibly at c).
2. $g'(x) \neq 0$ on this interval (except possibly at c).
3. $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = L$.

Then
$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = L.$$

Indeterminate limits : L'Hôpital's rule

In the situation of the indeterminate form, suppose the following conditions hold :

1. $f'(x)$ and $g'(x)$ exist on this interval (except possibly at c).
2. $g'(x) \neq 0$ on this interval (except possibly at c).
3. $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = L$.

Then $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = L$.

e.g. $\lim_{x \rightarrow 0} \frac{\log_e(1+x)}{x} = \lim_{x \rightarrow 0} \frac{\frac{1}{1+x}}{1} = \lim_{x \rightarrow 0} \frac{1}{1+x} = 1$.

$$\frac{d}{dx}(\log_e(1+x)) = \frac{1}{1+x}, \quad \frac{d}{dx}(x) = 1.$$

More examples

More examples

$$\lim_{x \rightarrow 2} \frac{x^2 - 5x + 6}{x - 2} = \lim_{x \rightarrow 2} \frac{2x - 5}{1} = \lim_{x \rightarrow 2} (2x - 5) = 2 \cdot 2 - 5 = -1.$$

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = \lim_{x \rightarrow 0} \frac{\cos(x)}{1} = \lim_{x \rightarrow 0} \cos(x) = \cos(0) = 1.$$

$$\lim_{x \rightarrow \infty} \frac{a + be^x}{c + de^x} = \lim_{x \rightarrow \infty} \frac{be^x}{de^x} = \lim_{x \rightarrow \infty} \frac{b}{d} = \frac{b}{d}.$$

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x^2} = \lim_{x \rightarrow 0} \frac{\sin(x)}{2x} = \frac{1}{2} \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = \frac{1}{2}.$$

Thank you