

Statistics for Data Science -1

Lecture 6.4: Probability- Properties of Probability

Usha Mohan

Indian Institute of Technology Madras

Learning objectives

Learning objectives

1. Understand uncertainty and concept of a random experiment.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.
6. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.
6. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
7. Distinguish between independent and dependent events.

Learning objectives

1. Understand uncertainty and concept of a random experiment.
2. Describe sample spaces, events of random experiments.
3. Understand the notion of simple event and compound events.
4. Basic laws of probability.
5. Calculate probabilities of events and use a tree diagram to compute probabilities.
6. Understand notion of conditional probability, i.e find the probability of an event given another event has occurred.
7. Distinguish between independent and dependent events.
8. Solve applications of probability.

Properties of Probability

Equally likely outcomes

The three main interpretations of probability

The three main interpretations of probability

1. Classical (Apriori or theoretical): Let S be the sample space of a random experiment in which there are n **equally likely** outcomes, and the event E consists of exactly m of these outcomes, then we say the probability of the event E is $\frac{m}{n}$ and represent it as $P(E) = \frac{m}{n}$

The three main interpretations of probability

1. Classical (Apriori or theoretical): Let S be the sample space of a random experiment in which there are n **equally likely** outcomes, and the event E consists of exactly m of these outcomes, then we say the probability of the event E is $\frac{m}{n}$ and represent it as $P(E) = \frac{m}{n}$
2. Relative frequency (Aposteriori or empirical): The probability of an event in an experiment is the proportion (or fraction) of times the event occurs in a very long (theoretically infinite) series of (independent) repetitions of experiment. In other words, if $n(E)$ is the number of times E occurs in n repetitions of the experiment, $P(E) = \lim_{n \rightarrow \infty} \frac{n(E)}{n}$

The three main interpretations of probability

1. Classical (Apriori or theoretical): Let S be the sample space of a random experiment in which there are n **equally likely** outcomes, and the event E consists of exactly m of these outcomes, then we say the probability of the event E is $\frac{m}{n}$ and represent it as $P(E) = \frac{m}{n}$
2. Relative frequency (Aposteriori or empirical): The probability of an event in an experiment is the proportion (or fraction) of times the event occurs in a very long (theoretically infinite) series of (independent) repetitions of experiment. In other words, if $n(E)$ is the number of times E occurs in n repetitions of the experiment, $P(E) = \lim_{n \rightarrow \infty} \frac{n(E)}{n}$
3. Subjective: The probability of an event is a “**best guess**” by a person making the statement of the chances that the event will happen. The probability measures an individual's degree of belief in the event.

Probability Axioms

Consider an experiment whose sample space is S . We suppose that for each event E there is a number, denoted $P(E)$ and called the probability of event E , that is in accord with the following three properties (axioms).

Probability Axioms

Consider an experiment whose sample space is S . We suppose that for each event E there is a number, denoted $P(E)$ and called the probability of event E , that is in accord with the following three properties (axioms).

1. For any event E , the probability of E is a number between 0 and 1. That is, $0 \leq P(E) \leq 1$.

Probability Axioms

Consider an experiment whose sample space is S . We suppose that for each event E there is a number, denoted $P(E)$ and called the probability of event E , that is in accord with the following three properties (axioms).

1. For any event E , the probability of E is a number between 0 and 1. That is, $0 \leq P(E) \leq 1$.
2. The probability of sample space S is 1. Symbolically, $P(S) = 1$.

Probability Axioms

Consider an experiment whose sample space is S . We suppose that for each event E there is a number, denoted $P(E)$ and called the probability of event E , that is in accord with the following three properties (axioms).

1. For any event E , the probability of E is a number between 0 and 1. That is, $0 \leq P(E) \leq 1$.
2. The probability of sample space S is 1. Symbolically, $P(S) = 1$. In other words, the outcome of the experiment will be an element of sample space S with probability 1.

Probability Axioms

Consider an experiment whose sample space is S . We suppose that for each event E there is a number, denoted $P(E)$ and called the probability of event E , that is in accord with the following three properties (axioms).

1. For any event E , the probability of E is a number between 0 and 1. That is, $0 \leq P(E) \leq 1$.
2. The probability of sample space S is 1. Symbolically, $P(S) = 1$. In other words, the outcome of the experiment will be an element of sample space S with probability 1.
3. For a sequence of mutually exclusive (disjoint) events, E_1, E_2, \dots ,

$$P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i)$$

Probability of union of disjoint events

The third property can be stated as:

Probability of union of disjoint events

The third property can be stated as:

The probability of the union of disjoint events is equal to the sum of the probabilities of these events.

Probability of union of disjoint events

The third property can be stated as:

The probability of the union of disjoint events is equal to the sum of the probabilities of these events.

For instance, if E_1 and E_2 are disjoint, then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2)$$

Probability of union of disjoint events

The third property can be stated as:

The probability of the union of disjoint events is equal to the sum of the probabilities of these events.

For instance, if E_1 and E_2 are disjoint, then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2)$$

In other words, if events E_1 and E_2 cannot simultaneously occur, then the probability that the outcome of the experiment is contained in either E_1 or E_2 is equal to the sum of the probability that it is in E_1 and the probability that it is in E_2 .

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$
 - ▶ E and E^c are disjoint. Also, $E \cup E^c = S$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$
 - ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
 - ▶ Apply Property 3 to LHS

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get
$$P(E \cup E^c) = P(E) + P(E^c) = P(S) = 1.$$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get
 $P(E \cup E^c) = P(E) + P(E^c) = P(S) = 1.$
Hence $P(E^c) = 1 - P(E)$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get
$$P(E \cup E^c) = P(E) + P(E^c) = P(S) = 1.$$

Hence $P(E^c) = 1 - P(E)$

2. $P(\Phi) = 0$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get
$$P(E \cup E^c) = P(E) + P(E^c) = P(S) = 1.$$

Hence $P(E^c) = 1 - P(E)$

2. $P(\Phi) = 0$

- ▶ $S^c = \Phi$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get
$$P(E \cup E^c) = P(E) + P(E^c) = P(S) = 1.$$

Hence $P(E^c) = 1 - P(E)$

2. $P(\Phi) = 0$

- ▶ $S^c = \Phi$
- ▶ Apply the above property, $P(S^c) = 1 - P(S)$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get
$$P(E \cup E^c) = P(E) + P(E^c) = P(S) = 1.$$

Hence $P(E^c) = 1 - P(E)$

2. $P(\Phi) = 0$

- ▶ $S^c = \Phi$
- ▶ Apply the above property, $P(S^c) = 1 - P(S)$
- ▶ Apply property 2, $P(S) = 1$

General properties of probability

Properties 1, 2, and 3 can be used to establish some general results concerning probabilities.

1. Probability of complement of an event: $P(E^c) = 1 - P(E)$

- ▶ E and E^c are disjoint. Also, $E \cup E^c = S$
- ▶ Apply Property 3 to LHS $P(E \cup E^c) = P(E) + P(E^c)$
- ▶ Apply Property 2 to RHS $P(S) = 1$
- ▶ Equating both, we get
$$P(E \cup E^c) = P(E) + P(E^c) = P(S) = 1.$$

Hence $P(E^c) = 1 - P(E)$

2. $P(\Phi) = 0$

- ▶ $S^c = \Phi$
- ▶ Apply the above property, $P(S^c) = 1 - P(S)$
- ▶ Apply property 2, $P(S) = 1$
- ▶ Hence $P(\Phi) = 0$


Addition rule of probability

The following formula relates the probability of the union of events E_1 and E_2 , which are not necessarily disjoint, to $P(E_1)$, $P(E_2)$, and the probability of the intersection of E_1 and E_2 . It is often called the addition rule of probability.

For any events E_1 and E_2 ,

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2)$$

Proof of addition rule



Section summary

- ▶ Probability axioms
- ▶ Properties of Probability
 - ▶ Addition rule